

KCNJ5 Mutations: Sex, Salt and Selection

Authors

T. A. Williams^{1,2}, J. W. M. Lenders^{3,4}, J. Burrello¹, F. Beuschlein², M. Reincke²

Affiliations

¹Division of Internal Medicine and Hypertension, Department of Medical Sciences, University of Turin, Turin, Italy

²Medizinische Klinik und Poliklinik IV, Klinikum der Ludwig-Maximilians-Universität München, Munich, Germany

³Department of Internal Medicine, Radboud University Medical Center, Nijmegen, The Netherlands

⁴Department of Internal Medicine III, Technische Universität Dresden, Dresden, Germany

Key words

- aldosterone-producing adenoma
- primary aldosteronism
- *KCNJ5*
- potassium channel

Abstract

Somatic mutations have been identified in the *KCNJ5* gene (encoding the potassium channel GIRK4) in aldosterone-producing adenomas (APA). Most of these mutations are located in or near the selectivity filter of the GIRK4 channel pore and

several have been shown to lead to the constitutive overproduction of aldosterone. *KCNJ5* mutations in APA are more frequent in women; however, this gender dimorphism is a reported phenomenon of Western but not East Asian populations. In this review we discuss some of the issues that could potentially underlie this observation.

Introduction

The original description of somatic aldosterone-producing adenoma (APA) mutations in the *KCNJ5* gene by Choi et al. in 2011 [1] marks one of the defining moments in primary aldosteronism (PA) research that initiated an extraordinary avalanche of studies in this field. Within 2 years of the discovery of *KCNJ5* mutations in APAs, the application of next-generation sequencing methods led to the identification of somatic APA mutations in 3 other genes. The affected genes were *ATP1A1* and *ATP2B3*, encoding Na⁺/K⁺-ATPase 1 and Ca²⁺-ATPase 3, respectively, and *CACNA1D* that encodes a subunit of an L-type voltage gated calcium channel, Cav1.3 [2–4]. However, in an intriguing development reported at the symposium *Progress in Primary Aldosteronism 4* held in June 2015 and published recently [5], somatic mutations in *ATP1A1*, *ATP2B3* and *CACNA1D* have also been identified in aldosterone-producing cell clusters (APCCs) in normal adrenal glands that stimulate *CYP11B2* activity and aldosterone production. In a large European collaboration, Fernandes-Rosa et al. [6] reported the combined prevalence of somatic mutations in APAs in *KCNJ5*, *ATP1A1*, *ATP2B3* and *CACNA1D* as 54% (○ Fig. 1). The functional effect of the thus far described somatic APA mutations all converge at the same intracellular level of the Ca²⁺ signalling pathway that ultimately lead to an increase in aldosterone biosynthesis [7–10]. The original identification of the somatic APA mutations in

KCNJ5 [1] and the combined efforts of several other groups highlighting the role of these mutations in unilateral PA in driving aldosterone excess is surely one of the major advances in hypertension research over the last decade.

G Protein-Activated Inwardly-Rectifying Potassium Channels

GIRK4 (encoded by *KCNJ5*, cytogenetic location: 11q24.3) is a member of the family of G protein-activated inwardly-rectifying potassium channels and is expressed at the plasma membrane in various different tissues, including the heart, central and peripheral neurons, various endocrine tissues, as well as in non-excitatory structures [11]. GIRK channels form transmembrane permeation pathways or pores with a high selectivity for K⁺ thereby preferentially allowing K⁺ to flow into the cell. However, the directional flow of K⁺ depends on where the ion channel is expressed. In adrenal zona glomerulosa cells, GIRK channels participate in maintaining the hyperpolarised state of the cell along with other K⁺ channels by allowing an outward flow of K⁺ conductance [12]. The K⁺ ion gradient across the cell membrane in GIRK channels is modulated in a voltage-dependent manner by the occlusion of the channel pore by intracellular polyamines and Mg²⁺ [13]. A GIRK channel subunit consists of 2 transmembrane helices with a large cytoplasmic domain and the N and C termini extruding into

received 19.08.2015

accepted 07.10.2015

Bibliography

DOI <http://dx.doi.org/10.1055/s-0035-1565090>

Published online:

November 13, 2015

Horm Metab Res 2015;

47: 953–958

© Georg Thieme Verlag KG

Stuttgart · New York

ISSN 0018-5043

Correspondence

T. A. Williams

Division of Internal Medicine

and Hypertension

Department of Medical

Sciences

University of Turin

Turin

Italy

Tel.: +39/011/6336 920

Fax: +39/011/6336 931

tracyannwilliams48@gmail.com





Fig. 1 Prevalence of somatic mutations in APA reported in a European cohort: The prevalence of somatic APA mutations in *KCNJ5*, *ATP1A1*, *ATP2B3* and *CACNA1D* reported in a European cohort of APA ($n=474$) recruited through the ENS@T registry (European Network for Adrenal Tumors, <http://www.ensat.org/>) [6].

the cytosol. An extracellular loop forms the selectivity filter that comprises the highly conserved signature sequence Gly-Tyr-Gly of K^+ channels. Functional GIRK channels are composed of 4 homo- or heterotetrameric subunits. Although GIRK4 can be expressed as a homotetramer, in most tissues and cell types it is found as a heterotetrameric complex with GIRK1, GIRK2 or GIRK3 [12].

The activation of GIRK channels is via direct interaction of the cytoplasmic domains with the G protein $\beta\gamma$ subunit and this process may also involve the anionic lipid phosphatidylinositol 4,5-bisphosphate [14]. In contrast, the inactivation of GIRK4 appears to be mediated by phospholipase C activation [15]. The crystal structure of GIRK4 has not been resolved, however, that of the related channel GIRK2 in the closed as well as a constitutively active conformation has been determined both in the absence and presence of phosphatidylinositol 4,5-bisphosphate providing insights into the molecular mechanisms of the multi-ligand regulation of GIRK channels [16].

Somatic *KCNJ5* Mutations in APA

Choi et al. [1] identified GIRK4-p.Gly151Arg and p.Leu168Arg substitutions in 8 out of 22 APAs. Both of these mutations interfere with the selectivity filter of the ion channel and lead to the indiscriminate conductance of Na^+ through the pore of the outer tunnel (see Fig. 2). The resultant membrane depolarisation causes the opening of Ca^{2+} voltage-gated ion channels, Ca^{2+} influx, activation of *CYP11B2* transcription and increased aldosterone biosynthesis. The ensuing renewed interest in PA research led to the rapid determination of the prevalence of somatic *KCNJ5* mutations in several cohorts of APAs [6, 17–29]. Foremost was the study by Boulkroun et al. [17] (see Table 1) that additionally reported the predominance of APA-*KCNJ5* mutations in women, an observation that was subsequently corroborated by others (see Table 1). Several novel mutations in *KCNJ5* associated with APAs were also identified (see Fig. 3), however, the mutations that were first described are by far the most prevalent [1]. A recent meta-analysis that incorporated 13 studies involving 1636 patients showed that female gender is a phenotype of APA patients carrying *KCNJ5* mutations along with pronounced aldosteronism, larger tumour size and younger age [30].



Mutation	Reference	Mutation	Reference
Gly151Arg	Choi 2011 [1]	Glu246Gly	Cheng 2015 [22]
Leu168Arg	Choi 2011 [1]	Ala139-Phe142dup	Xu 2015 [26]
delle157	Azizan 2012 [18]	Gly153-Gly164dup	Wang 2015 [28]
Glu145Gln	Akerström 2012 [19]	Thr148-Thr149insArg	Zheng 2015 [29]
Thr158Ala	Mulatero 2012 [20]	Ile157Lys	Scholl 2015 [53]
Trp126Arg	Williams 2014 [10]	Phe154Cys	Scholl 2015 [53]
insThr149	Kuppusamy 2014 [24]	Ile150_Gly151insMet	Scholl 2015 [53]
Arg115Trp	Cheng 2015 [22]	Ile144_Glu145insAlalle	Scholl 2015 [53]

Fig. 2 Cartoon of the pore module of GIRK4: The cartoon represents the pore module of a G protein inwardly rectifying K^+ channel. The first transmembrane segment M1 faces the lipid bilayer and the second transmembrane segment forms the inner pore helix. The K^+ permeation pathway comprises the narrow outer tunnel with the ion selectivity filter, the central cavity and the inner tunnel. Functional GIRK channels are composed of 4 homo- or heterotetrameric subunits and GIRK4 exists both as a homotetramer but more usually as a heterotetramer. The approximate positions of p.Gly151Arg in the selectivity filter of the outer tunnel and p.Leu168Arg in the inner helix are shown as green spheres. The somatic mutations identified to date with the reference that first reported them are shown in the panel below the figure. The figure was modified from a schematic diagram of the pore module of the *Streptomyces lividans* K^+ channel [54] according to the corresponding structure of the chicken *KCNJ12* channel [1], that shares 89% identity with human GIRK4 within the pore region and selectivity filter.

Reported Prevalence of *KCNJ5* Mutations in APA Patients in Different Populations

Unilateral PA patients from Western populations report a lower prevalence of somatic APA *KCNJ5* mutations compared to East Asian populations (see Table 1, 2, see Fig. 3, 39 vs. 73%, $p < 0.0001$). Bilateral adrenal hyperplasia (BAH or idiopathic aldosteronism) is generally reported to account for around 2-thirds of all cases of PA [31]. In contrast, a Japanese nationwide epidemiological study reported that the frequency of unilateral APA predominated over BAH accounting for 86% of cases of APA and 14% BAH patients ($n=1409$) [32]. It is likely that this indicates a selection bias for APA patients that display a more pronounced phenotype with higher blood pressure levels compared to BAH patients; in fact, in the study by Miyake et al. [32], the unilateral APA patients displayed higher blood pressure levels compared to the BAH patients. Further, in China, where accessibility to AVS centres is limited, a selection bias is likely to exist and favour patients with a more florid phenotype potentially meaning those with large adrenal nodules. Other factors influencing the selection bias for APA in Japanese patients could be the use of ACTH stimulation during adrenal venous sampling (AVS) for subtype diagnosis in all Japanese centres: this could theoretically increase the aldos-

terone production from the adenoma, thereby enhancing the probability of detecting an APA [33]. Also, APA carrying *KCNJ5* mutations are more frequently composed of zona fasciculata-like cells; in contrast to APA of “wild type” status, or of APA carrying *ATP1A1*, *ATP2B3* or *CACNA1D* mutations, that are comprised principally of zona glomerulosa-like cells [23,34,35]. Because

Table 1 Reported prevalence of *KCNJ5* mutations according to gender in Western populations.

Population	Sample size	<i>KCNJ5</i> mutations associated with unilateral PA	Reference
West Europe	380	129/380 (34%)	Boukroun 2012 [17]
Men-No. (%)	182/380 (48%)	32/182 (19%)	
Women-No. (%)	198/380 (52%)	97/198 (49%)	
West Europe-Australia	348	157/348 (47%)	Åkerström 2012 [19]
Men-No. (%)	151/348 (43%)	35/151 (23%)	
Women-No. (%)	197/348 (57%)	122/197 (62%)	
USA-Italy	42	13/42 (31%)	Monticone 2012 [9]
Men-No. (%)	18/42 (43%)	3/18 (17%)	
Women-No. (%)	24/42 (57%)	10/24 (42%)	
Norway	28	10/28 (36%)	Arnesen 2013 [21]
Men-No. (%)	20/28 (71%)	5/20 (25%)	
Women-No. (%)	8/28 (29%)	5/8 (63%)	
Italy	112[#]	44/112 (39%)	Williams 2014 [10]
Men-No. (%)	58/112 (52%)	17/58 (29%)	
Women-No. (%)	54/112 (48%)	27/54 (50%)	
The Netherlands	39[*]	22/39 (56%)	Dekkers 2014 [23]
Men-No. (%)	18/39 (46%)	7/18 (39%)	
Women-No. (%)	21/39 (54%)	15/21 (71%)	

* There were 53 samples in the original study but of these 39 samples had complete information for gender and genotype which are included here

[#]Thirty-three of these samples were included in the study of Boukroun [17] and were excluded from the subsequent analysis depicted in **Fig. 3**. These 33 samples comprised 16 women with 10 APA carrying *KCNJ5* mutations and 17 men with 5 APA carrying *KCNJ5* mutations

ACTH is the physiological regulator of hormone production in the zona fasciculata, whereas angiotensin II and K^+ mainly control hormone production in the zona glomerulosa, the use of ACTH stimulation during AVS in Japanese centres may also introduce a selection bias for APA carrying *KCNJ5* mutations. Therefore, these factors could contribute to the disproportionate selection of the APA subtype and potentially of *KCNJ5*-mutated APAs in Japanese populations compared to Western populations. In an interesting development, a recent meta-analysis demonstrated a correlation between mean daily urinary Na^+ excretion with the frequency of *KCNJ5* mutations in APA and hypothesised that higher average dietary Na^+ intake may cause a more severe phenotype and/or earlier detection of the disease in APA patients [30].

The INTERMAP population study illustrated the wide variations in salt consumption between East Asian (China and Japan) and Western (USA and UK) diets in which participants from East Asia on a self-reported reduced salt diet displayed higher daily urinary Na^+ excretion compared with Western participants on a non-reduced salt diet [36] (**Table 3**). Further, the Intersalt international study of electrolyte excretion and blood pressure from 52 centres, with a Na^+ excretion range of 0.2 mmol/24h (Yanomamo Indians, Brazil) to 242 mmol/24h (North China), a significant positive association between daily urinary Na^+ excretion and systolic blood pressure and between the Na^+/K^+ ratio and systolic blood pressure was found [37]. Four of the 52 centres were isolated populations that displayed very low sodium excretion, low sodium/potassium excretion, low body mass index, and low alcohol consumption. In these populations, high blood pressure was rare or non-existent [37,38].

In a global survey carried out in 2010, mean Na^+ intake was 3.95 g/day, twice the WHO recommended limit (2 g Na^+ /day). Intakes were highest in East Asia, Central Asia and Eastern Europe (mean > 4.2 g/day). In contrast, regional mean intakes in North America, Western Europe and Australia/New Zealand ranged from 3.4 to 3.8 g/day [39].




Fig. 3 Apparent prevalence of *KCNJ5* mutations according to gender and region: The Western populations group comprises Western European countries, Australia and the USA; the East Asia group comprises China and Japan. The pie charts represent the percentage of men or women with or without APA carrying *KCNJ5* mutations as a function of the total number of APA (men + women). The panel under the figure gives the percentage of men or women with APA carrying *KCNJ5* mutations as a function of the number of APA from the same gender as indicated. P-values were calculated using the Yates Chi-squared test and $p < 0.05$ was considered significant. *ns*: Not significant. The frequencies of *KCNJ5* mutations in APA in the men vs. men and women vs. women from the Western populations compared to the Eastern Asia groups were also significantly different ($p < 0.0001$).

Table 2 Reported prevalence of *KCNJ5* mutations according to gender in East Asian populations.

Population	Sample size	<i>KCNJ5</i> mutations associated with unilateral PA	Reference
Japan	5	5 (100%)	Monticone 2012 [9]
Men-No. (%)	2/5 (40%)	2/5 (100%)	
Women-No. (%)	3/5 (60%)	3/5 (100%)	
Japan	23	15/23 (65%)	Taguchi 2012 [25]
Men-No. (%)	10/23 (43%)	7/10 (70%)	
Women-No. (%)	13/23 (57%)	8/13 (62%)	
Japan	108*	75/108 (69%)	Kitamoto 2015 [27]
Men-No. (%)	37/103 (36%)	25/37 (68%)	
Women-No. (%)	66/103 (64%)	50/66 (76%)	
China	114	86/114 (75%)	Wang 2015 [28]
Men-No. (%)	52/114 (46%)	40/52 (77%)	
Women-No. (%)	62/114 (54%)	46/62 (74%)	
China	168	129/168 (77%)	Zheng 2015 [29]
Men-No. (%)	83/168 (49%)	56/83 (67%)	
Women-No. (%)	85/168 (51%)	73/85 (86%)	

* The original study comprised 108 patients and excluded 5 of them from their analysis due mutations in other genes (*ATP1A1*, *ATP2B3*, *CACNA1D*)

Table 3 Daily urinary Na⁺ excretion in all participants from the INTERMAP population study reporting non-reduced salt and reduced salt diets.

Nation	Non-RSD	RSD
UK	144.7 ± 2.0 (n=447)	155.2 ± 9.0 (n=24)
USA	163.6 ± 1.2 (n=2036)	148.9 ± 4.2 (n=159)
Japan	198.9 ± 1.5 (n=1109)	181.0 ± 8.4 (n=36)
China	228.3 ± 2.6 (n=828)	171.5 ± 22.5 (n=22.5)

Data were from the study by Okuda et al. [36]. Numbers refer to the 24-h urinary Na⁺ excretion in mmol (mean ± standard error). In parenthesis, n = number of participants. RSD: reduced salt diet; non-RSD, Non-reduced salt diet

According to the China national diabetes and metabolic disorders study, 26.6% of Chinese adults (20 years or older, approximately 254 million individuals), are hypertensive [40]. If dietary salt intake is correlated to a more severe phenotype and/or earlier detection of the disease in APA patients [30] then it is likely to have a more pronounced effect in East Asian patients. Finally, another factor that could influence the detection of PA and some of its subtypes in different populations, is the difference in blood pressure sensitivity to aldosterone among races [41] that could potentially affect phenotype, detection and referral in different ethnic groups. However, the relevance of this in East Asians compared to other ethnic groups is unknown.

Increased Prevalence of *KCNJ5* Mutations in APA in Western Women

KCNJ5 mutations in APA have consistently been reported as more frequent in women in Western populations (○ **Table 1**, ○ **Fig. 3**). ○ **Table 1** shows the prevalence of *KCNJ5* mutations in APA patients according to gender in cohorts from Europe, Australia and the USA (Western populations, ○ **Table 1**, ○ **Fig. 3**) with an overall predominance in women compared to men (55 vs. 22%, $p < 0.0001$), in agreement with a recent meta-analysis [30]. In contrast, this distinction was not evident in East Asia (China and Japan, ○ **Table 2**) where there was no significant difference between genders (women vs. men: 79 vs. 71%, ○ **Fig. 3**).

In a European population, in addition to being more prevalent in women, *KCNJ5* mutation status in APA patients was correlated with a younger age at diagnosis [6, 17]. It is tempting to speculate that the younger age at diagnosis may have been a gender effect, potentially related to the reluctance of men to seek medical help and failing to do so until the disease has progressed [42]. Consistent with this idea is a Danish study that demonstrated a lower contact rate of men with the general practitioner, and higher rates of hospitalisation and mortality [43]. It would be relevant to study this hypothesis by analysing the severity of hypertension, and associated target organ damage at study entry according to gender and genotype. Interestingly, in a study to assess trends in hypertension in US adults (NHANES 1988–1994 and 1999–2004), among the race/ethnic groups studied, women generally showed better awareness, treatment, and control rates than men [44].

The interrelation of blood pressure and sex steroids may also play a role in the gender dimorphism of *KCNJ5* mutation status in APA patients. Blood pressure levels are consistently reported to be higher in males compared to females [44–47]. This gender difference in blood pressure appears during adolescence and the relative prevalence of hypertension shifts to predominate in females in the elderly [48]. Interestingly, during the menopause transition, estradiol levels are 20% lower for Chinese and Japanese women compared to Caucasian, Hispanic or African-American women [49]. Estrogens are associated with protective cardiovascular effects [50] and 17 β -estradiol inhibits aldosterone synthesis in vitro via the estrogen β receptor [51]. The role of dietary Na⁺ excess in the pathogenesis of primary hypertension may have a more pronounced effect in men than in women [52]. Further, in a global survey, mean dietary Na⁺ intake in men was around 10% higher than in women [39].

Therefore, pre-menopausal Western women with PA may not be exposed to the deleterious effects of higher blood pressure and higher salt intake to the extent of their male counterparts. Further, they might benefit from the protective vascular effects of estrogen. It is attractive to hypothesise that these factors may contribute to shift the balance towards *KCNJ5* mutated APAs which are associated with a more severe phenotype and a higher likelihood of coming to medical attention.

Summary and Conclusions

Patients with APA harbouring *KCNJ5* mutations display a more severe form of aldosteronism and this may cause a selection bias in East Asian populations theoretically further exacerbated in Japanese centres by the use of ACTH stimulation during AVS. Against a background of lower blood pressure levels, lower salt intake and vascular protection by estrogens, pre-menopausal female APA patients from Western populations may present for medical examination with a more pronounced aldosteronism compared to their male counterparts. This could potentially contribute to the sexual dimorphism of APA-*KCNJ5* mutations in Western populations. In contrast, East Asian women have a higher dietary salt intake, higher blood pressure levels and lower estrogen levels than Western women.

To address any potential pathophysiological basis of these issues: 1) cohorts of APA-*KCNJ5* patients could be analysed according to clinical phenotype (blood pressure, biochemical and hormonal parameters) and gender; 2) the true prevalence of *KCNJ5* mutations could be analysed in a large (unbiased) autopsy study from

both Western and Asian populations within the same age range as the published cohort series; and 3) the mutational status of APA patients could be correlated with parameters of quality of life and psychological well-being in a multivariate adjusted analysis to study gender-dependent effects on hypertension presentation.

Acknowledgements

This work was funded by a grant from the Deutsche Forschungsgemeinschaft (RE 752/20-1 to M.R. and BE 2177/13-1 to F.B.) and by a grant of the Else Kröner-Fresenius Stiftung (2013_A182) to M.R.

Conflict of Interest

The authors declare no conflict of interest.

References

- Choi M, Scholl UI, Yue P, Björklund P, Zhao B, Nelson-Williams C, Ji W, Cho Y, Patel A, Men CJ, Lolis E, Wisgerhof MV, Geller DS, Mane S, Hellman P, Westin G, Åkerström G, Wang W, Carling T, Lifton RP. K+ channel mutations in adrenal aldosterone-producing adenomas and hereditary hypertension. *Science* 2011; 331: 768–772
- Beuschlein F, Boulkroun S, Osswald A, Wieland T, Nielsen HN, Lichtenauer UD, Penton D, Schack VR, Amar L, Fischer E, Walther A, Tauber P, Schwarzmayr T, Diener S, Graf E, Allolio B, Samson-Couterie B, Benecke A, Quinkler M, Fallo F, Plouin PF, Mantero F, Meitinger T, Mulatero P, Jeunemaitre X, Warth R, Vilsen B, Zennaro MC, Strom TM, Reincke M. Somatic mutations in ATP1A1 and ATP2B3 lead to aldosterone-producing adenomas and secondary hypertension. *Nat Genet* 2013; 45: 440–444
- Azizan EA, Poulsen H, Tuluc P, Zhou J, Clausen MV, Lieb A, Maniero C, Garg S, Bochukova EG, Zhao W, Shaikh LH, Brighton CA, Teo AE, Davenport AP, Dekkers T, Tops B, Kusters B, Ceral J, Yeo GS, Neogi SG, McFarlane I, Rosenfeld N, Marass F, Hadfield J, Margas W, Chaggar K, Solar M, Deinum J, Dolphin AC, Farooqi IS, Striessnig J, Nissen P, Brown MJ. Somatic mutations in ATP1A1 and CACNA1D underlie a common subtype of adrenal hypertension. *Nat Genet* 2013; 45: 1055–1060
- Scholl UI, Goh G, Stolting G, de Oliveira RC, Choi M, Overton JD, Fonseca AL, Korah R, Starker LF, Kunstman JW, Prasad ML, Hartung EA, Maura N, Benson MR, Brady T, Shapiro JR, Loring E, Nelson-Williams C, Libutti SK, Mane S, Hellman P, Westin G, Åkerström G, Björklund P, Carling T, Fahlke C, Hidalgo P, Lifton RP. Somatic and germline CACNA1D calcium channel mutations in aldosterone-producing adenomas and primary aldosteronism. *Nat Genet* 2013; 45: 1050–1054
- Nishimoto K, Tomlins SA, Kuick R, Cani AK, Giordano TJ, Hovelson DH, Liu CJ, Sanjanwala AR, Edwards MA, Gomez-Sanchez CE, Nanba K, Rainey WE. Aldosterone-stimulating somatic gene mutations are common in normal adrenal glands. *Proc Natl Acad Sci USA* 2015; 112: E4591–E4599
- Fernandes-Rosa FL, Williams TA, Riester A, Steichen O, Beuschlein F, Boulkroun S, Strom TM, Monticone S, Amar L, Meatchi T, Mantero F, Cicala MV, Quinkler M, Fallo F, Allolio B, Bernini G, Maccario M, Giacchetti G, Jeunemaitre X, Mulatero P, Reincke M, Zennaro MC. Genetic spectrum and clinical correlates of somatic mutations in aldosterone-producing adenoma. *Hypertension* 2014; 64: 354–361
- Oki K, Plonczynski MW, Luis Lam M, Gomez-Sanchez EP, Gomez-Sanchez CE. Potassium channel mutant KCNJ5 T158A expression in HAC-15 cells increases aldosterone synthesis. *Endocrinology* 2012; 153: 1774–1782
- Williams TA, Monticone S, Crudo V, Warth R, Veglio F, Mulatero P. Visinin-like 1 is upregulated in aldosterone-producing adenomas with KCNJ5 mutations and protects from calcium-induced apoptosis. *Hypertension* 2012; 59: 833–839
- Monticone S, Hattangady NG, Nishimoto K, Mantero F, Rubin B, Cicala MV, Pezzani R, Auchus RJ, Ghayee HK, Shibata H, Kurihara I, Williams TA, Giri JG, Bollag RJ, Edwards MA, Isales CM, Rainey WE. Effect of KCNJ5 mutations on gene expression in aldosterone-producing adenomas and adrenocortical cells. *J Clin Endocrinol Metab* 2012; 97: E1567–E1572
- Williams TA, Monticone S, Schack VR, Stindl J, Burrello J, Buffolo F, Annaratone L, Castellano I, Beuschlein F, Reincke M, Lucatello B, Ronconi V, Fallo F, Bernini G, Maccario M, Giacchetti G, Veglio F, Warth R, Vilsen B, Mulatero P. Somatic ATP1A1, ATP2B3, and KCNJ5 mutations in aldosterone-producing adenomas. *Hypertension* 2014; 63: 188–195
- Mulatero P, Monticone S, Rainey WE, Veglio F, Williams TA. Role of KCNJ5 in familial and sporadic primary aldosteronism. *Nat Rev Endocrinol* 2013; 9: 104–112
- Gomez-Sanchez CE, Oki K. Minireview: potassium channels and aldosterone dysregulation: is primary aldosteronism a potassium channelopathy? *Endocrinology* 2014; 155: 47–55
- Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, Kurachi Y. Inwardly rectifying potassium channels: their structure, function, and physiological roles. *Physiol Rev* 2010; 90: 291–366
- Sadja R, Smadja K, Alagem N, Reuveny E. Coupling Gbetagamma-dependent activation to channel opening via pore elements in inwardly rectifying potassium channels. *Neuron* 2001; 29: 669–680
- Mao J, Wang X, Chen F, Wang R, Rojas A, Shi Y, Piao H, Jiang C. Molecular basis for the inhibition of G protein-coupled inward rectifier K(+) channels by protein kinase C. *Congest Heart Fail* 2006; 12: 200–205
- Whorton MR, MacKinnon R. Crystal structure of the mammalian GIRK2 K+ channel and gating regulation by G proteins, PIP2, and sodium. *Cell* 2011; 147: 199–208
- Boulkroun S, Beuschlein F, Rossi GP, Golib-Dzib JF, Fischer E, Amar L, Mulatero P, Samson-Couterie B, Hahner S, Quinkler M, Fallo F, Letizia C, Allolio B, Ceolotto G, Cicala MV, Lang K, Lefebvre H, Lenzini L, Maniero C, Monticone S, Perrocheau M, Pilon C, Plouin PF, Rayes N, Seccia TM, Veglio F, Williams TA, Zinnamosca L, Mantero F, Benecke A, Jeunemaitre X, Reincke M, Zennaro MC. Prevalence, clinical, and molecular correlates of KCNJ5 mutations in primary aldosteronism. *Hypertension* 2012; 59: 592–598
- Azizan EA, Murthy M, Stowasser M, Gordon R, Kowalski B, Xu S, Brown MJ, O'Shaughnessy KM. Somatic mutations affecting the selectivity filter of KCNJ5 are frequent in 2 large unselected collections of adrenal aldosteronomas. *Hypertension* 2012; 59: 587–591
- Åkerström T, Crona J, Delgado Verdugo A, Starker LF, Cupisti K, Willenberg HS, Knoefel WT, Saeger W, Feller A, Ip J, Soon P, Anlauf M, Alesina PF, Schmid KW, Decaussin M, Levillain P, Wängberg B, Peix JL, Robinson B, Zedenius J, Bäckdahl M, Caramuta S, Iwen KA, Botling J, Stålbjerg P, Kraimps JL, Dralle H, Hellman P, Sidhu S, Westin G, Lehnert H, Walz MK, Åkerström G, Carling T, Choi M, Lifton RP, Björklund P. Comprehensive re-sequencing of adrenal aldosterone producing lesions reveal three somatic mutations near the KCNJ5 potassium channel selectivity filter. *PLoS One* 2012; 7: e41926
- Mulatero P, Tauber P, Zennaro MC, Monticone S, Lang K, Beuschlein F, Fischer E, Tizzani D, Pallauf A, Viola A, Amar L, Williams TA, Strom TM, Graf E, Bandulik S, Penton D, Plouin PF, Warth R, Allolio B, Jeunemaitre X, Veglio F, Reincke M. KCNJ5 mutations in European families with non-glucocorticoid remediable familial hyperaldosteronism. *Hypertension* 2012; 59: 235–240
- Arnesen T, Glomnes N, Strømøy S, Knappskog S, Heie A, Akslen LA, Grytaas M, Varhaug JE, Gimm O, Brauckhoff M. Outcome after surgery for primary hyperaldosteronism may depend on KCNJ5 tumor mutation status: a population-based study from Western Norway. *Langenbecks Arch Surg* 2013; 398: 869–874
- Cheng CJ, Sung CC, Wu ST, Lin YC, Sytwu HK, Huang CL, Lin SH. Novel KCNJ5 mutations in sporadic aldosterone-producing adenoma reduce Kir3.4 membrane abundance. *J Clin Endocrinol Metab* 2015; 100: E155–E163
- Dekkers T, ter Meer M, Lenders JW, Hermus AR, Schultze Kool L, Langenhuijsen JF, Nishimoto K, Ogishima T, Mukai K, Azizan EA, Tops B, Deinum J, Küsters B. Adrenal nodularity and somatic mutations in primary aldosteronism: one node is the culprit? *J Clin Endocrinol Metab* 2014; 99: E1341–E1351
- Kuppussamy M, Carocchia B, Stindl J, Bandulik S, Lenzini L, Gioco F, Fishman V, Zanotti G, Gomez-Sanchez C, Bader M, Warth R, Rossi GP. A novel KCNJ5-insT149 somatic mutation close to, but outside, the selectivity filter causes resistant hypertension by loss of selectivity for potassium. *J Clin Endocrinol Metab* 2014; 99: E1765–E1773
- Taguchi R, Yamada M, Nakajima Y, Satoh T, Hashimoto K, Shibusawa N, Ozawa A, Okada S, Rokutanda N, Takata D, Koibuchi Y, Horiguchi J, Oyama T, Takeyoshi I, Mori M. Expression and mutations of KCNJ5 mRNA in Japanese patients with aldosterone-producing adenomas. *J Clin Endocrinol Metab* 2012; 97: 1311–1319
- Xu S, Hardege I, Murthy M, Gordon R, Stowasser M, O'Shaughnessy K. A novel insertional somatic KCNJ5 mutation in an Australian patient with an aldosterone producing adenoma. *J Hypertens* 2015; 33: e120

- 27 Kitamoto T, Suematsu S, Matsuzawa Y, Saito J, Omura M, Nishikawa T. Comparison of cardiovascular complications in patients with and without KCNJ5 gene mutations harboring aldosterone-producing adenomas. *J Atheroscler Thromb* 2015; 22: 191–200
- 28 Wang B, Li X, Zhang X, Ma X, Chen L, Zhang Y, Lyu X, Tang Y, Huang Q, Gao Y, Fan Y, Ouyang J. Prevalence and characterization of somatic mutations in Chinese aldosterone-producing adenoma patients. *Medicine* 2015; 94: e708
- 29 Zheng FF, Zhu LM, Nie AF, Li XY, Lin JR, Zhang K, Chen J, Zhou WL, Shen ZJ, Zhu YC, Wang JG, Zhu DL, Gao PJ. Clinical characteristics of somatic mutations in Chinese aldosterone-producing adenoma. *Hypertension* 2015; 65: 622–628
- 30 Lenzi L, Rossitto G, Maiolino G, Letizia C, Funder JW, Rossi GP. A Meta-Analysis of Somatic KCNJ5 K⁺ channel mutations in 1636 patients with an aldosterone-producing adenoma. *J Clin Endocrinol Metab* 2015; 100: E1089–E1095
- 31 Schirpenbach C, Reincke M. Primary aldosteronism: current knowledge and controversies in Conn's syndrome. *Nat Clin Pract Endocrinol Metab* 2007; 3: 220–227
- 32 Miyake Y, Tanaka K, Nishikawa T, Naruse M, Takayanagi R, Sasano H, Takeda Y, Shibata H, Sone M, Satoh F, Yamada M, Ueshiba H, Katabami T, Iwasaki Y, Tanaka H, Tanahashi Y, Suzuki S, Hasegawa T, Katsumata N, Tajima T, Yanase T. Prognosis of primary aldosteronism in Japan: results from a nationwide epidemiological study. *Endocr J* 2014; 61: 35–40
- 33 Young WF, Stanson AW, Thompson GB, Grant CS, Farley DR, van Heerden JA. Role for adrenal venous sampling in primary aldosteronism. *Surgery* 2004; 136: 1227–1235
- 34 Azizan EA, Lam BY, Newhouse SJ, Zhou J, Kuc RE, Clarke J, Happerfield L, Marker A, Hoffman GJ, Brown MJ. Microarray, qPCR, and KCNJ5 sequencing of aldosterone-producing adenomas reveal differences in genotype and phenotype between zona glomerulosa- and zona fasciculata-like tumors. *J Clin Endocrinol Metab* 2012; 97: E819–E829
- 35 Monticone S, Castellano I, Versace K, Lucatello B, Veglio F, Gomez-Sanchez CE, Williams TA, Mulatero P. Immunohistochemical, genetic and clinical characterization of sporadic aldosterone-producing adenomas. *Mol Cell Endocrinol* 2015; 411: 146–154
- 36 Okuda N, Stamler J, Brown IJ, Ueshima H, Miura K, Okayama A, Saitoh S, Nakagawa H, Sakata K, Yoshita K, Zhao L, Elliott P, INTERMAP Research Group. Individual efforts to reduce salt intake in China, Japan, UK, USA: what did people achieve? The INTERMAP Population Study. *J Hypertens* 2014; 32: 2385–2392
- 37 Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *BMJ* 1988; 297: 319–328
- 38 Stamler J, Rose G, Elliott P, Dyer A, Marmot M, Kesteloot H, Stamler R. Findings of the International Cooperative INTERSALT Study. *Hypertension* 1991; 17: 19–15
- 39 Powles J, Fahimi S, Micha R, Khatibzadeh S, Shi P, Ezzati M, Engell RE, Lim SS, Danaei G, Mozaffarian D, Global Burden of Diseases Nutrition and Chronic Diseases Expert Group (NutriCoDE). Global, regional and national sodium intakes in 1990 and 2010: a systematic analysis of 24 h urinary sodium excretion and dietary surveys worldwide. *BMJ Open* 2013; 3: e003733
- 40 Gao Y, Chen G, Tian H, Lin L, Lu J, Weng J, Jia W, Ji L, Xiao J, Zhou Z, Ran X, Ren Y, Chen T, Yang W, and for the China National Diabetes and Metabolic Disorders Study Group. Prevalence of Hypertension in China: A Cross-Sectional Study. *PLoS One* 2013; 8: e65938
- 41 Tu W, Eckert GJ, Hannon TS, Liu H, Pratt LM, Wagner MA, Dimeglio LA, Jung J, Pratt JH. Racial differences in sensitivity of blood pressure to aldosterone. *Hypertension* 2014; 63: 1212–1218
- 42 Banks I. No man's land: men, illness, and the NHS. *BMJ* 2001; 323: 1058–1060
- 43 Juel K, Christensen K. Are men seeking medical advice too late? Contacts to general practitioners and hospital admissions in Denmark 2005. *J Public Health* 2008; 30: 111–113
- 44 Cutler JA, Sorlie PD, Wolz M, Thom T, Fields LE, Roccella EJ. Trends in hypertension prevalence, awareness, treatment, and control rates in United States adults between 1988–1994 and 1999–2004. *Hypertension* 2008; 52: 818–827
- 45 Os I, Oparil S, Gerds E, Høiegggen A. Essential hypertension in women. *Blood Press* 2004; 13: 272–278
- 46 Burt VL, Whelton P, Roccella EJ, Brown C, Cutler JA, Higgins M, Horan MJ, Labarthe D. Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988–1991. *Hypertension* 1995; 25: 305–313
- 47 Sandberg K, Ji H. Sex differences in primary hypertension. *Biol Sex Differ* 2012; 3: 7
- 48 Kotchen JM, McKean HE, Kotchen TA. Blood pressure trends with aging. *Hypertension* 1982; 4: III128–III134
- 49 Baber RJ. East is east and West is west: perspectives on the menopause in Asia and The West. *Climacteric* 2014; 17: 23–28
- 50 Rossi P, Francès Y, Kingwell BA, Ahimastos AA. Gender differences in artery wall biomechanical properties throughout life. *J Hypertens* 2011; 29: 1023–1033
- 51 Caroccia B, Seccia TM, Campos AG, Gioco F, Kuppasamy M, Ceolotto G, Guerzoni E, Simonato F, Mareso S, Lenzi L, Fassina A, Rossi GP. GPER-1 and estrogen receptor- β ligands modulate aldosterone synthesis. *Endocrinology* 2014; 155: 4296–4304
- 52 Hedayat SS, Minhajuddin AT, Ijaz A, Moe OW, Elsayed EF, Reilly RF, Huang CL. Association of urinary sodium/potassium ratio with blood pressure: sex and racial differences. *Clin J Am Soc Nephrol* 2012; 7: 315–322
- 53 Scholl UI, Healy JM, Thiel A, Fonseca AL, Brown TC, Kunstman JW, Horne MJ, Dietrich D, Riemer J, Kücüköylü S, Reimer EN, Reis AC, Goh G, Kristiansen G, Mahajan A, Korah R, Lifton RP, Prasad ML, Carling T. Novel somatic mutations in primary hyperaldosteronism are related to the clinical, radiological and pathological phenotype. *Clin Endocrinol* 2015, Aug 7 doi: doi: 10.1111/cen.12873 [Epub ahead of print]
- 54 Sepúlveda FV, Cid LP, Teulon J, Niemeyer MI. Molecular aspects of structure, gating, and physiology of pH-sensitive background K2P and Kir K⁺-transport channels. *Physiol Rev* 2015; 95: 179–217