


Iodine as a New Catalyst for the Condensation of 2-Hydroxy-2,2'-bisindan-1,1',3,3'-tetrone with Cyclic Enaminones: Synthesis of Spiro-dihydropyridine Derivatives under Acid-Free Conditions

Firouz Matloubi Moghaddam*

Vahid Saberi

Seyedmahmoudreza Keshavarz

Laboratory of Organic Synthesis and Natural Products,
Department of Chemistry, Sharif University of Technology,
Azadi Street, PO Box 11159516 Tehran, Iran
matloubi@sharif.edu



Received: 05.11.2017

Accepted after revision: 16.01.2018

Published online: 01.02.2018

DOI: 10.1055/s-0036-1591918; Art ID: so-2017-d0055-1

License terms:

Abstract Iodine has been used as a new catalyst for the synthesis of spiro-dihydropyridine derivatives by the condensation of cyclic enaminones and 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone.

Key words iodine, aminouracil, spiro-dihydropyridine

The properties of molecular iodine, including its Lewis acidity, lead to a range of applications in organic synthesis.¹⁻⁴ Nitrogen-containing heterocyclic compounds have been of interest for the development of organic synthesis through decades.⁵⁻⁸ Among them, 1,4-dihydropyridines (1,4-DHPs; Figure 1) are an important class of compounds in the field of vasodilation and bronchodilation, being potent calcium antagonists and calcium channel blockers.⁹⁻¹²




Figure 1 Selected examples of dihydropyridines


Spiro-heterocycles, due to their steric constraints, represent an important class of substances that often have interesting biological properties.¹³⁻¹⁵ Among them, spiro-dihydropyridine derivatives serve as important building

blocks in a wide range of biologically active compounds.¹⁶ There are several synthetic methods available for the preparation of functionalized dihydropyridines.¹⁷

2-Hydroxy-2,2'-bisindan-1,1',3,3'-tetrone can be readily generated by acid or base catalyzed condensation of ninhydrin with 1,3-indanedione.¹⁸ Activation of 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone for electrophilic reactions has been achieved under acidic conditions such as AcOH/H₂SO₄ for addition of phenols,¹⁹ AcOH for enol condensation,²⁰ triflic acid,²¹ acidic magnetic nanoparticle for synthesis of pyrazoles,²² and silica-sulfuric acid for synthesis of dihydropyridines.²³ However, the use of acidic conditions can cause rearrangements and can also require high temperatures in some cases.

In the present work, in a continuation of our ongoing research program in the field of synthesis of spiro-heterocyclic compounds, we report the reaction of 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone **1** and cyclic enaminones **2a-j** in the presence of molecular iodine under mild reaction conditions to form spiro-dihydropyridines **3a-j** (Scheme 1).

Firstly, 1,3-indandione was reacted with ninhydrin in the presence of triethylamine in EtOH at room temperature, to afford 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone **1**. The reaction of compound **1** was tested initially with 6-aminouracil in dimethyl sulfoxide (DMSO) at 120 °C without catalyst. However, these conditions did not afford the desired product **3a** (Table 1, entry 2). The use of an acidic catalyst such as *p*-TSA or acetic acid gave **3a** in moderate yields (entries 3 and 4). Our group has previously used molecular iodine as a catalyst^{1,24} and when we examined the use of iodine in DMSO at 120 °C the reaction proceeded in excellent yield (entry 5). To study the effect of temperature, an additional experiment was performed at 100 °C, resulting in 91% yield of **3a** after 4 hours (entry 8). However, when the



Scheme 1 Synthesis spiro-dihydropyridines


reaction was performed at lower temperatures, the yield decreased dramatically (entries 9 and 10). When solvents such as *N,N*-dimethylformamide (DMF) and toluene were examined, again a decrease in the yield of the reaction was observed (entries 11 and 12). Therefore, 20 mol% I_2 in DMSO at 100 °C was established as the optimal conditions for the current methodology.

To investigate the scope of this reaction, a range of cyclic enaminones was used; the results are summarized in Table 2. All derivatives **3a–j** were obtained in high yields, although in the case of compounds **3i** and **3j** the yield of the reaction decreased slightly, presumably because the nitrogen of the 6-aminocoumarin is further substituted and therefore more sterically hindered.

All novel compounds **3a–f** were fully characterized by elemental analysis, IR, 1H NMR, ^{13}C NMR spectroscopic analysis and HRMS. For example, the HRMS spectrum of **3a** displayed m/z $[M+H]^+$ at 398.0775 and the IR spectrum showed absorption bands at 3249, 1710, 1690, and 1666 cm^{-1} assigned to NH stretching and five-membered ketone, amide, and α,β -unsaturated ketone carbonyl groups, respectively. The 1H -decoupled ^{13}C NMR spectrum of **3a** showed 18 distinct resonances, with a signal at 56 ppm being attributed to the spiro- carbon.

Finally, to investigate the scalability of our reaction, the reaction was conducted on a gram scale under the optimized reaction conditions and no significant changes in either reaction time or yield were observed (Scheme 2).


Table 1 Optimization of Reaction Conditions for the Preparation of **3a**^a



Entry	Solvent	Catalyst	Catalyst (%)	Temp. (°C)	Time (h)	Yield (%) ^b
1	DMSO	none	–	100	24	–
2	DMSO	none	–	120	24	–
3	DMSO	<i>p</i> -TSA	100	120	12	42
4	DMSO	HOAc	100	120	12	37
5	DMSO	I_2	100	120	12	94
6	DMSO	I_2	20	120	12	92
7	DMSO	I_2	10	120	12	65
8	DMSO	I_2	20	100	4	91
9	DMSO	I_2	20	80	4	44
10	DMSO	I_2	20	50	4	23
11	DMF	I_2	20	100	4	67
12	toluene	I_2	20	100	4	35

^a Reagents and conditions: **1** (1 mmol), **2a** (1 mmol), solvent (4 mL), in a capped vial.

^b Isolated yield.

Table 2 Synthesis of 3a–j^a


Entry	Cyclic Enaminones	Product	Yield (%) ^b
1	2a 	3a 	91
2	2b 	3b 	88
3	2c 	3c 	85
5	2d 	3d 	92
5	2e 	3e 	86
6	2f 	3f 	80

Table 2 (continued)

Entry	Cyclic Enaminones	Product	Yield (%) ^b
7	2g 	3g 	73
8	2h 	3h 	83
9	2i 	3i 	61
10	2j 	3j 	57

^a Reagents and conditions: **1** (1 mmol), **2a-j** (1 mmol), DMSO (4 mL), I₂ (20 mol%) in a capped vial.


^b Isolated yield.



Scheme 2

Although we have not established the mechanism of the reaction experimentally, a plausible mechanism is proposed in Scheme 3. The reaction between molecular iodine and hydroxy compound **1** could produce two intermediates, Path A gives intermediate **4**, which is reported to be formed under acidic conditions,^{19,21} and Path B leads to intermediate **5**. Michael addition of enaminone **2a** to either the double bond of **4** or the carbocationic center of **5** gives intermediate **7**, which undergoes intramolecular cyclocondensation to produce the desired product.

In conclusion, we have developed an efficient methodology for the synthesis of spiro-dihydropyridines in high yield from 2-hydroxy-2,2'-biindan-1,1',3,3'-tetrone as starting material.^{25,26} Activation of 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone with molecular iodine as a mild catalyst under acid-free reaction conditions and a simple work-up procedure are features of this method.



Scheme 3 Proposed mechanism of formation of spiro-dihydropyridans **3**

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0036-1591918>.

References

- Moghaddam, F. M.; Khodabakhshi, M. R.; Aminae, M. *Tetrahedron Lett.* **2014**, *55*, 4720.
- Alizadeh, A.; Saberi, V.; Mokhtari, J. *Synlett* **2013**, *24*, 1825.
- Zhang, J.; Gao, Q.; Wu, X.; Geng, X.; Wu, Y. D.; Wu, A. *Org. Lett.* **2016**, *18*, 1686.
- Hao, W. J.; Wang, S. Y.; Ji, S. J. *ACS Catal.* **2013**, *3*, 2501.
- Deiters, A.; Martin, S. F. *Chem. Rev.* **2004**, *104*, 2199.
- Thansandote, P.; Lautens, M. *Chem. Eur. J.* **2009**, *15*, 5874.
- Zhu, L.; Cheng, L.; Zhang, Y.; Xie, R.; You, J. *J. Org. Chem.* **2007**, *72*, 2737.
- Martín, R.; Rodríguez, Rivero. M.; Buchwald, S. L. *Angew. Chem. Int. Ed.* **2006**, *45*, 7079.
- (a) Safak, C.; Simsek, R. *Mini-Rev. Med. Chem.* **2006**, *6*, 747. (b) Katoh, M.; Nakajima, M.; Shimada, N.; Yamazaki, H.; Yokoi, T. *Eur. J. Clin. Pharmacol.* **2000**, *55*, 843. (c) Ruggenenti, P.; Perna, A.; Benini, R.; Remuzzi, G. *J. Am. Soc. Nephrol.* **1998**, *9*, 2096.
- (a) Glossmann, H.; Ferry, D. R.; Goll, A.; Striessnig, J.; Zernig, G. *Arzneim.-Forsch.* **1985**, *35*, 1917. (b) Matowe, W. C.; Akula, M.; Knaus, E. E.; Wolowyk, M. W. *Proc. West. Pharmacol. Soc.* **1989**, *32*, 305.
- Vo, D.; Matowe, W. C.; Ramesh, M.; Iqbal, N.; Wolowyk, M. W.; Howlett, S. E.; Knaus, E. E. *J. Med. Chem.* **1995**, *38*, 2851.
- (a) Vo, D.; Nguyen, J. T.; McEwen, C.-A.; Shan, R.; Knaus, E. E. *Drug Dev. Res.* **2002**, *56*, 1. (b) Edraki, N.; Mehdipour, A. R.; Khoshneviszadeh, M.; Miri, R. *Drug Discovery Today* **2009**, *14*, 1058.
- Zhang, Y. L.; Li, Y. F.; Wang, J. W.; Yu, B.; Shi, Y. K.; Liu, H. M. *Steroids* **2016**, *109*, 22.
- Parameswarappa, S. G.; Pigge, F. C. *J. Org. Chem.* **2012**, *77*, 8038.
- Tejedor, D.; Cotos, L.; Méndez-Abt, G.; García-Tellado, F. *J. Org. Chem.* **2014**, *79*, 10655.
- Auria-Luna, F.; Marqués-López, E.; Mohammadi, S.; Heiran, R.; Herrera, R. P. *Molecules* **2015**, *20*, 15807.
- (a) Senczyszyn, J.; Brice, H.; Clayden, J. *Org. Lett.* **2013**, *15*, 1922. (b) Debnath, K.; Singha, K.; Pramanik, A. *RSC Adv.* **2015**, *5*, 31866. (c) Sarkar, P.; Mukhopadhyay, C. *Tetrahedron Lett.* **2016**, *57*, 4306.
- (a) Campagna, F.; Carotti, A.; Casini, G.; Ferappi, M. *Gazz. Chim. Ital.* **1983**, *113*, 507. (b) Schoenberg, A.; Singer, E. *Chem. Ber.* **1970**, *103*, 3871.
- Das, S.; Pramanik, A.; Fröhlich, R.; Patra, A. *Tetrahedron* **2004**, *60*, 10197.
- Das, S.; Fröhlich, R.; Pramanik, R. *J. Chem. Res.* **2005**, *9*, 572.
- Das, S.; Fröhlich, R.; Pramanik, R. *J. Chem. Res.* **2007**, *1*, 5.
- Ashis, K.; Mukherjee, S.; Pramanik, R. *RSC Adv.* **2015**, *130*, 107847.
- Ashis, K. R.; Pramanik, R. *Mol. Diversity* **2015**, *3*, 459.
- (a) Moghaddam, F. M.; Bardajee, R. G.; Ismaili, H.; Dokht, M.; Taimoory, S. *Synth. Commun.* **2006**, *36*, 2543. (b) Moghaddam, F. M.; Khodabakhshi, M. R.; Aminae, M. *Tetrahedron Lett.* **2014**, *55*, 4720.
- Typical procedure for the synthesis of spiro-dihydropyridines 3a–j:** A mixture of 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone **1** (1 mmol), cyclic enaminone **2a–f** (1 mmol) and molecular iodine (20 mol%) in DMSO (4 mL) was stirred at 100 °C for 4 h. After completion of the reaction (monitored by TLC, ethyl acetate/*n*-hexane, 1:2) the reaction mixture was allowed to cool to room temperature. Water was added and the precipitate was filtered off and washed with acetone to give the product **3a–j**.
- Synthesis of 2-hydroxy-2,2'-bisindan-1,1',3,3'-tetrone 1:** A mixture of 1,3-indandione (10 mmol), ninhydrin (10 mmol), and triethylamine (1 mmol) in EtOH (50 mL) was stirred at room temperature for 5 h. The precipitate was filtered and washed with EtOH (2 × 5 mL) to give **1**. Yellow powder; mp 187–190 °C; ¹H NMR (500 MHz, CDCl₃): δ = 8.00–7.93 (m, 2 H, Ar), 7.92–7.83 (m, 6 H, Ar), 5.47 (s, 1 H, OH), 3.96 (s, 1 H, CH); ¹³C NMR (125 MHz, CDCl₃): δ = 197, 196, 142, 141, 137, 136, 124, 124, 76, 53.

Analytical data for spiro[indene-2,5'-indeno[2',1':5,6]pyrido[2,3-d]pyrimidine]-1,2',3,4',6'(1H,3H,11H)-pentaone (3a): Yield: 0.361 g (91%); red powder; mp 212–215 °C (dec.); IR (KBr): 3249, 2917, 1710, 1690, 1666 cm⁻¹; ¹H NMR (500 MHz, D₂O): δ = 7.89–7.85 (m, 4 H, Ar-H), 7.36 (d, *J* = 6.9 Hz, 1 H, Ar-H), 7.24–7.17 (m, 2 H, Ar-H), 7.00 (d, *J* = 6.9 Hz, 1 H, Ar-H); ¹³C NMR (125 MHz, D₂O): δ = 209, 187, 173, 170, 165, 162, 141, 138, 137, 136, 131, 130, 122, 119, 119, 100, 92, 56; Anal. Calcd for C₂₂H₁₁N₃O₅: C, 66.50; H, 2.79; N, 10.58. Found: C, 65.90; H, 2.81; N, 10.61; HRMS: *m/z* [M+H]⁺ calcd. for C₂₂H₁₁N₃O₅: 398.0777; found: 398.0775.