

# Intralesional Endoscopy and Septectomy as a Diagnostic Tool and Treatment Method for Lymphatic Malformations

Anne-Sophie Holler<sup>1</sup> Jan Gödeke<sup>1</sup> Veronika Engel<sup>1</sup> Oliver J. Muensterer<sup>1</sup>

<sup>1</sup>Department of Pediatric Surgery, University Medicine of the Johannes Gutenberg University Mainz, Mainz, Germany

Eur J Pediatr Surg Rep 2017;5:e60–e61.

**Address for correspondence** Anne-Sophie Holler, MD, Department of Pediatric Surgery, University Medicine of the Johannes Gutenberg University Mainz, Langenbeckstr. 1, Mainz 55131, Germany (e-mail: anne-sophie.holler@unimedizin-mainz.de).

## New Insights and the Importance for the Pediatric Surgeon


Current treatment options for lymphatic malformations are expectant management, sclerotherapy, and surgical resection. Intralesional endoscopy and intercystic septectomy constitute a promising supplementary treatment method for subcutaneous lymphatic malformations. By creating a single, communicating cavity, the efficacy of subsequent sclerotherapy may be increased, and thus the need for multiple treatment sessions may be avoidable.

## Case Report


Sclerotherapy and surgery are both effective treatment methods for lymphatic malformations.<sup>1,2</sup> However, recurrence due to incomplete resection is a common problem, often necessitating multiple treatment sessions.<sup>2</sup> Intralesional

endoscopy has been described as a diagnostic approach and potential therapeutic tool.<sup>1,3</sup>

We report a case of a 12-year-old male patient who presented with a mainly subcutaneous mixed lymphatic malformation located on the right flank (→Fig. 1). After suffering a direct trauma, the lesion had increased markedly in size and surgical intervention was indicated due to the associated pain. Intralesional endoscopy was performed that showed a mixed macro-/microcystic lymphatic malformation with hemorrhage (→Fig. 2). Intercystic septa were dissected under endo-



**Fig. 1** Coronal T2 magnetic resonance imaging (MRI) with a subcutaneous mixed macro-/microcystic lymphatic malformation on the patient's right flank (white arrow).



**Fig. 2** Intraoperative image showing a macrocystic cavity and many small fluid-filled cysts (asterisks), which are separated by thin septae.

received  
January 3, 2017  
accepted after revision  
February 23, 2017

DOI <https://doi.org/10.1055/s-0037-1606388>.  
ISSN 2194-7619.

© 2017 Georg Thieme Verlag KG  
Stuttgart · New York

License terms



scopic visualization. At the end of the procedure, a single macrocystic cavity had been artificially created. Picibanil (OK-432) was inserted into the cavity and left in situ for 24 hours (► **Video 1**). The patient had no visible swelling, no pain, and merely two small, well-healed scars at 2 months of follow-up.

### Video 1

Under sonographic guidance, the largest cyst of the lymphatic malformation was cannulated, and an 8 French pigtail catheter was advanced into the cyst over a guidewire. The cyst was filled with 0.9% sodium chloride solution to facilitate the placement of two 3-mm trocars into the cyst. Intralesional endoscopy showed a mixed macro-/microcystic lymphatic malformation with hemorrhage. Dissection of intercystic septa was performed under endoscopic visualization using hook electrocautery and blunt dissection. At the end of the procedure, a single macrocystic cavity had been artificially created. Picibanil (OK-432) was instilled through the formerly inserted pigtail catheter into the now solitary cyst and left in situ for 24 hours. Online content including video sequences viewable at: [www.thieme-connect.com/products/ejournals/html/10.1055/s-0037-1606388](http://www.thieme-connect.com/products/ejournals/html/10.1055/s-0037-1606388).

Intralesional endoscopy and intercystic septectomy constitute an interesting novel approach for the diagnosis and treatment of mixed lymphatic malformations. By creating a single, communicating cavity, the efficacy of subsequent sclerotherapy may be increased, and thus the need for multiple treatment sessions may be avoidable.

### Conflict of Interest

None.

### References

- 1 Defnet AM, Bagrodia N, Hernandez SL, Gwilliam N, Kandel JJ. Pediatric lymphatic malformations: evolving understanding and therapeutic options. *Pediatr Surg Int* 2016;32(05):425–433
- 2 Cheng J. Doxycycline sclerotherapy in children with head and neck lymphatic malformations. *J Pediatr Surg* 2015;50(12):2143–2146
- 3 Eivazi B, Teymoortash A, Wiegand S, et al. Intralesional endoscopy of advanced lymphatic malformations of the head and neck: a new diagnostic approach and a potential therapeutic tool. *Arch Otolaryngol Head Neck Surg* 2010;136(08):790–795