

Aneurysmal Subarachnoid Hemorrhage: Strategies for Preventing Vasospasm in the Intensive Care Unit

Michael N. Diring, MD¹ Allyson R. Zazulia, MD¹

¹Department of Neurology, Washington University School of Medicine, St. Louis, Missouri

Semin Respir Crit Care Med 2017;38:760–767.

Address for correspondence Michael N. Diring, MD, Department of Neurology, Washington University School of Medicine, Campus Box 8111, 660 S Euclid Avenue, St Louis, MO 63110 (e-mail: diringerm@neuro.wustl.edu).

Abstract

This article addresses the intensive care unit (ICU) management of patients with aneurysmal subarachnoid hemorrhage (SAH), with an emphasis on the prevention of cerebral vasospasm and delayed cerebral ischemia (DCI), which are major contributors to morbidity and mortality. Interventions addressing various steps in the development of vasospasm have been attempted, with variable success. Enteral nimodipine remains the only approved measure to potentially prevent DCI. Since oral and intravenous administrations are limited by hypotension, direct administration via sustained-release pellets and intraventricular administration of sustained-release microparticles are being investigated. Studies of other calcium channel blockers have been disappointing. Efforts to remove blood from the subarachnoid space via cisternal irrigation, cisternal or ventricular thrombolysis, and lumbar cerebrospinal fluid drainage have met with limited and variable success, and they remain an area of active investigation. Several interventions that had early promise have failed to show benefit when studied in large trials; these include tirilazad, magnesium, statins, clazosentan, transluminal angioplasty, and hypervolemia.

Keywords

- ▶ subarachnoid hemorrhage
- ▶ aneurysm
- ▶ vasospasm
- ▶ hypertension
- ▶ treatment

Subarachnoid hemorrhage (SAH) refers to bleeding that occurs primarily within the subarachnoid space. The most common cause of SAH is trauma, but among cases of spontaneous SAH, rupture of an intracranial saccular aneurysm accounts for approximately 85%.¹ Other causes of spontaneous SAH include bleeding from arteriovenous malformations, moyamoya syndrome, bleeding disorders, cocaine and stimulant abuse, and extension of intracerebral hematomas. In up to one-fifth of cases, no source of bleeding is identified.²

Acute aneurysmal SAH results from the sudden extrusion of blood from a defect in the wall of an arterial aneurysm. The duration of the bleeding event determines the impact of the primary injury, and ranges from a severe headache to syncope to sudden death in about a quarter of patients.³ The amount of blood in the subarachnoid space sets the stage for secondary injury from hydrocephalus,⁴ vasospasm, and

delayed cerebral ischemia (DCI).^{5,6} The focus of treatment is anticipating, preventing, and managing these secondary complications.

In addition to the neurologic consequences of SAH, intense activation of the sympathetic nervous system occurs, which can produce a variety of extracranial consequences including hypertension, stunned myocardium,⁷ neurogenic pulmonary edema,⁸ abnormal renal function, and systemic inflammatory response syndrome (SIRS).⁹

Risk Factors

Several risk factors for SAH have been identified, although it is not clear whether these factors identify risk of developing an aneurysm or risk of rupture of an aneurysm that already exists. Modifiable risk factors include hypertension, cigarette

Issue Theme Advancements in Neurocritical Care and Emergency Neurology; Guest Editors: David Y. Hwang, MD, FNCS, and David M. Greer, MD, MA, FCCM, FAHA, FNCS, FAAN, FANA

Copyright © 2017 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662.

DOI <https://doi.org/10.1055/s-0037-1607990>. ISSN 1069-3424.

smoking,¹⁰ and heavy alcohol consumption.¹¹ Genetic factors also play an important role in SAH.¹² The odds ratio of SAH for individuals with one affected first-degree relative is 2; this increases to 51 for individuals with two affected first-degree relatives.¹³ Aneurysm development is more common in patients with connective tissue disease and polycystic kidney disease, and identifying other genetic markers is an active area of investigation.^{14–17}

Pathophysiology

Most aneurysms are located near the circle of Willis at the base of the brain and near arterial bifurcations (aneurysms secondary to bacterial endocarditis [mycotic aneurysms] tend to form more distally). About 85% of aneurysms occur in the carotid circulation, with the most common sites being the origin of the posterior communicating artery from the internal carotid artery (41%), anterior communicating artery/anterior cerebral artery complex (34%), and middle cerebral artery (20%).¹ Up to 20% of patients have multiple aneurysms.¹⁸

Presentation

The typical presentation of acute SAH is the sudden onset of a severe headache, nausea, vomiting, and syncope. Presenting with focal neurologic deficits is very unusual, but may be seen due to mass effect from a giant aneurysm, parenchymal hemorrhage, subdural hematoma, or a large localized subarachnoid clot. In addition, third cranial nerve palsies (dilated pupil, ptosis, and limited eye movements) may be present due to aneurysmal compression of the nerve, particularly with posterior communicating artery aneurysms. Seizures at onset may be reported by family, but it is not clear how many of these episodes represent true epileptic events versus abnormal posturing.¹⁹

Initial Management

The initial steps in managing a patient who is suspected of having a SAH focus on airway evaluation, early computed tomographic (CT) imaging, blood pressure control, serial assessment of neurological function, and stabilization to undergo additional diagnostic studies (CT angiography, catheter angiography). Clinical status is usually described using the method proposed by Hunt and Hess²⁰ and World Federation of Neurological Surgeons (WFNS) Scales²¹ (–Table 1 and 2).

Diagnosis

A noncontrast CT scan remains the diagnostic test of choice to detect SAH. Current CT scanners have a sensitivity of 98.7% when performed within 6 hours of onset in patients presenting with thunderclap headache and a normal neurological examination.²² Blood appears as a hyperdense signal in the cisterns surrounding the brainstem. CT may be falsely negative if the volume of blood is very small, if the hemorrhage occurred several days prior, if the blood is primarily

Table 1 Hunt–Hess scale

Grade	Symptoms
1	Asymptomatic or mild headache
2	Moderate to severe headache, nuchal rigidity, with or without cranial nerve deficits
3	Confusion, lethargy, or mild focal symptoms
4	Stupor and/or hemiparesis
5	Deep coma, decerebrate rigidity

Table 2 World Federation of Neurologic Surgeons

Grade	Glasgow Coma Scale	Motor deficits
I	15	Absent
II	13–14	Absent
III	13–14	Present
IV	7–12	Present or absent
V	3–6	Present or absent

located in the posterior fossa, or if the hematocrit is extremely low. Traditional teaching has been that if the CT is normal and suspicion of SAH remains strong, a lumbar puncture (LP) should be performed. More recent studies suggest that the yield of LP in this setting is quite low.²³

Conventional catheter angiography is the gold standard for the detection of intracranial aneurysms. CT angiography has recently improved to the point where some centers perform it routinely.^{24,25} Magnetic resonance angiography techniques are rapidly advancing, but there can be some limitations of its use in the acute care setting.

In 15 to 20% of cases, angiography fails to demonstrate the cause of SAH.²⁶ If a high-quality complete angiography does not identify a source of bleeding, patients have a very low incidence of rebleeding, especially if the blood is limited to the perimesencephalic and ambient cisterns.²⁷

If the patient is lethargic or agitated, consideration should be given to elective intubation to facilitate performing safe and rapid angiography.

Medical Complications

Blood pressure is typically elevated following SAH due to pain and anxiety and sympathetic activation. Left untreated, hypertension is thought to increase the risk of aneurysmal rerupture; however, few data exist to support this notion.²⁸ Still, most recommend normalization of blood pressure.²⁹ Analgesics alone may be effective; otherwise, rapidly acting antihypertensive agents are employed. Preferred agents include labetalol, hydralazine, and nicardipine.^{30,31} An important exception to treatment of hypertension is when hydrocephalus is present. In that situation, blood pressure should be addressed only after the hydrocephalus is treated.

Cardiovascular abnormalities are common in SAH. Electrocardiographic changes include sinus arrhythmia, tall

peaked “cerebral” T-waves, ST segment depression, and prolonged QT segments.³² Cardiac enzymes are often mildly elevated. Arrhythmias occur frequently, but are most commonly benign unless associated with electrolyte abnormalities; however, some patients may have more serious arrhythmias, sometimes leading to cardiac arrest.

In rare cases, patients may present with severe cardiac dysfunction. Myocardial contractility is markedly impaired, leading to a fall in cardiac output and blood pressure as well as pulmonary edema.^{33,34} This condition has been referred to as “stunned myocardium,” and may also include an element of neurogenic pulmonary edema.^{8,35} The typical pattern on echocardiography is that of Takotsubo cardiomyopathy,³⁶ and management is similar to other causes of acute pump failure with inotropic agents, diuretics, high concentrations of oxygen, and PEEP.^{8,36} The condition is remarkably transient and completely resolves in a few days.^{37,38} The severity of the SAH is the most important predictor of cardiac dysfunction.

Initial ICU Management

Patients may be admitted to the intensive care unit (ICU) directly from the emergency department or following surgical or endovascular repair of the aneurysm. Routine monitoring includes serial neurological examinations; continuous electrocardiographic monitoring; and frequent determinations of blood pressure, electrolytes, body weight, fluid balance, and, in many centers, transcranial Doppler (TCD) measurement of cerebral blood flow velocities and continuous electroencephalography.

Prevention of Rebleeding

The risk of rebleeding is highest immediately following SAH, occurring in 8 to 10% within the first 24 hours, and declining over the next few days. Rates are highest in women and those with a poor clinical grade, in poor medical condition, and with elevated systolic blood pressure. More than half of the patients who rebleed die.

Prior to aneurysm repair, factors associated with rebleeding (e.g., cough, Valsalva) should be minimized. Excessive stimulation should be avoided. Agitated patients should be sedated with short-acting agents, but should remain responsive enough for assessment of neurologic status. Over-sedation should be avoided, as it may mask clinical deterioration. Definitive prevention of rebleeding is by repair of the aneurysm, either by surgical or endovascular means. This should be completed within 24 hours of presentation.

Vasospasm and Delayed Cerebral Ischemia

While in general the term “vasospasm” implies vasoconstriction leading to a reduction in the caliber of a vessel, it has many other meanings when used in the context of SAH. For the purposes of this discussion, “vasospasm” will be used to refer to the blood vessel changes that occur and are seen on cerebral angiography. Delayed neurologic deficits with or

without vasospasm will be referred to as delayed neurologic ischemia (DCI).

The relationship between SAH and arterial narrowing seen on cerebral angiography was first described by Ecker and Riemenschneider in 1951.³⁹ Previously, Robertson ascribed ischemic brain lesions identified on autopsy to probable spasm of arteries.⁴⁰ Fisher and colleagues published a sentinel paper in 1977 in which they associated neurologic deficits with vasospasm.⁴¹ From that point until very recently, arterial vasospasm was considered the proximate cause of delayed neurologic deterioration, ischemia, and infarction.⁴² Over the past decade, several observations have weakened that association and opened up many new areas of investigation.^{43,44}

Histologically, following SAH there are structural alterations in endothelial and smooth muscle cells of the arterial wall.⁴⁵ The release of oxyhemoglobin into the subarachnoid space seems to instigate these changes.⁴⁶ Oxyhemoglobin stimulates the secretion of endothelin (ET)-1, a potent vasoconstrictor;⁴⁷ inhibits the vasodilatory effects of nitric oxide (NO); and produces oxygen-free radicals,⁴⁸ which may play a role in lipid peroxidation, possibly mediating the structural changes in the vessel wall.

The role of inflammation in the development of vasospasm is an active area of investigation. The incidence of vasospasm is increased in the presence of SIRS,⁹ and cerebrospinal fluid (CSF) levels of cytokines are elevated in patients who develop vasospasm.^{49,50} Endothelial NO synthase (eNOS) gene polymorphisms are associated with vasospasm.⁵¹

Vasospasm has been clearly linked to neurologic deficits, reduced cerebral blood flow, and ischemia and infarction; yet, all of these can occur in its absence.^{52,53} Large trials of an endothelin antagonist demonstrated a reduction in the incidence of vasospasm, but had no effect on clinical outcome.^{54,55} Regions of reduced cerebral blood flow and cerebral infarction are frequently seen independent of vasospasm.^{56,57} Several alternative hypotheses to explain this discrepancy have been proposed and are under investigation; however, no insights to date have provided a clear direction for clinical intervention.

Reducing the Impact of Vasospasm/DCI

Due to their delayed onset, vasospasm and DCI are very attractive targets for preventive therapies. By targeting various steps, their development may be entirely averted or their impact reduced. Since the volume of subarachnoid blood is a strong predictor of vasospasm and DCI, reduction of SAH volume was a focus of early interventions.

Cisternal Thrombolysis and Lumbar CSF Drainage

Cisternal irrigation with saline is routinely performed at the time of surgery to clear subarachnoid blood. Its efficacy has never been directly tested, however. Since the rates of vasospasm and DCI appear to be the same whether the aneurysm is repaired by surgical or endovascular means, the effectiveness of this practice is in doubt. Attempts have been made to enhance clearance by instilling tissue

plasminogen activator (t-PA) into the cisterns and ventricles, and meta-analyses suggest some potential benefit.^{58,59} In a large randomized trial, there was no overall effect on vasospasm, but in patients with thick subarachnoid clots, there was a 56% relative risk reduction of severe vasospasm in the t-PA-treated group.⁶⁰ An alternative approach is to instill t-PA into a ventricular drain,⁶¹ which may accelerate the clearance of both ventricular and subarachnoid blood, especially when administered early.⁶²

Attempts have been made to apply this approach to patients undergoing coil embolization of aneurysms. In those patients, a microcatheter was inserted in the lumbar region and navigated up to the cisterns where urokinase was instilled; the treatment appeared to reduce symptomatic vasospasm.⁶³

Another approach to clearing blood from the subarachnoid space is through lumbar CSF drainage. In a non-randomized controlled-cohort study, CSF drainage reduced the incidence of clinical vasospasm, the use of angioplasty, and vasospasm-related infarction.⁶⁴ A subsequent single-center, prospective, randomized controlled trial found reduced prevalence of DCI and improved early outcome, but no improvement in outcome at 6 months.⁶⁵ Additional studies are underway.

Calcium Channel Blockers

Enteral

Nimodipine is the only drug approved by the U.S. Food and Drug Administration for use in the treatment of vasospasm. A dihydropyridine, it blocks calcium influx through the L-type calcium channels in blood vessels and neurons. It reduces the risk of poor outcome and secondary ischemia after aneurysmal SAH.^{66–68}

How nimodipine exerts its beneficial effects is incompletely understood. It may involve neuronal as well as vascular effects, acting as a neuroprotective and vasodilatory agent.⁶⁹ It does not appear to reverse angiographic vasospasm, but this has not been well studied. Nimodipine is routinely administered to all aneurysmal SAH patients beginning from the time of presentation through 21 days. The approved dose is 60 mg orally every 4 hours, the dose being limited by hypotension. Some patients do not tolerate the recommended dose, especially when being treated with induced hypertension for DCI. This appears to be exacerbated by hypovolemia. Dose adjustment to 30 mg every 2 hours may limit hypotension, but in some patients, nimodipine is discontinued due to refractory hypotension. This has its downside, however; high compliance with scheduled nimodipine is associated with better outcome.⁷⁰

In a large randomized trial, oral nicardipine decreased the incidence of DCI, reduced the use of rescue therapy for refractory DCI, and reduced angiographic vasospasm; yet, it did not improve overall outcome at 3 months.⁷¹ While, based on this trial, the use of nicardipine was largely abandoned in the United States, some argue that the lack of benefit seen in the trial was due to increased use of rescue therapies in the control group. Phase I and II safety studies of diltiazem in SAH demonstrated safety but no effect on vasospasm.⁷²

Intravenous

While intravenous calcium channel blockers are often used for blood pressure control in SAH patients,^{30,73} they have also been administered in an attempt to reduce vasospasm and DCI. In Europe, nimodipine is also used as a continuous intravenous infusion, although this is often associated with hypotension requiring dose adjustment.^{74,75}

Intrathecal

Nimodipine and nicardipine cross the blood–brain barrier, but they do so in very low concentrations. As doses are limited by hypotension, attempts have been made to administer the drugs directly into the subarachnoid space. This approach is challenging as it must be available for days, requiring repeated administration or use of a sustained-release formulation.

One approach has been the use of prolonged-release implants placed at the time of aneurysm clipping.⁷⁶ In a small phase II study, use of implants reduced the incidence of cerebral vasospasm and DCI and improved clinical outcome.⁷⁷ Early studies of intrathecal nicardipine have shown encouraging results.⁷⁸ The main limitation to the use of implants is that they must be surgically implanted. The growth of endovascular methods to repair aneurysms makes this option unavailable to many SAH patients.

An alternative approach that could be used in both surgical and endovascular patients is to administer the drug directly into the ventricular system via an external ventricular drain. The feasibility of this approach has been greatly enhanced by the development of sustained-release microparticles.⁷⁹ They have the advantages of requiring only a single administration, instilling the drug directly into the site of action, and not requiring a craniotomy. In a recently completed phase I/IIa study, intraventricular nimodipine sustained-release microparticles were compared with standard dose oral nimodipine. The microparticles were safe and well tolerated without hypotension. They were associated with reduced DCI and needed for rescue therapy, as well as improved clinical outcome.⁸⁰

Prophylactic Hypervolemia

It is widely accepted that SAH patients can spontaneously develop hypovolemia and that hypovolemia can impair cerebral perfusion and increase risk of DCI. Thus, it was proposed that prophylactic hypervolemia may reduce DCI and improve outcome in SAH patients. However, in prospective controlled studies, it failed to reduce the incidence of DCI and vasospasm, did not improve cerebral blood flow, and had no effect on outcome.^{81–84} In addition, costs and complications were higher in the group treated with prophylactic hypervolemia.⁸³ To date, prophylactic hypervolemic therapy has not been shown to effectively raise cerebral blood flow or improve neurological outcome. In addition, there is evidence for harm using overly aggressive hydration; however, careful monitoring of volume status is recommended to avoid hypovolemia.⁸⁵

Prophylactic Transluminal Balloon Angioplasty

Based on success in a dog model of SAH, a pilot study of prophylactic transluminal balloon angioplasty in patients

with Fisher grade 3 SAH was performed, which had promising results.⁸⁶ In a larger trial, fewer patients developed DCI after treatment with transluminal balloon angioplasty and there was a reduced need for therapeutic angioplasty.⁸⁷ While the procedure appeared to be beneficial, this was offset by a high rate of procedural complications resulting in three deaths, and the approach was abandoned. Since that time, advances in angioplasty balloon design have renewed interest in the approach.

Tirilazad

Following promising results in primate vasospasm models, tirilazad, a non-glucocorticoid 21 amino-steroid free radical scavenger, was studied in SAH. The effect on outcome was variable, in part related to gender differences in drug metabolism and an interaction with phenytoin.^{88–91}

Magnesium

Magnesium is a noncompetitive calcium antagonist and has potentially neuroprotective vascular effects. It can lead to vasodilatation by blocking the voltage-dependent calcium channel and decreasing entry of calcium into the cell. In addition, magnesium attenuates the effect of various potent vasoconstrictors, such as endothelin 1, and blocks the formation of reactive oxygen species. Several small trials of aggressive intravenous magnesium administration suggested benefit.⁹² In a phase II trial enrolling 283 patients, magnesium treatment reduced the risk of DCI by 34% and poor outcome by 23%.⁹³ However, a follow-up phase III study of 1,204 patients failed to show any benefit.⁹⁴

More recently, attempts have been made to administer magnesium directly into the CSF via continuous cisternal irrigation. In a preliminary trial of 73 patients, a reduction in vasospasm was noted, but there was neither a reduced incidence of DCI nor improved outcome.⁹⁵

Statins

Statins appear to have neuroprotective properties independent of cholesterol reduction due to their ability to upregulate endothelial nitric oxide synthase. Early small randomized trials found improved outcome in patients treated with simvastatin and pravastatin.^{96–98} In an international, multicenter, randomized, double-blind trial, 803 patients were randomly assigned to receive either simvastatin 40 mg or placebo for 21 days. The trial did not find any benefit for short- or long-term outcome.⁹⁹

Clazosentan

Endothelin is a potent vasoconstrictor and is thought to play a key role in the development of arterial vasospasm. Clazosentan is an endothelin receptor antagonist designed to inhibit endothelin-mediated cerebral vasospasm. Two early trials demonstrated the ability of the drug to reduce angiographic vasospasm.^{100,101} They were followed by two large trials powered to detect an effect on clinical outcome.^{54,55} Both of these studies demonstrated a clear reduction in angiographic vasospasm, but while the higher dose reduced short-term vasospasm-related morbidity and all-cause mor-

tality in one study, neither study demonstrated improved 3-month clinical outcome.

Summary

Vasospasm and DCI following aneurysmal SAH are major contributors to morbidity and mortality. The delayed nature of their onset has made them an attractive target for preventive therapies. Interventions addressing various steps in the development of vasospasm have been attempted with variable success.

Enteral nimodipine remains the only approved measure to prevent DCI. Since oral and intravenous administration is limited by hypotension, direct administration via sustained-release pellets placed during surgery or intraventricular administration of sustained-release microparticles has been investigated. These approaches show promise of being even more effective, and confirmatory trials are underway.

Efforts to remove blood from the subarachnoid space via cisternal irrigation, cisternal or ventricular thrombolysis, and lumbar CSF drainage have met with limited and variable success, and they remain an active area of investigation.

Several interventions that had early promise have failed to show benefit when studied in large trials, including tirilazad, magnesium, statins, clazosentan, transluminal angioplasty, and hypervolemia.

References

- 1 van Gijn J, Kerr RS, Rinkel GJ. Subarachnoid haemorrhage. *Lancet* 2007;369(9558):306–318
- 2 van Gijn J, van Dongen KJ, Vermeulen M, Hijdra A. Perimesencephalic hemorrhage: a nonaneurysmal and benign form of subarachnoid hemorrhage. *Neurology* 1985;35(04):493–497
- 3 Heros RC, Kistler JP. Intracranial arterial aneurysm—an update. *Stroke* 1983;14(04):628–631
- 4 Mohr JP, Kistler JP, Lost D. Intracranial aneurysms. In: Lost D, Mohr JP, eds. *Stroke: Pathophysiology, Diagnosis and Management*. London: Churchill Livingstone; 1985:643–677
- 5 Rabinstein AA, Weigand S, Atkinson JL, Wijndicks EF. Patterns of cerebral infarction in aneurysmal subarachnoid hemorrhage. *Stroke* 2005;36(05):992–997
- 6 Hijdra A, Van Gijn J, Stefanko S, Van Dongen KJ, Vermeulen M, Van Crevel H. Delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage: clinicoanatomic correlations. *Neurology* 1986;36(03):329–333
- 7 Lee VH, Connolly HM, Fulgham JR, Manno EM, Brown RD Jr, Wijndicks EF. Takotsubo cardiomyopathy in aneurysmal subarachnoid hemorrhage: an underappreciated ventricular dysfunction. *J Neurosurg* 2006;105(02):264–270
- 8 Bruder N, Rabinstein A; Participants in the International Multidisciplinary Consensus Conference on the Critical Care Management of Subarachnoid Hemorrhage. Cardiovascular and pulmonary complications of aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2011;15(02):257–269
- 9 Dhar R, Diringer MN. The burden of the systemic inflammatory response predicts vasospasm and outcome after subarachnoid hemorrhage. *Neurocrit Care* 2008;8(03):404–412
- 10 Anderson CS, Feigin V, Bennett D, Lin RB, Hankey G, Jamrozik K; Australasian Cooperative Research on Subarachnoid Hemorrhage Study (ACROSS) Group. Active and passive smoking and the risk of subarachnoid hemorrhage: an international population-based case-control study. *Stroke* 2004;35(03):633–637

- 11 Anderson C, Ni Mhurchu C, Scott D, Bennett D, Jamrozik K, Hankey G; Australasian Cooperative Research on Subarachnoid Hemorrhage Study Group. Triggers of subarachnoid hemorrhage: role of physical exertion, smoking, and alcohol in the Australasian Cooperative Research on Subarachnoid Hemorrhage Study (ACROSS). *Stroke* 2003;34(07):1771–1776
- 12 Raaymakers TW. Aneurysms in relatives of patients with subarachnoid hemorrhage: frequency and risk factors. MARS Study Group. Magnetic resonance angiography in relatives of patients with subarachnoid hemorrhage. *Neurology* 1999;53(05):982–988
- 13 Bor AS, Rinkel GJ, Adami J, et al. Risk of subarachnoid haemorrhage according to number of affected relatives: a population based case-control study. *Brain* 2008;131(Pt 10):2662–2665
- 14 Martínez-González NA, Sudlow CL. Effects of apolipoprotein E genotype on outcome after ischaemic stroke, intracerebral haemorrhage and subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry* 2006;77(12):1329–1335
- 15 Mohan D, Munteanu V, Coman T, Ciurea AV. Genetic factors involves in intracranial aneurysms—actualities. *J Med Life* 2015;8(03):336–341
- 16 Lonjon M, Pennes F, Sedat J, Bataille B. Epidemiology, genetic, natural history and clinical presentation of giant cerebral aneurysms. *Neurochirurgie* 2015;61(06):361–365
- 17 Debette S, Germain DP. Neurologic manifestations of inherited disorders of connective tissue. *Handb Clin Neurol* 2014;119:565–576
- 18 Nehls DG, Flom RA, Carter LP, Spetzler RF. Multiple intracranial aneurysms: determining the site of rupture. *J Neurosurg* 1985;63(03):342–348
- 19 Lanzino G, D'Urso PI, Suarez J; Participants in the International Multi-Disciplinary Consensus Conference on the Critical Care Management of Subarachnoid Hemorrhage. Seizures and anticonvulsants after aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2011;15(02):247–256
- 20 Hunt WE, Hess RM. Surgical risk as related to time of intervention in the repair of intracranial aneurysms. *J Neurosurg* 1968;28(01):14–20
- 21 Report of World Federation of Neurological Surgeons Committee on a universal subarachnoid hemorrhage grading scale. *J Neurosurg* 1988;68(06):985–986
- 22 Dubosh NM, Bellolio MF, Rabinstein AA, Edlow JA. Sensitivity of early brain computed tomography to exclude aneurysmal subarachnoid hemorrhage: a systematic review and meta-analysis. *Stroke* 2016;47(03):750–755
- 23 Bakr A, Silva D, Cramb R, Flint G, Foroughi M. Outcomes of CSF spectrophotometry in cases of suspected subarachnoid haemorrhage with negative CT: two years retrospective review in a Birmingham hospital. *Br J Neurosurg* 2017;31(02):223–226
- 24 Grasso G, Alafaci C, Macdonald RL. Management of aneurysmal subarachnoid hemorrhage: State of the art and future perspectives. *Surg Neurol Int* 2017;8:11
- 25 Lawton MT, Vates GE. Subarachnoid hemorrhage. *N Engl J Med* 2017;377(03):257–266
- 26 Qu F, Aiyagari V, Cross DT III, Dacey RG Jr, Diringer MN. Untreated subarachnoid hemorrhage: who, why, and when? *J Neurosurg* 2004;100(02):244–249
- 27 Dalyai R, Chalouhi N, Theofanis T, et al. Subarachnoid hemorrhage with negative initial catheter angiography: a review of 254 cases evaluating patient clinical outcome and efficacy of short- and long-term repeat angiography. *Neurosurgery* 2013;72(04):646–652, discussion 651–652
- 28 Starke RM, Connolly ES Jr; Participants in the International Multi-Disciplinary Consensus Conference on the Critical Care Management of Subarachnoid Hemorrhage. Rebleeding after aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2011;15(02):241–246
- 29 Diringer MN, Bleck TP, Claude Hemphill J III, et al; Neurocritical Care Society. Critical care management of patients following aneurysmal subarachnoid hemorrhage: recommendations from the Neurocritical Care Society's Multidisciplinary Consensus Conference. *Neurocrit Care* 2011;15(02):211–240
- 30 Ortega-Gutierrez S, Thomas J, Reccius A, et al. Effectiveness and safety of nicardipine and labetalol infusion for blood pressure management in patients with intracerebral and subarachnoid hemorrhage. *Neurocrit Care* 2013;18(01):13–19
- 31 Woloszyn AV, McAllen KJ, Figueroa BE, DeShane RS, Barletta JF. Retrospective evaluation of nicardipine versus labetalol for blood pressure control in aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2012;16(03):376–380
- 32 Sommargren CE. Electrocardiographic abnormalities in patients with subarachnoid hemorrhage. *Am J Crit Care* 2002;11:48–56
- 33 Chen Z, Venkat P, Seyfried D, Chopp M, Yan T, Chen J. Brain-heart interaction: cardiac complications after stroke. *Circ Res* 2017;121(04):451–468
- 34 Kerro A, Woods T, Chang JJ. Neurogenic stunned myocardium in subarachnoid hemorrhage. *J Crit Care* 2017;38:27–34
- 35 Pinnamaneni S, Aronow WS, Frishman WH. Neurocardiac injury after cerebral and subarachnoid hemorrhages. *Cardiol Rev* 2017;25(02):89–95
- 36 Wartenberg KE, Schmidt JM, Claassen J, et al. Impact of medical complications on outcome after subarachnoid hemorrhage. *Crit Care Med* 2006;34(03):617–623, quiz 624
- 37 Deibert E, Barzilai B, Braverman AC, et al. Clinical significance of elevated troponin I levels in patients with nontraumatic subarachnoid hemorrhage. *J Neurosurg* 2003;98(04):741–746
- 38 Deibert E, Aiyagari V, Diringer MN. Reversible left ventricular dysfunction associated with raised troponin I after subarachnoid haemorrhage does not preclude successful heart transplantation. *Heart* 2000;84(02):205–207
- 39 Ecker A, Riemenschneider PA. Arteriographic demonstration of spasm of the intracranial arteries, with special reference to saccular arterial aneurysms. *J Neurosurg* 1951;8(06):660–667
- 40 Robertson EG. Cerebral lesions due to intracranial aneurysms. *Brain* 1949;72(Pt 2):150–185
- 41 Fisher CM, Roberson GH, Ojemann RG. Cerebral vasospasm with ruptured saccular aneurysm—the clinical manifestations. *Neurosurgery* 1977;1(03):245–248
- 42 Macdonald RL. Origins of the concept of vasospasm. *Stroke* 2016;47(01):e11–e15
- 43 Macdonald RL. Does prevention of vasospasm in subarachnoid hemorrhage improve clinical outcome? Yes. *Stroke* 2013;44(06, Suppl 1):S31–S33
- 44 Diringer MN. Controversy: does prevention of vasospasm in subarachnoid hemorrhage improve clinical outcome? *Stroke* 2013;44(06, Suppl 1):S29–S30
- 45 Smith RR, Clower BR, Grotendorst GM, Yabuno N, Cruse JM. Arterial wall changes in early human vasospasm. *Neurosurgery* 1985;16(02):171–176
- 46 Macdonald RL, Weir BK. A review of hemoglobin and the pathogenesis of cerebral vasospasm. *Stroke* 1991;22(08):971–982
- 47 Fuwa I, Mayberg M, Gadjusek C, Harada T, Luo Z. Enhanced secretion of endothelin by endothelial cells in response to hemoglobin. *Neurol Med Chir (Tokyo)* 1993;33(11):739–743
- 48 Misra HP, Fridovich I. The generation of superoxide radical during the autoxidation of hemoglobin. *J Biol Chem* 1972;247(21):6960–6962
- 49 Zeiler FA, Thelin EP, Czosnyka M, Hutchinson PJ, Menon DK, Helmy A. Cerebrospinal fluid and microdialysis cytokines in aneurysmal subarachnoid hemorrhage: a scoping systematic review. *Front Neurol* 2017;8:379
- 50 Rodríguez-Rodríguez A, Egea-Guerrero JJ, Ruiz de Azúa-López Z, Murillo-Cabezas F. Biomarkers of vasospasm development and

- outcome in aneurysmal subarachnoid hemorrhage. *J Neurol Sci* 2014;341(1-2):119–127
- 51 Khurana VG, Fox DJ, Meissner I, Meyer FB, Spetzler RF. Update on evidence for a genetic predisposition to cerebral vasospasm. *Neurosurg Focus* 2006;21(03):E3
 - 52 Dankbaar JW, Rijdsdijk M, van der Schaaf IC, Velthuis BK, Wermer MJ, Rinkel GJ. Relationship between vasospasm, cerebral perfusion, and delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage. *Neuroradiology* 2009;51(12):813–819
 - 53 Vergouwen MD, Ildigwe D, Macdonald RL. Cerebral infarction after subarachnoid hemorrhage contributes to poor outcome by vasospasm-dependent and -independent effects. *Stroke* 2011;42(04):924–929
 - 54 Macdonald RL, Higashida RT, Keller E, et al. Randomized trial of clazosentan in patients with aneurysmal subarachnoid hemorrhage undergoing endovascular coiling. *Stroke* 2012;43(06):1463–1469
 - 55 Macdonald RL, Higashida RT, Keller E, et al. Randomised trial of clazosentan, an endothelin receptor antagonist, in patients with aneurysmal subarachnoid hemorrhage undergoing surgical clipping (CONSCIOUS-2). *Acta Neurochir Suppl (Wien)* 2013;115:27–31
 - 56 Dhar R, Diringer MN. Relationship between angiographic vasospasm, cerebral blood flow, and cerebral infarction after subarachnoid hemorrhage. *Acta Neurochir Suppl (Wien)* 2015;120:161–165
 - 57 Brown RJ, Kumar A, Dhar R, Sampson TR, Diringer MN. The relationship between delayed infarcts and angiographic vasospasm after aneurysmal subarachnoid hemorrhage. *Neurosurgery* 2013;72(05):702–707, discussion 707–708
 - 58 Hänggi D, Steiger HJ. The influence of cisternal and ventricular lavage on cerebral vasospasm in patients suffering from subarachnoid hemorrhage: analysis of effectiveness. *Acta Neurochir Suppl (Wien)* 2011;110(Pt 2):95–98
 - 59 Amin-Hanjani S, Ogilvy CS, Barker FG II. Does intracisternal thrombolysis prevent vasospasm after aneurysmal subarachnoid hemorrhage? A meta-analysis. *Neurosurgery* 2004;54(02):326–334, discussion 334–335
 - 60 Findlay JM, Kassell NF, Weir BK, et al. A randomized trial of intraoperative, intracisternal tissue plasminogen activator for the prevention of vasospasm. *Neurosurgery* 1995;37(01):168–176, discussion 177–178
 - 61 Kramer AH, Roberts DJ, Holodinsky J, et al. Intraventricular tissue plasminogen activator in subarachnoid hemorrhage patients: a prospective, randomized, placebo-controlled pilot trial. *Neurocrit Care* 2014;21(02):275–284
 - 62 Kramer AH, Jenne C, Holodinsky JK, et al. Pharmacokinetics and pharmacodynamics of tissue plasminogen activator administered through an external ventricular drain. *Neurocrit Care* 2015;23(03):386–393
 - 63 Hamada J, Kai Y, Morioka M, et al. Effect on cerebral vasospasm of coil embolization followed by microcatheter intrathecal urokinase infusion into the cisterna magna: a prospective randomized study. *Stroke* 2003;34(11):2549–2554
 - 64 Klimo P Jr, Kestle JR, MacDonald JD, Schmidt RH. Marked reduction of cerebral vasospasm with lumbar drainage of cerebrospinal fluid after subarachnoid hemorrhage. *J Neurosurg* 2004;100(02):215–224
 - 65 Al-Tamimi YZ, Bhargava D, Feltbower RG, et al. Lumbar drainage of cerebrospinal fluid after aneurysmal subarachnoid hemorrhage: a prospective, randomized, controlled trial (LUMAS). *Stroke* 2012;43(03):677–682
 - 66 Allen GS, Ahn HS, Preziosi TJ, et al. Cerebral arterial spasm—a controlled trial of nimodipine in patients with subarachnoid hemorrhage. *N Engl J Med* 1983;308(11):619–624
 - 67 Pickard JD, Murray GD, Illingworth R, et al. Effect of oral nimodipine on cerebral infarction and outcome after subarachnoid haemorrhage: British aneurysm nimodipine trial. *BMJ* 1989;298(6674):636–642
 - 68 Velat GJ, Kimball MM, Mocco JD, Hoh BL. Vasospasm after aneurysmal subarachnoid hemorrhage: review of randomized controlled trials and meta-analyses in the literature. *World Neurosurg* 2011;76(05):446–454
 - 69 Feigin VL, Rinkel GJ, Algra A, Vermeulen M, van Gijn J. Calcium antagonists in patients with aneurysmal subarachnoid hemorrhage: a systematic review. *Neurology* 1998;50(04):876–883
 - 70 Wessell A, Kole MJ, Badjatia N, et al. High compliance with scheduled nimodipine is associated with better outcome in aneurysmal subarachnoid hemorrhage patients cotreated with heparin infusion. *Front Neurol* 2017;8:268
 - 71 Haley EC Jr, Kassell NF, Torner JC. A randomized trial of nicardipine in subarachnoid hemorrhage: angiographic and transcranial Doppler ultrasound results. A report of the Cooperative Aneurysm Study. *J Neurosurg* 1993;78(04):548–553
 - 72 Saunders FW, Marshall WJ. Diltiazem: dose it affect vasospasm? *Surg Neurol* 1986;26(02):155–158
 - 73 Li Y, Fang W, Tao L, et al. Efficacy and safety of intravenous nimodipine administration for treatment of hypertension in patients with intracerebral hemorrhage. *Neuropsychiatr Dis Treat* 2015;11:1231–1238
 - 74 Bele S, Proescholdt MA, Hochreiter A, et al. Continuous intra-arterial nimodipine infusion in patients with severe refractory cerebral vasospasm after aneurysmal subarachnoid hemorrhage: a feasibility study and outcome results. *Acta Neurochir (Wien)* 2015;157(12):2041–2050
 - 75 Rinkel GJ, Feigin VL, Algra A, Vermeulen M, van Gijn J. Calcium antagonists for aneurysmal subarachnoid haemorrhage. *Cochrane Database Syst Rev* 2002;(04):CD000277
 - 76 Zhang YP, Shields LB, Yao TL, Dashti SR, Shields CB. Intrathecal treatment of cerebral vasospasm. *J Stroke Cerebrovasc Dis* 2013;22(08):1201–1211
 - 77 Barth M, Capelle HH, Weidauer S, et al. Effect of nicardipine prolonged-release implants on cerebral vasospasm and clinical outcome after severe aneurysmal subarachnoid hemorrhage: a prospective, randomized, double-blind phase IIa study. *Stroke* 2007;38(02):330–336
 - 78 Ehtisham A, Taylor S, Bayless L, Samuels OB, Klein MW, Janzen JM. Use of intrathecal nicardipine for aneurysmal subarachnoid hemorrhage-induced cerebral vasospasm. *South Med J* 2009;102(02):150–153
 - 79 Hänggi D, Etminan N, Macdonald RL, et al. NEWTON: nimodipine microparticles to enhance recovery while reducing toxicity after subarachnoid hemorrhage. *Neurocrit Care* 2015;23(02):274–284
 - 80 Hänggi D, Etminan N, Aldrich F, et al; NEWTON Investigators. Randomized, open-label, phase 1/2a study to determine the maximum tolerated dose of intraventricular sustained release nimodipine for subarachnoid hemorrhage (NEWTON [Nimodipine Microparticles to Enhance Recovery While Reducing Toxicity After Subarachnoid Hemorrhage]). *Stroke* 2017;48(01):145–151
 - 81 Rinkel GJ, Feigin VL, Algra A, van Gijn J. Circulatory volume expansion therapy for aneurysmal subarachnoid haemorrhage. *Cochrane Database Syst Rev* 2004;(04):CD000483
 - 82 Lennihan L, Mayer SA, Fink ME, et al. Effect of hypervolemic therapy on cerebral blood flow after subarachnoid hemorrhage: a randomized controlled trial. *Stroke* 2000;31(02):383–391
 - 83 Egge A, Waterloo K, Sjøholm H, Solberg T, Ingebrigtsen T, Romner B. Prophylactic hyperdynamic postoperative fluid therapy after aneurysmal subarachnoid hemorrhage: a clinical, prospective, randomized, controlled study. *Neurosurgery* 2001;49(03):593–605, discussion 605–606
 - 84 Capampangan DJ, Wellik KE, Aguilar MI, Demaerschalk BM, Wingerchuk DM. Does prophylactic postoperative hypervolemic therapy prevent cerebral vasospasm and improve clinical outcome after aneurysmal subarachnoid hemorrhage? *Neurologist* 2008;14(06):395–398

- 85 Wolf S; Participants in the International Multi-Disciplinary Consensus Conference on the Critical Care Management of Subarachnoid Hemorrhage. Routine management of volume status after aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2011;15(02):275–280
- 86 Muizelaar JP, Zwienerberg M, Rudisill NA, Hecht ST. The prophylactic use of transluminal balloon angioplasty in patients with Fisher Grade 3 subarachnoid hemorrhage: a pilot study. *J Neurosurg* 1999;91(01):51–58
- 87 Zwienerberg-Lee M, Hartman J, Rudisill N, et al; Balloon Prophylaxis for Aneurysmal Vasospasm (BPAV) Study Group. Effect of prophylactic transluminal balloon angioplasty on cerebral vasospasm and outcome in patients with Fisher grade III subarachnoid hemorrhage: results of a phase II multicenter, randomized, clinical trial. *Stroke* 2008;39(06):1759–1765
- 88 Kassell NF, Haley EC Jr, Apperson-Hansen C, Alves WM. Randomized, double-blind, vehicle-controlled trial of tirilazad mesylate in patients with aneurysmal subarachnoid hemorrhage: a cooperative study in Europe, Australia, and New Zealand. *J Neurosurg* 1996;84(02):221–228
- 89 Lanzino G, Kassell NF, Dorsch NW, et al. Double-blind, randomized, vehicle-controlled study of high-dose tirilazad mesylate in women with aneurysmal subarachnoid hemorrhage. Part I. A cooperative study in Europe, Australia, New Zealand, and South Africa. *J Neurosurg* 1999;90(06):1011–1017
- 90 Lanzino G, Kassell NF. Double-blind, randomized, vehicle-controlled study of high-dose tirilazad mesylate in women with aneurysmal subarachnoid hemorrhage. Part II. A cooperative study in North America. *J Neurosurg* 1999;90(06):1018–1024
- 91 Haley EC Jr, Kassell NF, Apperson-Hansen C, Maile MH, Alves WM. A randomized, double-blind, vehicle-controlled trial of tirilazad mesylate in patients with aneurysmal subarachnoid hemorrhage: a cooperative study in North America. *J Neurosurg* 1997;86(03):467–474
- 92 Suarez JJ; Participants in the International Multidisciplinary Consensus Conference on the Critical Care Management of Subarachnoid Hemorrhage. Magnesium sulfate administration in subarachnoid hemorrhage. *Neurocrit Care* 2011;15(02):302–307
- 93 van den Bergh WM, Algra A, van Kooten F, et al; MASH Study Group. Magnesium sulfate in aneurysmal subarachnoid hemorrhage: a randomized controlled trial. *Stroke* 2005;36(05):1011–1015
- 94 Dorhout Mees SM, Algra A, Vandertop WP, et al; MASH-2 Study Group. Magnesium for aneurysmal subarachnoid haemorrhage (MASH-2): a randomised placebo-controlled trial. *Lancet* 2012;380(9836):44–49
- 95 Yamamoto T, Mori K, Esaki T, Nakao Y, Tokugawa J, Watanabe M. Preventive effect of continuous cisternal irrigation with magnesium sulfate solution on angiographic cerebral vasospasms associated with aneurysmal subarachnoid hemorrhages: a randomized controlled trial. *J Neurosurg* 2016;124(01):18–26
- 96 Tseng MY, Czosnyka M, Richards H, Pickard JD, Kirkpatrick PJ. Effects of acute treatment with pravastatin on cerebral vasospasm, autoregulation, and delayed ischemic deficits after aneurysmal subarachnoid hemorrhage: a phase II randomized placebo-controlled trial. *Stroke* 2005;36(08):1627–1632
- 97 Lynch JR, Wang H, McGirt MJ, et al. Simvastatin reduces vasospasm after aneurysmal subarachnoid hemorrhage: results of a pilot randomized clinical trial. *Stroke* 2005;36(09):2024–2026
- 98 Chou SH, Smith EE, Badjatia N, et al. A randomized, double-blind, placebo-controlled pilot study of simvastatin in aneurysmal subarachnoid hemorrhage. *Stroke* 2008;39(10):2891–2893
- 99 Kirkpatrick PJ, Turner CL, Smith C, Hutchinson PJ, Murray GD; STASH Collaborators. Simvastatin in aneurysmal subarachnoid haemorrhage (STASH): a multicentre randomised phase 3 trial. *Lancet Neurol* 2014;13(07):666–675
- 100 Vajkoczy P, Meyer B, Weidauer S, et al. Clazosentan (AXV-034343), a selective endothelin A receptor antagonist, in the prevention of cerebral vasospasm following severe aneurysmal subarachnoid hemorrhage: results of a randomized, double-blind, placebo-controlled, multicenter phase IIa study. *J Neurosurg* 2005;103(01):9–17
- 101 Macdonald RL, Kassell NF, Mayer S, et al; CONSCIOUS-1 Investigators. Clazosentan to overcome neurological ischemia and infarction occurring after subarachnoid hemorrhage (CONSCIOUS-1): randomized, double-blind, placebo-controlled phase 2 dose-finding trial. *Stroke* 2008;39(11):3015–3021