



C. BOLM, R. MOCCI, C. SCHUMACHER, M. TURBERG, F. PUCCHETTI, J. G. HERNÁNDEZ*
(RWTH AACHEN, GERMANY; CITTADELLA UNIVERSITARIA, MONSERRATO AND
UNIVERSITY OF FIRENZE, SESTO FIORENTINO, ITALY)

Mechanochemical Activation of Iron Cyano Complexes: A Prebiotic Impact Scenario for the Synthesis of α -Amino Acid Derivatives

Angew. Chem. Int. Ed. **2018**, *57*, 2423–2426.

Mechanochemical Generation of HCN from $K_3[Fe(CN)_6]$: A Novel Strecker Protocol



Significance: A Strecker reaction of carbonyl compounds, primary amines, and potassium ferricyanide was carried out in the presence of SiO_2 under ball-milling conditions to give the corresponding α -aminonitriles in 56–73% yield (eq. 1, 11 examples). The hydration of α -aminonitrile **A** also proceeded under ball-milling conditions to afford the corresponding amino amide in 51% yield (eq. 2).

Comment: In situ generation of HCN was found to take place through mechanochemical activation of potassium ferricyanide $\{K_3[Fe(CN)_6]\}$ by ball-milling in the presence of SiO_2 . The resulting HCN was trapped in situ by a Strecker reaction with benzaldehyde and benzylamine to give **A**.

Category

Polymer-Supported
Synthesis

Key words

Strecker reaction

mechanochemical
activation

carbonyls

amines

potassium
ferricyanide

aminonitriles

Synfact
of the month

SYNFACTS Contributors: Yasuhiro Uozumi, Go Hamasaka
Synfacts 2018, 14(05), 0537 Published online: 17.04.2018
DOI: 10.1055/s-0037-1609466; **Reg-No.:** Y03318SF

2018 © THIEME STUTTGART • NEW YORK

537