













# Formylation of Amines with CO<sub>2</sub> Using a Polymeric Frustrated Lewis Pair



**Significance:** A polymeric carbon dioxide bridged frustrated Lewis pair (**A**) was prepared by treatment of an electron-deficient triarylborane immobilized on polystyrene with a sterically demanding triarylphosphine immobilized on polystyrene in the presence of CO<sub>2</sub> (eq. 1). The polymeric frustrated Lewis pair **A** catalyzed the formylation of amines with carbon dioxide in the presence of phenylsilane at room temperature to give the corresponding formamides in up to 99% yield (eq. 2). The turnover number for the formylation reached as high as 14800.

**Comment:** The polymeric frustrated Lewis pair **A** was characterized by <sup>11</sup>B, <sup>19</sup>F, and <sup>31</sup>P NMR, DLS, TEM, FT-IR, UV-Vis, and conductivity measurements. In the formylation of diethylamine, morpholine, aniline, benzophenone imine, or imidazole, the catalyst was reused seven times with slight loss of the catalytic activity.

**SYNFACTS Contributors:** Yasuhiro Uozumi, Go Hamasaka  
Synfacts 2018, 14(10), 1097 Published online: 17.09.2018  
DOI: 10.1055/s-0037-1610983; Reg-No.: Y10218SF