

K. LIAO, Y.-F. YANG, Y. LI, J. N. SANDERS, K. N. HOUK, D. G. MUSAEV, H. M. L. DAVIES\*  
 (EMORY UNIVERSITY, ATLANTA AND UNIVERSITY OF CALIFORNIA, LOS ANGELES, USA;  
 ZHEIJANG UNIVERSITY OF TECHNOLOGY, HANGZHOU, P. R. OF CHINA)  
 Design of Catalysts for Site-Selective and Enantioselective Functionalization of Non-activated Primary C–H Bonds  
*Nat. Chem.* **2018**, *10*, 1048–1055.

# Enantioselective Functionalization of Primary C–H Bonds


Category

Metal-Catalyzed  
 Asymmetric  
 Synthesis and  
 Stereoselective  
 Reactions

Key words

rhodium catalysis  
 C–H activation  
 carbenes

Synfact  
*of the month*



**Significance:** The authors previously described Rh-catalyzed functionalizations of secondary or tertiary C–H bonds (Liao et al. *Nature* **2016**, 533, 230; Liao et al. *Nature* **2017**, 551, 609). By tuning the catalyst, non-activated primary C–H bonds are functionalized with excellent site and enantioselectivities.

**SYNFACTS Contributors:** Mark Lautens, José F. Rodríguez  
 Synfacts 2018, 14(11), 1143 Published online: 18.10.2018  
 DOI: 10.1055/s-0037-1611078; Reg-No.: L13218SF

**Comment:** Subjecting chiral substrates to the reaction conditions allows access to either set of diastereomeric products when the appropriate catalyst enantiomer is employed. Stigmasteryl acetate could be selectively functionalized at its most sterically accessible primary C–H bond.