


E. J. COREY*, N. M. WEINSHENKER, T. K. SCHAAF, W. HUBER (HARVARD UNIVERSITY, CAMBRIDGE, USA)

Stereo-Controlled Synthesis of Prostaglandins F_{2α} and E₂ (dl)

J. Am. Chem. Soc. 1969, 91, 5675–5677.

Synthesis of Prostaglandins F_{2α} and E₂



Significance: In 1969, Corey and co-workers reported a landmark total synthesis of prostaglandins PGF_{2α} and PGE₂ which play an important role as hormones in mammalian tissue. Strategic application of a [2.2.1]-bicycloheptane enabled expedient access to the highly substituted cyclopentane core.

Comment: Bicycloheptane **D** was accessed through Diels-Alder reaction between diene **B** and **C**. Subsequent Baeyer-Villiger oxidation followed by hydrolysis and iodolactonization furnished lactone **G**, which was transformed into the natural products by a sequence involving two olefinations.

SYNFACTS Contributors: Erick M. Carreira, Helene Wolleb
Synfacts 2019, 15(01), 0007 Published online: 14.12.2018
DOI: 10.1055/s-0037-1611377; Reg-No.: C07518SF

2019 © Georg Thieme Verlag Stuttgart · New York

Category

Synthesis of Natural Products and Potential Drugs

Key words

prostaglandin F_{2α}
prostaglandin E₂
bicycloheptane
Diels-Alder reaction
Baeyer-Villiger oxidation
iodolactonization

Synfact Classic

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.