


A Short, Efficient, and Stereoselective Synthesis of Piperine and its Analogues

Adriano Bauer [◇]
 Jun-Hyun Nam [◇]
 Nuno Maulide* [◇]

Institute of Organic Chemistry, University of Vienna, Währinger
 Straße 38, 1090 Vienna, Austria
 nuno.maulide@univie.ac.at

[◇] These authors contributed equally to this work

Published as part of the 30 Years SYNLETT – Pearl Anniversary Issue



- ✓ Piperine synthesized in quantitative yield
- ✓ Full stereocontrol
- ✓ Modulation of the aryl and the amide moiety

Received: 26.11.2018

Accepted after revision: 16.12.2018

Published online: 14.01.2019

DOI: 10.1055/s-0037-1611652; Art ID: st-2018-b0768-I

License terms:

Abstract A quantitative synthesis of piperine from commercially available starting material is presented. The synthesis relies on a stereoselective nucleophilic attack of an *in situ* generated cuprate onto a cyclobutene lactone. The so-formed aryl-substituted cyclobutene spontaneously undergoes a conrotatory 4π -electrocyclic ring opening to form the 4-aryl pentadienoic acid as a single diastereoisomer. The high-yielding synthesis can be easily modulated on the aryl and on the amide moiety for the synthesis of a wide range of piperine analogues.

Key words total synthesis, piperine, alkaloid, cyclobutene, pericyclic reaction, stereoselective, organometallic chemistry


In 1820, the Danish physicist and chemist Hans Christian Ørstedt, pursuing an interest in the isolation of ‘new alkalis’, reported a new alkaloid from pepper (*piper nigrum*), which he called piperine.¹ Piperine would ultimately gain the attention of the chemistry and physiology communities² mostly due to its wide range of biological activities. This was foreshadowed by the original communication itself, where Ørstedt noted that an ethanolic solution of piperine has an ‘exceptionally pungent taste’.¹ The pungency of piperine can be attributed to its agonistic nature towards the heat- and acidity-sensing TRPV ion channels, which are associated with temperature and pain regulation in the human body.^{2a,b} A range of human disorders are linked to the overexpression of TRPV1, including inflammatory bowel disease (ulcerative colitis and Crohn’s disease)

and chronic breast pain.^{2c} Studies have shown that piperine is a potent desensitizer of human TRPV1, rendering its structure a potent scaffold for the design of improved TRPV1 agonists.^{2c,d}

Moreover, piperine has been recently identified as an allosteric modulator of the γ -amino butyric acid type A (GABA_A) receptor.^{3a} The mode of action of piperine is thus analogous to commonly used drugs such as benzodiazepines, widely used as sleep-inducing agents.^{3b,c} In addition, it has been shown that piperine derivatives are efficient inhibitors of vascular smooth muscle cell proliferation.⁴ Anti-depressant^{5b} and antitumor^{5c} activities along with insecticidal properties^{5d} are also attributed to this intriguing molecule which can be technically found in almost every modern kitchen in the world.

Despite the numerous demonstrated beneficial therapeutic properties of piperine, biological applications are limited by its poor solubility in aqueous media.⁶ This implies that new synthetic routes towards piperine analogues are highly desirable.

Piperine can be easily extracted⁵ and its basic hydrolysis, yielding piperic acid, opens up the preparation of many amide analogues for biological evaluation.⁷ This approach is unfortunately limited by nonexchangeability of the aryl moiety. Most approaches for the synthesis of 1-carbonyl-4-aryl-substituted dienes typically make use of a (2 + 2), (3 + 1), or a (1 + 2 + 1) carbon disconnection. In this regard Wittig olefination,⁸ olefin metathesis,⁹ palladium-catalyzed cross-coupling,¹⁰ or ruthenium-catalyzed vinyl-alkyne coupling¹¹ have been employed as dominant strategies.



Scheme 1 Previous work and this work

A rather unconventional approach is the direct coupling of the aryl moiety with the diene or a diene precursor. Mihovilovic and coworkers reported an efficient Heck reaction approach in which an aryl bromide is coupled to a pentadienoic amide (Scheme 1, a).^{4,12} Another intriguing, earlier approach, relies on the addition of a Grignard reagent to a furfural hydrazone, which rearranges to the corresponding pentadienal under the reaction conditions (Scheme 1, b).¹³


Herein we would like to present a different strategy towards the synthesis of piperine analogues. The bicyclo [2.2.0] lactone **2** and its derivatives have been deployed in previous work by our group and others as a versatile electrophile.¹⁴ In particular, we have shown that copper-mediated nucleophilic addition is a very robust method for a *trans*-selective allylic substitution of **2**.^{14f}

The installation of an electron-rich moiety (such as $-OR$ or $-N_3$) in this position has been earlier shown to facilitate a subsequent, spontaneous 4π -electrocyclic opening. This is likely due to a push-pull relationship between the carboxylic acid and the electron-donating substituent.¹⁵

We hypothesized that an aryl moiety might be sufficiently electron donating in order to induce a similar push-pull effect and enable facile electrocyclic ring opening, either spontaneously at room temperature or upon mild heating. Importantly, the transient *trans*-configured cyclobutene should undergo opening according to a thermally allowed, conrotatory movement torquoselective for the *E,E*-diene product.


Lactone **2** was prepared in quantitative yield photochemically, as previously reported.^{14f} In the event, we found that addition of the *in situ* formed cuprate (from its corresponding Grignard reagent **3a**) directly led to piperic acid (**4a**) as the sole product in a single, quantitative step.¹⁶ As expected, **4a** was formed exclusively as the *E,E*-diene isomer in a clean reaction.¹⁷ Straightforward amide formation

via acyl chloride substitution with piperidine afforded piperine in more than 95% isolated yield.¹⁸ Through the route presented herein, this alkaloid was thus available in only three quantitative steps from pyrone **1** and with full stereo-selectivity (Scheme 2).




Scheme 2 A simple three-step synthesis of piperine

Encouraged by these results, we investigated the synthesis of three different piperic acid analogues by using different Grignard reagents for the ring opening of bicyclic lactone **2**. The 5-phenylpentadienoic acid (**4b**) was prepared in quantitative yield, while the *para*-methoxyphenyl- and the 2-thiophenyl- analogues were formed in slightly lower yields. Nevertheless, geometric selectivity was excellent in all cases (Scheme 3).¹⁹ Finally, the corresponding amides were formed as before *via* standard acyl chloride substitution (Scheme 4).¹⁸ It should be noted that amides **5bb** and **5cb** have been previously reported to enhance GABA_A-induced chloride currents more strongly than natural piperine ($789\% \pm 72\%$ and $883\% \pm 70\%$, respectively).¹²



Scheme 3 Synthesis of piperic acid analogues 4



Scheme 4 Synthesis of piperine analogues 5

In conclusion, we herein presented a conceptually new approach to the synthesis of 4-aryl-substituted pentadienoic acids and their amides in excellent yield and geometrical stereoselectivity.^{20–26} This enabled a preparation of the natural product piperine in quantitative yield over three steps from commercially available 2-pyrone **1**. Analogues can be readily synthesized through this modular and operationally simple procedure.

Funding Information

Funding by the Austrian Research Fund (FWF Project P27194) is gratefully acknowledged.

Acknowledgment

We are grateful to the University of Vienna for continued generous support of our research programs.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0037-1611652>.

References and Notes

- (1) Ørstedt, H. C. *Journal für Chemie und Physik* **1820**, 29, 80.
- (2) (a) Bley, K. R. *Exper. Opin. Invest. Drugs* **2004**, 13, 1445. (b) Gavva, N. R.; Bannon, A. W.; Surapaneni, S.; Hovland, D. N. Jr; Lehto, S. G.; Gore, A.; Juan, T.; Deng, H.; Han, B.; Klionsky, L.; Kuang, R.; Le, A.; Tamir, R.; Wang, J.; Youngblood, B.; Zhu, D.; Norman, M. H.; Magal, E.; Treanor, J. J. S.; Louis, J.-C. *J. Neurosci.* **2007**, 27, 3366. (c) Szallasi, A. *Trends in Pharm. Sci.* **2005**, 26, 437. (d) Gunthorpe, M. J.; Szallasi, A. *Curr. Pharm. Design* **2008**, 14, 32.
- (3) (a) Zaugg, J.; Baburin, I.; Strommer, B.; Kim, H.-J.; Hering, S.; Hamburger, M. *J. Nat. Prod.* **2010**, 73, 185. (b) Whiting, P. *J. Curr. Opin. Pharmacol.* **2006**, 6, 24. (c) Riss, J.; Cloyd, J.; Gates, J.; Collins, S. *Acta Neurol. Scand.* **2008**, 118, 781.
- (4) Mair, E. M.; Liu, R.; Atanasov, A. G.; Wimmer, L.; Nemetz-Fiedler, D.; Sider, N.; Heiss, E. H.; Mihovilovic, M. D.; Dirsch, V. M.; Rollinger, J. M. *Planta Med.* **2015**, 81, 1065.
- (5) (a) Vasavirna, K.; Upender, M. *Int. J. Pharm. Pharm. Sci.* **2014**, 6, 34. (b) Li, S.; Wang, C.; Koike, K.; Nikaido, R.; Wang, M. W. *J. Asian Nat. Prod. Res.* **2007**, 9, 421. (c) Sunila, E. S.; Kuttan, G. *J. Ethnopharmacol.* **2004**, 90, 339. (d) Tavares, W. S.; Cruz, I.; Petacci, F.; Freitas, S. S.; Serrao, J. E.; Zanoncio, J. C. *J. Med. Plants Res.* **2011**, 5, 5301.
- (6) Gorgani, L.; Mohammadi, M.; Najafpour, G. D.; Nikzad, M. *Compr. Rev. Food Sci. Food Saf.* **2017**, 16, 124.
- (7) Umadevi, P.; Deepti, K.; Venugopal, V. R. *Med. Chem. Res.* **2013**, 22, 5466.
- (8) (a) Olsen, R. A.; Spessard, G. O. *J. Agric. Food Chem.* **1981**, 29, 942. (b) Schobert, R.; Siegfried, S.; Gordon, G. J. *J. Chem. Soc., Perk. Trans. 1* **2001**, 2393. (c) O'Brien, C. J.; Lavigne, F.; Coyle, E. E.; Holohan, A. J.; Doonan, B. J. *Chem. Eur. J.* **2013**, 19, 5854.
- (9) Lipshutz, B. H.; Ghorai, S.; Boskovic, Z. V. *Tetrahedron* **2008**, 64, 6949.
- (10) (a) Abarbri, M.; Parrain, J.-L.; Duchene, A. *Synth. Commun.* **1998**, 28, 239. (b) Naskar, D.; Roy, S. *Tetrahedron* **2000**, 1369.
- (11) Schabel, T.; Plietker, B. *Chem. Eur. J.* **2013**, 19, 6938.
- (12) Wimmer, L.; Schönbauer, D.; Pakfeifer, P.; Schöffmann, A.; Khom, S.; Hering, S.; Mihovilovic, M. D. *Org. Biomol. Chem.* **2015**, 13, 990.
- (13) Chandrasekhar, S.; Venkat Reddy, M.; Srinivasa Reddy, K.; Ramarao, C. *Tetrahedron Lett.* **2000**, 41, 2667.
- (14) (a) Frébault, F.; Luparia, M.; Oliveira, M. T.; Goddard, R.; Maulide, N. *Angew. Chem. Int. Ed.* **2010**, 49, 5672. (b) Luparia, M.; Audisio, D.; Maulide, N. *Synlett* **2011**, 735. (c) Luparia, M.; Oliveira, M. T.; Audisio, D.; Frébault, F.; Goddard, R.; Maulide, N. *Angew. Chem. Int. Ed.* **2011**, 50, 12631. (d) Audisio, D.; Luparia, M.; Oliveira, M. T.; Kluett, D.; Maulide, N. *Angew. Chem. Int. Ed.* **2012**, 51, 7314. (e) Souris, C.; Frébault, F.; Audisio, D.; Farès, C.; Maulide, N. *Synlett* **2013**, 24, 1286. (f) Souris, C.; Misale, A.; Chen, Y.; Luparia, M.; Maulide, N. *Org. Lett.* **2015**, 17, 4486. (g) Gutekunst, W. R.; Baran, P. S. *J. Am. Chem. Soc.* **2011**, 133, 19076. (h) Nistanaki, S.; Boralsky, L. A.; Pan, R. D.; Nelson, H. *ChemRxiv* **2018**, DOI: 10.26434/chemrxiv.7245167.v2.
- (15) For spontaneous ring opening of cyclobutenes, see: (a) Binns, F.; Hayes, R.; Ingham, S.; Saengchantara, S. T.; Turner, R. W.; Wallace, T. W. *Tetrahedron* **1992**, 48, 515. (b) Binns, F.; Hayes, R.;

- Hodgetts, K. J.; Saengchantara, S. T.; Wallace, T. W.; Wallis, C. J. *Tetrahedron* **1996**, *52*, 3631. (c) Sheldrake, H. M.; Wallace, T. W.; Wilson, C. P. *Org. Lett.* **2005**, *7*, 4233. (d) Ingham, S.; Turner, R. W.; Wallace, T. J. *Chem. Soc., Chem. Commun.* **1985**, 1664.
- (16) Magnesium turnings (1.60 mmol, 3.0 equiv.) were suspended in a flamed-dried Schlenk tube containing anhydrous THF (3 mL) which was previously flushed with argon. A solution of an aryl-bromide in dry THF (1.60 mmol, 1.0 equiv. in 2 mL) was added dropwise under stirring. The mixture was stirred at room temperature until the magnesium was fully solubilized. LiCl (3.20 mmol, 6 equiv.) was flame-dried in another Schlenk tube under vacuum. After flushing with argon and cooling to room temperature CuCN (1.60 mmol, 3 equiv.) was added to the solid, suspended in anhydrous THF (5 mL), and stirred for 20 min at room temperature. The green solution was cooled down to -40°C , followed by an addition of the Grignard (1.60 mmol, 3 equiv.) and further stirred for 40 min. Afterwards, the reaction mixture was cooled down to -78°C . After stirring for 15 min, an ethereal solution of lactone **2** (0.53 mmol, 1 equiv.) was added, and the reaction mixture was allowed to warm up slowly to -30°C . The reaction was quenched by the dropwise addition of aqueous HCl (1 M, 5 mL), and the obtained solution was warmed up quickly to room temperature (the dry ice bath was removed). Upon dilution with EtOAc (10 mL), the reaction mixture was extracted and washed with aqueous HCl (1 M) whereby the aqueous layer was neutralized with sufficient NaOH (5 M) in order to deprotonate the formed hydrocyanic acid. The organic layer was dried over Na_2SO_4 , filtered, and the solvent was removed under reduced pressure. Purification of the crude acid was performed *via* column chromatography (EtOAc/heptane/acetic acid 10%:89%:1%) to yield the 4-aryl-substituted pentadienoic acid as a solid.
- (17) Copper(I) iodide lead only to messy reactions with traces of desired product.
- (18) In a flame-dried Schlenk tube the carboxylic acid **4** (0.20 mmol, 1.0 equiv.) was dissolved in anhydrous DCM (1 mL), followed by the addition of thionyl chloride (0.40 mmol, 2.0 equiv.). The reaction was stirred at room temperature until the solid dissolved completely. Heating to reflux can be applied for a faster conversion. Thereafter, volatiles were removed under reduced pressure, and the flask was flushed with argon again (an argon-filled balloon was attached to the rotary evaporator). To the resulting residue anhydrous DCM was added (1 mL), and the amine was added dropwise (0.40 mmol, 2.0 equiv.). Afterwards, the solution was stirred for 1 h at room temperature. Finally, the reaction mixture was quenched and washed with saturated aqueous NaHCO_3 solution, and the aqueous layer was extracted 3 times with DCM. The combined organic layer was dried over Na_2SO_4 , filtered, and volatiles were removed under reduced pressure to yield the pure amide. Further purification *via* flash chromatography gave the desired amide. (Gradient: EtOAc/heptane 10:90 to EtOAc/heptane 60:40).
- (19) The crude NMR of **4b**, **4c**, and **4d** suggested that traces (<5% NMR yield) of another geometrical isomer was present which was easily separated by column chromatography. We were not able to find this side product after purification.
- (20) Kotretsou, S. I.; Georgiadis, M. P. *Org. Prep. Proced. Int.* **2000**, *32*, 161.
- (21) Wani, N. A.; Singh, S.; Farooq, S.; Shankar, S.; Koul, S.; Khan, I. A.; Rai, R. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 4174.
- (22) Wang, Z.; He, Z.; Zhang, L.; Huang, Y. *J. Am. Chem. Soc.* **2018**, *140*, 735.
- (23) Yasir, A.; Ishtiaq, S.; Jahangir, M.; Ajajib, M.; Salar, U.; Khan, K. M. *Med. Chem.* **2018**, *14*, 269.
- (24) Shen, Y.; Zhou, Y. *J. Chem. Soc., Perkin Trans. 1* **1992**, *22*, 3081.
- (25) Spectroscopic data of **1**,²⁰ **2**,^{14f} **4a**,²¹ **4b**,²² **4c**,^{10b} **4d**,^{10b} **5aa**,²³ **5ba**,²⁴ **5ca**,⁴ **5bb**,¹² and **5cb**¹² matches previously reported spectra.
- (26) **Characterization of 5da**
 ^1H NMR (400 MHz, CDCl_3): δ = 7.37 (ddd, J = 14.6, 11.1, 0.6 Hz, 1 H), 7.23 (d, J = 5.0 Hz, 1 H), 7.07 (d, J = 3.5 Hz, 1 H), 7.01–6.93 (m, 2 H), 6.70 (dd, J = 15.3, 11.2 Hz, 1 H), 6.45 (d, J = 14.7 Hz, 1 H), 3.60 (d, J = 24.6 Hz, 2 H), 3.52 (s, 2 H), 1.71–1.55 (m, 7 H). ^{13}C NMR (151 MHz, CDCl_3): δ = 165.40, 142.03, 131.18, 127.97, 127.90, 126.80, 125.99, 120.62, 47.04, 43.38, 26.86, 25.74, 24.78. HRMS (ESI): m/z calcd for $[\text{M} + \text{Na}]^+$: 270.0923; found: 270.0926. ATR-FTIR: 3001, 2935, 2854, 2237, 1631, 1588, 1513, 1435, 1358, 1295, 1254, 1222, 1190, 1133, 1118, 1043, 1018, 990, 953, 907, 872, 853, 833, 805, 726, 644, 575, 543, 532 cm^{-1} .