


# Air-Stable Secondary Phosphine Oxides for Nickel-Catalyzed Cross-Couplings of Aryl Ethers by C–O Activation

Debasish Ghorai<sup>a,b</sup>Joachim Loup<sup>a</sup>Giuseppe Zanon<sup>b</sup>Lutz Ackermann<sup>\*a</sup> 

<sup>a</sup> Institut für Organische und Biomolekulare Chemie, Georg-August-Universität, Tammannstraße 2, 37077 Göttingen, Germany  
Lutz.Ackermann@chemie.uni-goettingen.de

<sup>b</sup> Department of Chemistry, University of Pavia, Viale Taramelli 10, 27100 Pavia, Italy

Published as part of the 30 Years SYNLETT – Pearl Anniversary Issue



Received: 02.12.2018

Accepted after revision: 06.01.2019

Published online: 15.01.2019

DOI: 10.1055/s-0037-1611663; Art ID: st-2018-b0782-I

License terms: 


**Abstract** Air- and moisture-stable secondary phosphine oxides (SPOs) enabled nickel-catalyzed Kumada–Corriu cross-couplings of various arylmethyl ethers at room temperature by challenging C–O activation.

**Key words** C–O activation, arylation, cross-coupling, secondary phosphine oxide, nickel

Transition-metal-catalyzed cross-coupling reactions have emerged as a uniquely powerful tool for the assembly of substituted biaryl motifs.<sup>1</sup> Thus far, these cross-couplings have heavily relied on aryl halides as electrophilic coupling reagents. In contrast, easily accessible phenol-based electrophiles have recently undergone a renaissance as attractive alternatives.<sup>2</sup> On the basis of Wenkert's early studies from 1979,<sup>3</sup> the considerable potential of phenol-derived substrates has only recently been fully recognized. Thus, versatile cross-couplings have been realized with challenging carbamates, carbonates, sulfamates, silyloxyarenes, esters and ethers, among others, prominently featuring nickel catalysis.<sup>4</sup> Generally, these nickel catalysts largely require electron-rich tertiary phosphines as stabilizing ligands to guarantee efficacy in the key C–O bond scission.<sup>4</sup> Unfortunately, these electron-rich tertiary phosphines are usually highly air-sensitive, with a documented half-life for the aerobic oxidation of tri-*t*-butyl-phosphine of a few minutes.<sup>5</sup>

The (heteroatom-substituted) secondary phosphine oxides (HA)SPOs represent uniquely powerful ancillary preligands for metal catalysis because of their unique features, including the air- and moisture-stable nature, among oth-


ers.<sup>6</sup> Notably, air-stable SPOs undergo a self-assembly process in the presence of transition metals to generate a monoanionic bidentate chelate coordination environment (Scheme 1, a).<sup>6</sup> While Ackermann and others have unraveled the considerable potential of SPO complexes towards a wealth of efficient cross-coupling reactions with various aryl halides,<sup>7</sup> the possibility of employing air-stable SPO preligands for more challenging C–O activations with aryl ethers has thus far proven elusive. Within our program on sustainable transition-metal-catalyzed transformations<sup>8</sup> and selective C–O activation,<sup>9</sup> we hence became attracted to probing the unprecedented use of air-stable SPOs preligands for cross-couplings with easily available aryl ethers, the result of which we report herein. Notable features of our findings include (i) air- and moisture-stable SPOs for efficient C–O activations, (ii) earth-abundant nickel catalysis, and (iii) exceedingly mild reaction conditions at room temperature (Scheme 1, b).



**Scheme 1** (a) Self-assembly with SPOs, (b) nickel/SPO-catalyzed C–O activation

We initiated our studies by probing reaction conditions for the envisioned cross-coupling of ether **1a** with Ni(acac)<sub>2</sub> and Ph<sub>2</sub>P(O)H (**L1**) in toluene at a room temperature of 23 °C (Table 1, entry 1). Among a variety of preligands and solvents, the electron-rich HASPO **L7** as well as (*n*-Bu)<sub>2</sub>P(O)H (**L8**) and THF gave optimal results, respectively (entries 2–13). NiCl<sub>2</sub>(DME) proved to be most effective (entries 14–17). It is noteworthy that under otherwise identical reaction conditions, the bidentate ligand dppp featured a significantly inferior performance (entry 18). A control experiment verified the essential role of the nickel catalyst (entry 19).

**Table 1** Optimization of the Nickel/SPO-Catalyzed C–O Activation of Ether **1a**<sup>a</sup>



Entry	Ni Catalyst	SPO	Solvent	Yield (%)
1	Ni(acac) <sub>2</sub>	<b>L1</b>	toluene	10
2	Ni(acac) <sub>2</sub>	<b>L2</b>	toluene	12
3	Ni(acac) <sub>2</sub>	<b>L3</b>	toluene	25
4	Ni(acac) <sub>2</sub>	<b>L4</b>	toluene	35
5	Ni(acac) <sub>2</sub>	<b>L5</b>	toluene	23
6	Ni(acac) <sub>2</sub>	<b>L6</b>	toluene	50
7	Ni(acac) <sub>2</sub>	<b>L6</b>	THF	64
8	Ni(acac) <sub>2</sub>	<b>L1</b>	THF	15
9	Ni(acac) <sub>2</sub>	<b>L5</b>	THF	21
10	Ni(acac) <sub>2</sub>	<b>L3</b>	THF	60
11	Ni(acac) <sub>2</sub>	<b>L4</b>	THF	48
12	Ni(acac) <sub>2</sub>	<b>L7</b>	THF	69
13	Ni(acac) <sub>2</sub>	<b>L8</b>	THF	83
14	Ni(OTf) <sub>2</sub>	<b>L8</b>	THF	53
15	NiBr <sub>2</sub>	<b>L8</b>	THF	n.r.


Entry	Ni Catalyst	SPO	Solvent	Yield (%)
<b>16</b>	<b>NiCl<sub>2</sub>(DME)</b>	<b>L8</b>	<b>THF</b>	<b>90</b>
17	NiCl <sub>2</sub> (DME)	<b>L8</b>	THF	68 <sup>b</sup>
18	NiCl <sub>2</sub> (DME)	dppp	THF	39 <sup>c</sup>
19	–	<b>L8</b>	THF	n.r.

<sup>a</sup> Reaction conditions: **1a** (0.50 mmol), *p*-TolMgBr (0.75 mmol), [Ni] (5.0 mol%), (HA)SPO (10 mol%), solvent (1.5 mL), 23 °C, 16 h; yield of isolated product given; n.r. = no reaction.

<sup>b</sup> SPO **L8** (5.0 mol%).


<sup>c</sup> dppp (5.0 mol%).

Having the optimized reaction conditions for the nickel/SPO-catalyzed C–O activation in hand, we tested its versatility with a representative set of ethers **1** (Scheme 2). Thus, a variety of naphthyl ethers **1** were identified as viable substrates for the Kumada–Corriu cross-coupling to deliver the desired products **2** with high catalytic efficacy. Notably, the nickel catalyst derived from the air-stable SPO **L8** even proved amenable to the chemoselective synthesis of biaryl **2b** and the sterically congested mesityl nucleophiles with comparable levels of activity (**2d** and **2i**).



**Scheme 2** Scope of SPO/nickel-catalyzed C–O activation; <sup>a</sup> with NiCl<sub>2</sub>(DME) (10 mol%) and **L8** (20 mol%)

Based on our previous literature reports,<sup>6c-d,10</sup> the working mode of the air-stable SPO-enabled C–O activation is suggested to initially involve the formation of complex **3** through self-assembly, along with the subsequent C–O activation by the key hetero-bimetallic intermediate **4** (Scheme 3).



**Scheme 3** Plausible working mode of SPOs for C–O activation

In summary, we have reported on the first use of air-stable secondary phosphine oxides (SPOs) for challenging cross-couplings of aryl ethers by C–O activation.<sup>11</sup> Thus, in situ generated nickel catalysts enabled efficient Kumada–Corriu arylations of naphthyl ethers at room temperature, even when using sterically hindered aryl nucleophiles.

## Funding Information

Generous support by the European Research Council under the European Community's Seventh Framework Program (FP7 2007–2013)/ERC Grant agreement no. 307535, and the Regione Lombardia – Cariplo Foundation is gratefully acknowledged.

## Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/s-0037-1611663>.

## References and Notes

- (1) (a) Cherney, A. H.; Kadunce, N. T.; Reisman, S. E. *Chem. Rev.* **2015**, *115*, 9587. (b) Johansson Seechurn, C. C.; Kitching, M. O.; Colacot, T. J.; Snieckus, V. *Angew. Chem. Int. Ed.* **2012**, *51*, 5062. (c) *Modern Arylation Methods*, 2nd ed; Ackermann, L., Ed.; Wiley-VCH: Weinheim, **2009**. (d) Beller, M.; Bolm, C. *Transition Metals for Organic Synthesis*; Wiley-VCH: Weinheim, **2004**. (e) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457.
- (2) Kozhushkov, S. I.; Potukuchi, H. K.; Ackermann, L. *Catal. Sci. Technol.* **2013**, *3*, 562.
- (3) (a) Wenkert, E.; Michelotti, E. L.; Swindell, C. S.; Tingoli, M. *J. Org. Chem.* **1984**, *49*, 4894. (b) Wenkert, E.; Michelotti, E. L.; Swindell, C. S. *J. Am. Chem. Soc.* **1979**, *101*, 2246.
- (4) Representative reviews: (a) Tobisu, M.; Chatani, N. *Acc. Chem. Res.* **2015**, *48*, 1717. (b) Su, B.; Cao, Z.-C.; Shi, Z.-J. *Acc. Chem. Res.* **2015**, *48*, 886. (c) Tollefson, E. J.; Hanna, L. E.; Jarvo, E. R. *Acc. Chem. Res.* **2015**, *48*, 2344. (d) Tasker, S. Z.; Standley, E. A.; Jamison, T. F. *Nature* **2014**, *509*, 299. (e) Cornella, J.; Zarate, C.; Martin, R. *Chem. Soc. Rev.* **2014**, *43*, 8081. (f) Li, B. J.; Yu, D. G.; Sun, C. L.; Shi, Z. J. *Chem. Eur. J.* **2011**, *17*, 1728. (g) Rosen, B. M.; Quasdorf, K. W.; Wilson, D. A.; Zhang, N.; Resmerita, A.-M.; Garg, N. K.; Percec, V. *Chem. Rev.* **2011**, *111*, 1346. (h) Yu, D.-G.; Li, B.-J.; Shi, Z.-J. *Acc. Chem. Res.* **2010**, *43*, 1486. Selected examples: (i) Wang, T.-H.; Ambre, R.; Wang, Q.; Lee, W.-C.; Wang, P.-C.; Liu, Y.; Zhao, L.; Ong, T.-G. *ACS Catal.* **2018**, *8*, 11368. (j) Cao, Z.-C.; Luo, Q.-Y.; Shi, Z.-J. *Org. Lett.* **2016**, *18*, 5978. (k) Zhang, J.; Xu, J.; Xu, Y.; Sun, H.; Shen, Q.; Zhang, Y. *Organometallics* **2015**, *34*, 5792. (l) Iglesias, M. J.; Prieto, A.; Nicasio, M. C. *Org. Lett.* **2012**, *14*, 4318. (m) Xie, L.-G.; Wang, Z.-X. *Chem. Eur. J.* **2011**, *17*, 4972. (n) Dankwardt, J. W. *Angew. Chem. Int. Ed.* **2004**, *43*, 2428. For general reviews on nickel catalyzed transformations, see: (o) Castro, L. C. M.; Chatani, N. *Chem. Lett.* **2015**, *44*, 410. (p) Yamaguchi, J.; Muto, K.; Itami, K. *Eur. J. Org. Chem.* **2013**, *19*. (q) Nakao, Y. *Chem. Rec.* **2011**, *11*, 242, and references cited therein.
- (5) Netherton, M. R.; Fu, G. C. *Org. Lett.* **2001**, *3*, 4295.
- (6) Select reviews: (a) Hérault, D.; Nguyen, D. H.; Nuel, D.; Buono, G. *Chem. Soc. Rev.* **2015**, *44*, 2508. (b) Shaikh, T. M.; Weng, C.-M.; Hong, F.-E. *Coord. Chem. Rev.* **2012**, *256*, 771. (c) Ackermann, L. *Isr. J. Chem.* **2010**, *50*, 652. (d) Ackermann, L. *Synthesis* **2006**, 1557. (e) Dubrovina, N. V.; Börner, A. *Angew. Chem. Int. Ed.* **2004**, *43*, 5883.
- (7) (a) Ghorai, D.; Müller, V.; Keil, H.; Stalke, D.; Zanoni, G.; Tkachenko, B. A.; Schreiner, P. R.; Ackermann, L. *Adv. Synth. Catal.* **2017**, *359*, 3137. (b) Hu, C.-Y.; Chen, Y.-Q.; Lin, G.-Y.; Huang, M.-K.; Chang, Y.-C.; Hong, F.-E. *Eur. J. Inorg. Chem.* **2016**, 3131. (c) Cano, I.; Tschan, M. J. L.; Martínez-Prieto, L. M.; Philippot, K.; Chaudret, B.; van Leeuwen, P. W. N. M. *Catal. Sci. Technol.* **2016**, *6*, 3758. (d) Wellala, N. P.; Guan, H. *Org. Biomol. Chem.* **2015**, *13*, 10802. (e) Cano, I.; Huertos, M. A.; Chapman, A. M.; Buntkowsky, G.; Gutmann, T.; Groszewicz, P. B.; van Leeuwen, P. W. N. M. *J. Am. Chem. Soc.* **2015**, *137*, 7718. (f) Ackermann, L.; Kapdi, A. R.; Fenner, S.; Kornhaass, C.; Schulzke, C. *Chem. Eur. J.* **2011**, *17*, 2965. (g) Ackermann, L.; Potukuchi, H. K.; Kapdi, A. R.; Schulzke, C. *Chem. Eur. J.* **2010**, *16*, 3300. (h) Ackermann, L.; Vicente, R.; Hofmann, N. *Org. Lett.* **2010**, *11*, 4274. (i) Achard, T.; Giordano, L.; Tenaglia, A.; Gimbert, Y.; Buono, G. *Organometallics* **2010**, *29*, 3936. (j) Christiansen, A.; Selent, D.; Spannenberg, A.; Baumann, W.; Franke, R.; Börner, A. *Organometallics* **2010**, *29*, 3139. (k) Christiansen, A.; Li, C.; Garland, M.; Selent, D.; Ludwig, R.; Spannenberg, A.; Baumann, W.; Franke, R.; Börner, A. *Eur. J. Org. Chem.* **2010**, 2733. (l) Ackermann, L.; Barfüßer, S. *Synlett* **2009**, 808. (m) Yang, D. X.; Colletti, S. L.; Wu, K.; Song, M.; Li, G. Y.; Shen, H. C. *Org. Lett.* **2009**, *11*, 381. (n) Billingsley, K. L.; Buchwald, S. L. *Angew. Chem., Int. Ed. Engl.* **2008**, *47*, 4695. (o) Ackermann, L.; Born, R.; Spatz, J. H.; Meyer, D. *Angew. Chem. Int. Ed.* **2005**, *44*, 7216.
- (8) (a) Gandeepan, P.; Müller, T.; Zell, D.; Cera, G.; Warratz, S.; Ackermann, L. *Chem. Rev.* **2019**, in press; doi: 10.1021/acs.chemrev.8b00507. (b) Lorion, M. M.; Maïndan, K.; Kapdi, A. R.; Ackermann, L. *Chem. Soc. Rev.* **2017**, *46*, 7399. (c) Moselage, M.; Li, J.; Ackermann, L. *ACS Catal.* **2016**, *6*, 498. (d) Liu, W.; Ackermann, L. *ACS Catal.* **2016**, *6*, 3743.
- (9) (a) Sauermann, N.; Loup, J.; Kootz, D.; Berkessel, A.; Ackermann, L. *Synthesis* **2017**, 49, 3476. (b) Song, W.; Ackermann, L. *Angew. Chem. Int. Ed.* **2012**, *51*, 8251. (c) Ackermann, L.; Pospesch, J.; Potukuchi, H. K. *Org. Lett.* **2012**, *14*, 2146. (d) Ackermann, L.;

Althammer, A.; Born, R. *Angew. Chem. Int. Ed.* **2006**, *45*, 2619.  
(e) Moselage, M.; Sauermann, N.; Richter, S. C.; Ackermann, L. *Angew. Chem. Int. Ed.* **2015**, *54*, 6352.

(10) Ackermann, L. *Synlett* **2007**, 507.

(11) **Representative Experimental Procedure and Characterization Data**

A mixture of 2-methoxynaphthalene (**1a**) (79 mg, 0.5 mmol), [NiCl<sub>2</sub>(DME)] (6.0 mg, 0.025 mmol, 5.0 mol%), and **L8** (8.0 mg, 0.05 mmol, 10.0 mol%) was stirred in THF (1.5 mL) for 2 min at ambient temperature under N<sub>2</sub>. Then, *p*-TolMgBr (1.0 M in THF, 0.75 mL, 0.75 mmol) was added, and the resulting solution was stirred for 16 h at ambient temperature. To the reaction was added aqueous HCl (1 M, 5 mL) and then EtOAc (5 mL), and the separated aqueous phase was extracted with EtOAc (2 × 5 mL).

The combined organic layers were dried with anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated in vacuo. The remaining residue was purified by column chromatography on silica gel (*n*-hexane) to yield **2a** (98 mg, 90%) as a colorless solid. Mp 93–95 °C. IR (ATR): 3054, 3024, 1501, 1351, 893, 856, 811, 748 cm<sup>-1</sup>. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ = 8.14 (d, *J* = 1.4 Hz, 1 H), 8.03–7.93 (m, 3 H), 7.85 (dd, *J* = 8.5, 1.9 Hz, 1 H), 7.74 (d, *J* = 8.1 Hz, 2 H), 7.64–7.54 (m, 2 H), 7.40 (dd, *J* = 8.5, 0.6 Hz, 2 H), 2.53 (s, 3 H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>): δ = 138.5 (C<sub>q</sub>), 138.3 (C<sub>q</sub>), 137.2 (C<sub>q</sub>), 133.8 (C<sub>q</sub>), 132.5 (C<sub>q</sub>), 129.6 (CH), 128.4 (CH), 128.2 (CH), 127.7 (CH), 127.3 (CH), 126.3 (CH), 125.8 (CH), 125.6 (CH), 125.5 (CH), 21.2 (CH<sub>3</sub>). MS (EI): *m/z* (relative intensity) = 218 [M]<sup>+</sup> (100), 217 (41), 202 (35). HRMS (EI): *m/z* [M]<sup>+</sup> calcd for [C<sub>17</sub>H<sub>14</sub>]<sup>+</sup>: 218.1096; found: 218.1094.