

Tuberculum Meningioma: Orbitopterional Approach

Alexander Yang¹ Mohammed Aref¹ A. Samy Youssef¹

¹Department of Neurosurgery, University of Colorado, Aurora, Colorado, United States

Address for correspondence Samy Youssef, MD, PhD, Department of Neurosurgery, University of Colorado, Mail Stop C307, 12631 East 17th Avenue, Aurora, CO 80045, United States
(e-mail: Samy.Youssef@UCDenver.edu).

J Neurol Surg B 2018;79(suppl S2):S219–S220.

Abstract

This is a case of an extensive tuberculum sella meningioma involving the circle of Willis down to the basilar artery that presented with bilateral visual loss worse on the right than left side. A one-piece right orbitopterional approach along the worse eye was used to gain access to the three cranial fossae. The orbitotomy facilitates access to the midline structures and contralateral base of the tumor with minimal brain retraction. Tumor resection is initiated by first identifying the tumor capsule, followed by piecemeal debulking via ultrasonic aspiration. Early decompression of the ipsilateral optic nerve was performed. Gross total resection of the tumor was achieved through multiple windows as follows: prechiasmatic, opticocarotid, and carotid oculomotor. Sharp dissection is performed around critical neurovascular structures to reduce strain and vascular injury. The circle of Willis including the small perforators was completely preserved. Postoperative examination at follow-up demonstrated improvement in vision less on the right side and resolution of postoperative partial third nerve palsy. The link to the video can be found at: <https://youtu.be/XfEh8CjkvA0>.

Keywords

- ▶ orbitopterional
- ▶ meningioma
- ▶ vision
- ▶ preservation
- ▶ tuberculum sella
- ▶ tuberculum
- ▶ sella

Conflict of Interest

None.



www.thieme.com/skullbasevideos

www.thieme.com/jnlsbvideos

received
October 20, 2017
accepted after revision
November 29, 2017
published online
January 16, 2018

DOI <https://doi.org/10.1055/s-0037-1618599>.
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms






Fig. 1 Preoperative, postoperative, and 6-month follow-up imaging.




Fig. 2 Anterior communicating artery (ACoA) complex demonstrating vascular segments of the Left A1, Left A2, Right A1, and Right A2 vessels, as well as perforating vessels.