

# Tuberculum Sellae Meningioma Resection: Technical Nuances on the Frontopterional Approach

Oriela Rustemi<sup>1</sup> Renato Scienza<sup>1</sup> Alessandro Della Puppa<sup>1</sup>

<sup>1</sup>Department of Neurosurgery, Padua University Hospital, Padua, Italy

**Address for correspondence** Oriela Rustemi, MD, Department of Neurosurgery, Padua University Hospital, Azienda Ospedaliera di Padova, via Giustiniani, 2-35128, Padova, Italy (e-mail: orielarustemi@libero.it).

J Neurol Surg B 2018;79(suppl S2):S225–S226.

## Abstract

Tuberculum sellae meningioma remains a surgical challenge. Deep location of tumor, vascular and nerve encasement, and pituitary stalk involvement are the main technical issues. The frontopterional approach represents a natural, simple, and elegant approach to this area enabling surgeon to have a direct control on all anatomical structures. A 42-year-old woman was referred with a delayed diagnosis of tuberculum sellae meningioma due to the presence of HLA-B27-associated uveitis. She presented with 1/10 visual acuity in the left eye and no right visual function. A left frontopterional craniotomy was performed. Visual function improved postoperatively. The video illustrates the cisternal anatomy via pterional approach.

The link to the video can be found at: <https://youtu.be/Hmbf5bt7A64>.

## Keywords

- ▶ tuberculum sellae meningioma
- ▶ pterional approach
- ▶ cisternal anatomy
- ▶ optic canal

## Conflict of Interest

None.

## Funding

None.



[www.thieme.com/skullbasevideos](http://www.thieme.com/skullbasevideos)

[www.thieme.com/jnlsbvideos](http://www.thieme.com/jnlsbvideos)


received  
July 8, 2017  
accepted  
December 12, 2017  
published online  
January 15, 2018

DOI <https://doi.org/10.1055/s-0037-1620245>.  
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG  
Stuttgart · New York

License terms





**Fig. 1** (A) Preoperative MRI shows a contrast-enhanced tuberculum sellae lesion, suspicious of meningioma. (B) The left optic nerve (II c.n.) is evidenced through a left pterional approach. The II c.n. is compressed by the meningioma. (C) The left optic canal is unroofed and the sign of optic nerve compression is indicated by blue asterisks (\*). (D) Intraoperative image after removing the left part of the meningioma showing the left optic nerve, internal carotid artery (ICA) and posterior communicating artery (PCom). (E) Intraoperative image after the removal of the meningioma showing bilateral optic nerves and ICAs, and the tuberculum meningioma's base. (F) Final intraoperative view showing bilateral oculomotor nerve (III c.n.), left II c.n., left ICA, and the conserved pituitary stalk. (G) Postoperative CT shows the complete removal of the meningioma. CT, computed tomography; MRI, magnetic resonance imaging.