

Gross Total Resection of Chordoid Glioma of the Third Ventricle via Anterior Interhemispheric Transcallosal Transforaminal Approach at Two Stages

Ihsan Dogan¹ Melih Ucer¹ Mustafa Kemal Başkaya¹

¹ Department of Neurological Surgery, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, United States

J Neurol Surg B 2018;00:S281–S282.

Address for correspondence Mustafa Kemal Başkaya, MD, Department of Neurological Surgery, University of Wisconsin School of Medicine and Public Health, 600 Highland Avenue, K4/8 CSC, Box 8660, Madison, WI 53792, United States (e-mail: baskaya@neurosurgery.wisc.edu).

Abstract

Suprasellar tumors in particular tumors located in the retrochiasmatic area and anterior third ventricle are challenging cases in terms of optimal surgical exposure. Several approaches have been described including transsylvian, translamina terminalis, endoscopic endonasal, and anterior interhemispheric. Each approach has advantages and disadvantages. In this video, we present a case of retrochiasmatic anterior third ventricular tumor that was operated via anterior interhemispheric transcallosal transforaminal approach. The patient is a 42-year-old female who presented with sudden onset of severe headache and depressed level of consciousness. Computed tomography (CT) scan of the head showed a hemorrhage in the third ventricle and suprasellar cisterns. CT angiogram and magnetic resonance imaging (MRI) confirmed diagnosis of hemorrhagic mass lesion in the third ventricle. Upon further questioning of her family, we found out that she was having excessive urination and short-term memory problems for last 2 weeks. First, ventriculostomy was placed for obstructive hydrocephalus. She then underwent surgical resection via anterior interhemispheric transcallosal transforaminal approach. Foramen of Monro was enlarged by performing transchoroidal dissection. Using transforaminal route, tumor was resected. Due to the narrow surgical corridor and high vascularity of the tumor, decision was made to come back at a second stage. Using same surgical approach, in the second stage, gross total

Keywords

- suprasellar tumor
- interhemispheric transcallosal approach
- third ventricle tumor
- chordoid glioma



www.thieme.com/skullbasevideos

www.thieme.com/jnlsvideos

received
October 17, 2017
accepted
December 3, 2017
published online
February 16, 2018

DOI <https://doi.org/10.1055/s-0038-1625967>.
ISSN 2193-6331.

© 2018 Georg Thieme Verlag KG
Stuttgart · New York

License terms



resection was performed. Postoperative MRI confirmed gross total resection. Histopathology was chordoid glioma of the third ventricle. She made excellent recovery with persistent diabetes insipidus. Currently, she is completing radiation therapy. In this video, we demonstrate techniques and pitfalls of anterior interhemispheric transcallosal approach to anterior third ventricular tumor.

The link to the video can be found at: <https://youtu.be/CI5c6Zup8sY>.

Conflict of Interest

None.