People, Trends and Views in Chemical Synthesis

2020/08

Synthesis of a Mechanically Planar Chiral **Rotaxane Ligand for Enantioselective Catalysis**

Highlighted article by A. W. Heard, S. M. Goldup



enantiomeric mechanically planar chiral rotaxanes

Contact

Your opinion about Synform is welcome, please correspond if you like: marketing@thieme-chemistry.com



Dear Readers,

Exactly one year ago my Editorial for the August 2019 issue of SYNFORM was centered on the Spring American Chemical Society National Meeting & Exposition held in Orlando, FL, USA, from March 31 through April 4, 2019, which I attended sponsored by Thieme Chemistry. This year not just the ACS Meetings but almost every conference on this planet has gone virtual, including the Thieme Chemistry Editorial Board Meeting 2020 - which is better than no meeting at all, but - oh my goodness - I am missing so much the real thing... I am not saying you can't do useful stuff on videoconference - actually if it wasn't for these invaluable online tools we would have all been doomed during the lockdown – but it's only a pale mimic of an actual meeting, an ectoplasm of real life. I have never been a lover of crowds, but I cannot even explain how much I am craving a good old-style meeting with real people, in person, possibly without those lifesaving – but nevertheless cumbersome – face masks... Meanwhile, I am afraid my only consolation is to look at the mask-free pictures of this issue's authors... and enjoy the great science tales they are telling, of course! Starting with the amazing molecular machines developed by S. Goldup (UK) and continuing with the informative Young Career Focus interview with S. G. Modha (India). The next article covers a new catalytic process for the efficient chlorination of (hetero)arenes published by N. Jiao (P. R. China), while the closing article explores the three-component aminomethylation of α -diazo ketones developed by D. Xing and W. Hu (P. R. China).

Stay safe and enjoy your reading!!

Another fonde

In this issue

Literature Coverage Synthesis of a Mechanically Planar Chiral Rotaxane Ligand for Enantioselective Catalysis
Young Career Focus Young Career Focus: Dr. Sachinkumar G. Modha (Uka Tarsadia University, India)
Literature Coverage DMSO-Catalyzed Late-Stage Chlorination of (Hetero)arenes
Literature Coverage Enantioselective Three-Component Aminomethyl- ation of α-Diazo Ketones with Alcohols and 1,3,5-Tri- azines
Coming soon

Contact

If you have any questions or wish to send feedback, please write to Matteo Zanda at: synform@outlook.com

Synthesis of a Mechanically Planar Chiral Rotaxane Ligand for Enantioselective Catalysis

A110

Chem 2020, 6, 994–1006

Rotaxanes are interlocked molecules in which a molecular ring is threaded onto a dumbbell-shaped axle which bears end groups too large to fit through the cavity of the ring.¹ Professor Steve Goldup from the University of Southampton (UK) explained: "This arrangement prevents the ring and axle from separating without a covalent bond being broken and so, even though there is no covalent bond between the ring and the axle, rotaxanes are molecular rather than supramolecular species. Rotaxanes are most famous as components of molecular machines,² which function by exploiting the controlled movement of the ring along the axle. However, there is a growing interest in how the chemical properties of rotaxanes, and other interlocked molecules, can be exploited to solve chemical problems, for example in catalysis,³ sensing,⁴ materials science⁵ and medicinal chemistry."⁶ As part of a research project focused on investigating the unusual stereochemical properties of interlocked molecules, Professor Steve Goldup and postgraduate student Andrew Heard developed a rotaxane-based gold complex that carries out an enantioselective cyclopropanation reaction.

Rotaxanes have the unusual property that they can exhibit molecular chirality even if the ring and axle are achiral (Scheme 1, a).⁷ Professor Goldup explained that this was first discussed over 50 years ago,8 but until very recently it was extremely hard to make large quantities of such "mechanically planar chiral" rotaxanes as there were no methods that did not require HPLC separation of the enantiomers. In previous work, the Goldup research group took the first steps towards solving this problem by developing methods in which a covalent chiral auxiliary is included in the synthesis to allow the stereoisomers to be separated⁹ and even direct the stereoselective formation of the mechanical bond.¹⁰ Professor Goldup said: "We've been pushing hard to develop ways to make mechanically chiral rotaxanes for almost 10 years. In parallel, we have been investigating the use of rotaxanes as ligands in homogeneous catalysis. In this project, we brought these two ideas together to investigate whether a mechanically planar chiral rotaxane could provide a chiral environment for a gold-mediated reaction in order to generate the products enantioselectively."



Scheme 1 a) Schematic showing the construction of enantiomeric mechanically planar chiral rotaxanes from achiral components. b) Synthesis of mechanically planar chiral rotaxane gold complexes (R_{mp})-**5** and (S_{mp})-**5**.

Literature Coverage

To make rotaxane-based gold complexes 5, Andrew Heard used an active template copper-mediated alkyne-azide cycloaddition reaction (Scheme 1, b).11 "In this reaction, a Culion bound in the cavity of a bipyridine macrocycle (3) mediates the reaction of the alkyne (2) and azide ([S]-1) half-axles. Because the Cu¹ ion is held in the cavity of the ring, the new bonds, and thus the new axle, are formed through the ring. The key feature of the synthesis is that azide half-axle (S)-1 is chiral and enantiopure and so the product rotaxane can be formed as two diastereomers (S, S_{mp}) -4 and (S, R_{mp}) -4 (mp refers to the mechanically planar chiral stereogenic unit) that differ only in the orientation of the ring on the axle," said Andrew Heard, adding: "This was the hardest part of the project. Even though previous work in our group suggested one of the stereoisomers would be formed preferentially, I observed very little diastereoselectivity. Worse, the covalent stereocentre could be epimerised very easily. Ultimately, I was able to separate the diastereomers using silica gel chromatography, but I had to pre-saturate the column eluent with water to avoid epimerising the products and work quickly. It was an exciting day when I saw the HPLC analysis of my products that confirmed they were separated with excellent stereochemical purity and that my procedure was reproducible!" With the rotaxane diastereomers separated, the synthesis was completed by alkylation of the covalent stereocentre to give the separated rotaxane enantiomers, reduction of the phosphine oxide to give the gold-binding phosphine ligand, and coordination of AuCl to give pre-catalysts (R_{mn}) -**5** and (S_{mn}) -**5** in 98% ee.

"With pre-catalysts 5 in hand, Andrew then investigated whether they could be used in an Au-mediated cyclopropanation reaction first developed by Toste and co-workers (Scheme 2, a).¹² I'm often asked why we focused on this particular reaction. As with many things in research, it is partly by design and partly by chance," said Professor Goldup. He continued: "It is often hard to impose stereocontrol in gold-catalysed reactions because the ligand and bound substrate are disposed at 180° to one another around the gold ion.13 A number of years ago I was interested in the effect of the mechanical bond on the diastereoselectivity of Au-mediated reactions; the mechanical bond creates a very crowded three-dimensional space in which the Au¹ ion can be embedded and this might help solve the problem. Marzia Galli, a new PhD student in my group at the time, suggested this cyclopropanation reaction for her study because the products are formed as a mixture of cis- and trans-isomers. Because this reaction was so successful in our previous study,¹⁴ we decided to investigate whether chiral rotaxane complex 5 could carry out the cyclopropanation reaction enantioselectively." Excitingly, the first experiments with (R)-5 led to the formation of cyclopropane 8 in reasonable enantioselectivity and yield. "Our reaction conditions are similar to those reported by Toste and co-workers, except we include a Cu¹ additive which binds into the cavity of the bipyridine ring and prevents the bipyridine binding to the Au¹ ion and inhibiting the reaction," explained Professor Goldup. He remarked: "We have of course checked and the Cu¹ salt itself is not a catalyst for the reaction!" Importantly, when the opposite hand of the complex, (S)-5, was used, the opposite enantiomer of 8 was formed as the major product. "That was a relief, as it reassured me that the enantioselectivity was due to the configuration of the mechanical bond rather than an adventitious chiral impurity left over from the synthesis," said Andrew. Having demonstrated that (R)-5 can act as an enantioselective catalyst for the reaction to produce 8, the study was then expanded to other substrates (Scheme 2, b). Andrew Heard said: "The substrate scope didn't tell us much more than that the catalyst is relatively general for propargylic benzoates, which is obviously nice. The high point of this part of the study was that by putting bulky substituents on the benzene ring of the substrate, we could enhance the enantioselectivity of the reaction to generate cyclopropane 10 in 76% ee. The low point was finding out that pivaloyl esters are not tolerated by the catalyst; these substrates gave higher enantioselectivities with Toste's catalysts and we were hoping we would see the same but cyclopropane 12 was produced as a near-racemate. Nevertheless, comparing like-for-like, rotaxane-based complex (R)-5 gives comparable enantioselectivity to chiral covalent catalysts reported for the same reaction."

Goldup and Heard completed their study by carrying out computational modelling to try and provide some insight into how the structure of 5 led to the observed enantioselectivity. Professor Goldup said: "I am always very careful about how I describe this part of the study when I give lectures. Rotaxane **5** is relatively large and flexible, and this means that modelling its catalytic behaviour with a high degree of rigour is very hard and would be a significant project in its own right. The modelling we have done is very preliminary, as we make clear in the paper. It was only designed to probe the kind of interactions that may be occurring between the ligand framework and the substrate to impart stereocontrol on the reaction and provide a visual representation of how the mechanical bond can create a chiral environment for the reaction to take place within." The modelling found that the substrate and ligand framework can interact by weak Cu– π , CH– π and C–H hydrogen bonds and that the mechanical bond can indeed create a crowded, chiral environment around the substrate (Scheme 2, c). "Given the relative simplicity of our approach, the modelling and experimental data agree remarkably well but this has to be interpreted carefully," said Professor Goldup. "That



Scheme 2 a) Enantioselective reaction of propargylic ester **6** with styrene (**7**) mediated by (R_{mp}) -**5** to give cyclopropane **8** in 58% *ee*. b) Examples of substrates examined. c) Modelled (stick representation, left; spacefilling representation, right) transition state for the reaction leading to observed major enantiomer of **8** with selected intercomponent interactions highlighted.

said, our initial results suggest that modelling these kinds of complex systems is definitely possible."

Professor Goldup and his group are excited about the future of rotaxanes and other interlocked molecules as catalysts. Andrew Heard said: "Although the synthesis was challenging in this case, our methods are improving all the time. In fact, testing them in new, challenging scenarios has driven a lot of our recent progress in this area. Now we have demonstrated that chiral mechanical bonds are useful, we are even more motivated to make functionalised systems readily available." The Goldup group are now expanding their studies to investigate the effect of rotaxane structure on enantioselectivity in the gold-mediated cyclopropanation reaction, as well as investigating other catalytic systems. They are also looking at other applications of mechanically planar chiral rotaxanes in sensing and materials chemistry. "Mechanically planar chiral rotaxanes aren't the only type of interlocked molecule in which the mechanical bond provides the only source of stereochemistry," explained Professor Goldup (Figure 1).⁷ "Catenanes can display topological and axial chirality, and there are intriguing forms of dynamic stereochemistry that arise as a result of the motion of the two components. Although there are some interesting preliminary data,^{15,16} along with mechanical planar chirality, the applications of these chiral molecules have yet to be investigated properly because they have been too hard to make. We have already developed ways to make topologically chiral catenanes¹⁷ and are working on some of the other classes now. In conclusion, our aim is to find what mechanically chiral molecules are useful for and, in particular, what problems they can solve that are hard or impossible to solve in other ways at present."



Figure 1 Other examples of mechanically chiral molecules include the enantiomers of: a) topologically chiral catenanes, b) axially chiral catenanes, and c) co-conformationally mechanically planar chiral rotaxanes.



REFERENCES

(1) C. J. Bruns, J. F. Stoddart *The Nature of the Mechanical Bond – From Molecules to Machines*, John Wiley & Sons: Hoboken, **2017**.

(2) E. R. Kay, D. A. Leigh, F. Zerbetto Angew. Chem. Int. Ed. **2007**, 46, 72–191.

(3) N. Pairault, H. Zhu, D. Jansen, A. Huber, C. G. Daniliuc, S. Grimme, J. Niemeyer *Angew. Chem. Int. Ed.* **2020**, *59*, 5102–5107.

(4) J. Y. C. Lim, I. Marques, V. Félix, P. D. Beer J. Am. Chem. Soc. **2017**, 139, 12228–12239.

(5) Y. Okumura, K. Ito Adv. Mater. 2001, 13, 485–487.

(6) O. Koniev, I. Dovgan, B. Renoux, A. Ehkirch, J. Eberova, S. Cianférani, S. Kolodych, S. Papot, A. Wagner *Med. Chem. Commun.* **2018**, *9*, 827–830.

(7) E. M. G. Jamieson, F. Modicom, S. M. Goldup *Chem. Soc. Rev.* **2018**, *47*, 5266–5311.

(8) G. Schill *Catenanes, Rotaxanes and Knots*, Academic Press: New York, **1971**.

(9) R. J. Bordoli, S. M. Goldup J. Am. Chem. Soc. **2014**, 136, 4817–4820.

(10) M. A. Jinks, A. de Juan, M. Denis, C. J. Fletcher, M. Galli, E. M. G. Jamieson, F. Modicom, Z. Zhang, S. M. Goldup *Angew. Chem. Int. Ed.* **2018**, *57*, 14806–14810.

(11) M. Denis, S. M. Goldup *Nat. Rev. Chem.* **2017**, *1*, 0061;
DOI: 10.1038/s41570-017-0061.
(12) M. L. Ishenson, P. J. Carrie, S. T. Stellan, F. D. Tarta,

(12) M. J. Johansson, D. J. Gorin, S. T. Staben, F. D. Toste J. Am. Chem. Soc. **2005**, 127, 18002–18003.

About the authors



A. Heard

Andrew Heard studied for his MChem under Prof. Andrew Dove at the University of Warwick (UK) and began his PhD in the Goldup group in 2017 studying applications of mechanically planar chiral molecules.



2015, 54, 13545-13549.

Chem 2019, 5, 1512–1520.

sym10010020.

9133.

A113

Stephen Goldup studied for his MChem at the University of Oxford (UK), supervised by Prof. Sir Jack Baldwin and undertook his PhD at Imperial College (UK), supervised by Prof. Tony Barrett before completing his PDRA at the University of Edinburgh (UK) with Prof. David Leigh. In 2008 he began his independent career at Queen Mary, University of London (UK), first as a Leverhulme Trust Early Career Fellow and then Royal Society

Prof. S. Goldup

Fellow and Senior Lecturer. In 2014 the Goldup group moved to the University of Southampton (UK) where Steve works with an outstanding group of young scientists developing methods for the synthesis of challenging chiral interlocked molecules with catalytic, materials and sensing applications. In 2017 Steve was promoted to Professor and as of 2019 is a Royal Society Wolfson Research Fellow.

(13) W. Zi, F. D. Toste Chem. Soc. Rev. 2016, 45, 4567-4589.

(14) M. Galli, J. E. M. Lewis, S. M. Goldup Angew. Chem. Int. Ed.

(15) K. Hirose, M. Ukimi, S. Ueda, C. Onoda, R. Kano, K. Tsuda,

Y. Hinohara, Y. Tobe Symmetry 2018, 10, 20; DOI: 10.3390/

M. Baroncini, A. Credi J. Am. Chem. Soc. 2019, 141, 9129-

(17) M. Denis, J. E. M. Lewis, F. Modicom, S. M. Goldup

(16) S. Corra, C. de Vet, J. Groppi, M. La Rosa, S. Silvi,

Young Career Focus: Dr. Sachinkumar G. Modha (Uka Tarsadia University, India)

Background and Purpose. SYNFORM regularly meets young up-and-coming researchers who are performing exceptionally well in the arena of organic chemistry and related fields of research, in order to introduce them to the readership. This Young Career Focus presents Dr. Sachinkumar G. Modha (Uka Tarsadia University, India).

Biographical Sketch



Sachinkumar G. Modha obtained his M.Sc. (organopharmaceutical chemistry) from Saurashtra University, Rajkot, Gujarat (India). He then worked in an organization named Oxygen Healthcare Research Pvt. Ltd. (Contract Research Organization), Ahmedabad, Gujarat (India) for 15 months. After his short stay in the group of Prof. Anamik K. Shah at Saurashtra University he went to Belgium to pursue his doctoral stu-

Dr. S. G. Modha

dies at the University of Leuven (KU Leuven) under the supervision of Prof. Erik E. Van der Eycken. After completing his PhD in 2012, he went on to work at the University of Manchester (UK) under the supervision of Prof. Michael F. Greaney as a postdoctoral research associate. In 2016, Sachin obtained the prestigious Alexander von Humboldt postdoctoral fellowship to work with Prof. Thorsten Bach at the Technical University of Munich (Germany), where his stay was extended by one year. He took up a position at Uka Tarsadia University, Bardoli, Gujarat (India) in 2019 as assistant professor.

INTERVIEW

SYNFORM What is the focus of your current research activity?

Dr. S. G. Modha I am fascinated by photochemistry. Harnessing the energy from electromagnetic irradiation and using it for chemical transformations that are difficult, if not impossible, to achieve otherwise, is exciting and challenging at the same time. Lots of progress has been seen in the broad field of photochemistry over the last few decades, but still many unanswered questions remain, which motivates me to contribute in this field. At the same time, I am also interested in the development of new atom-economical routes towards interesting organic molecules. In particular, I like multicomponent reactions (general concept shown in Scheme 1) and reactions at room temperature and thus some of my efforts are focused on developing new reactions towards this goal.

SYNFORM When did you get interested in synthesis?

Dr. S. G. Modha After completion of my postgraduate studies, I worked in a Contract Research Organization for 15 months. Different projects with reactions of scale as small as 10 mg and as big as 1.4 kg fascinated me. In particular, the idea of synthesizing a molecule which has never been synthesized before gave me goose bumps! Thus, I would say I became interested in synthesis during my work in the industry in 2006–2007.

SYNFORM What do you think about the modern role and prospects of organic synthesis?

Dr. S. G. Modha Chemistry is divided into more sub-fields than ever before and thus this question is more relevant than ever. As an organic chemist I believe that the time is gone when one could just synthesize new molecules and be happy



A115

with that. One must realize that new molecules without any applications are as good as nothing. On the other hand, if one needs organic molecules, he/she has to go to organic synthesis, and thus demand for organic chemistry is never diminishing. The ever-increasing need for new and better antibiotics to fight multi-drug resistance is one of the most pressing issues suggesting the modern role of organic synthesis to be of high importance. Under the current scenario of COVID-19, I think people and governments alike have understood the importance of science. Hopefully this unfortunate experience will make them realize that funding science should be of highest priority and if that happens then prospects of organic synthesis and science in general are brighter.

SYNFORM Could you tell us more about your group's areas of research and your aims?

Dr. S. G. Modha My group's area of research is mainly development of new and improved atom-economical routes under mild reaction conditions. In particular, combination of multicomponent reactions with other atom-economical and sustainable reactions is the broad theme of my group's research. My aim is to open my students towards vast possibilities that organic chemistry brings and how they can contribute to society by developing new and improved chemical reactions. As our knowledge of chemistry and science in general is expanding with each passing day, I aim towards designing alternate reaction pathways which are atom-economical and sustainable for some organic reactions that do not fit into these categories right now.

SYNFORM What is your most important scientific achievement to date and why?

Dr. S. G. Modha Diaryliodonium reagents have been known since 1894, but there were no reports till 2015 wherein both aryls of this reagent were utilized in an organic transformation. As a post-doctoral researcher at School of Chemistry, University of Manchester (UK), I took up this challenge and successfully developed the first pathway towards atomeconomical use of diaryliodonium reagents. We reported sequential C- and N-arylation of indole via copper catalysis using just one equivalent of the iodonium reagent (*J. Am. Chem. Soc.* **2015**, *137*, 1416–1419). The most interesting part of this research was that we could utilize an unsymmetrical diaryliodonium reagent and could do selective C- and N-arylation with high selectivity.



DMSO-Catalyzed Late-Stage Chlorination of (Hetero)arenes

A116

Nat. Catal. 2020, 3, 107-115

Chlorinated (hetero)arenes are widely used in organic synthesis and in pharmaceuticals. Professor Ning Jiao - an organic synthesis expert at Peking University (P. R. of China) pointed out that Pfizer systematically screened more than 220,000 aryl derivatives and found that aromatic chlorination may change the physicochemical properties of drugs, such as pK_{a} , dipole moment, as well as metabolic rate, therefore improving pharmacokinetic and pharmacological properties (Drug Metab. Lett. 2011, 5, 232–242). "As the Nobel Laureate James Black stated, 'the most fruitful basis for the discovery of a new drug is to start with an old drug' (Lancet 2000, 355, 1022), therefore, late-stage chlorination may provide a reliable shortcut for the discovery of new drugs," said Professor Jiao. Although some elegant chlorination reactions have been developed these protocols suffer from either the high cost of the chlorinating reagents or harsh conditions. "Therefore, the efficient and practical late-stage chlorination of bioactive molecules remains formidably challenging," he added.

Previously, Professor Jiao's group had developed the HX/ DMSO (X = Br, I) system for late-stage aromatic bromination and iodination of bioactive molecules, in which DMSO acted as a mild oxidant (Org. Lett. 2015, 17, 2886–2889). "The mild reaction conditions and operational simplicity make this method very attractive. Since its publication, the HX/DMSO halogenation system has been widely applied in total synthesis of natural products, halogenation of heteroarenes, and modification of bioactive molecules," remarked Professor Jiao. He continued: "After bromination and iodination, we investigated the oxidative chlorination of arenes with the HCl/DMSO system; unfortunately, the efficiency was very low because the electrode potential of Cl⁺/Cl⁻ is higher than that of Br⁺/Br⁻." The failure of this oxidative approach motivated the group to develop other strategies that could lead to a mild and efficient late-stage chlorination of arenes.

During the study of aromatic bromination with HBr/ DMSO, Professor Jiao's group found that the reaction rate increased if there was a slight excess of DMSO relative to HBr. "Since HBr is oxidized to bromonium ion *in situ*, we speculated that a suitable amount of DMSO may promote the halogenation. We reasoned that if that was actually the case, DMSO may be used as a Lewis base to catalyze the challenging chlorination," explained Professor Jiao. With the goal of proving this hypothesis, the group tested the chlorination of arenes with commercially available NCS as the chlorine source. "At the beginning of this study, the chlorination was performed with DMSO as the solvent," Professor Jiao said. "Unfortunately, only trace amounts of chlorination product were detected. Surprisingly, the chlorination of xanthotoxin, a natural product extracted from Ammi majus plant, afforded the product in 90% yield with 20 mol% of DMSO as the catalyst. Then, we achieved the late-stage chlorination of drugs and natural products (Scheme 1, A)." Gratifyingly, the group found that many functional groups, including -OH and -NH₂, could be perfectly tolerated in this system. These experiments indicated that the reactivity of NCS was strongly improved by DMSO, which made the chlorination possible. Moreover, the newly discovered DMSO catalysis was compared to some other reported Lewis acid and Lewis base promoted catalytic approaches. "For the chlorination of single amino acid N-Bz-Tyr-OMe and approved drug diclofenac, low efficiencies were obtained with other catalysts including Ph₂P=S, FeCl₂, ZrCl₄, AuCl₂ and Ru(bpy)₃Cl₂, respectively," noted Professor Jiao. He continued: "Conversely, these experiments revealed that DMSO catalysis featured high efficiency and good functional-group tolerance. In addition, this DMSO/NCS chlorination system performed very well on a multi-gram scale. Naproxen, gemfibrozil, and indole substrates were chlorinated in high yields, which shows great potential for industrial applications (up to 54.5 g)."

Professor Jiao noted that chemical modification of peptides or proteins has emerged as an invaluable tool in biochemistry (*Chem. Rev.* **2015**, *115*, 2174–2195). "Tyrosine is essential in proteins and many peptide drugs such as alarelin and oxytocin," he explained. "Importantly, a series of structurally diverse peptides with tyrosine residue could also be chlorinated efficiently by this novel DMSO catalytic protocol (Scheme 1, B). It is noteworthy that most of the chlorination reactions did not perform well in the absence of DMSO." This chemistry provides a practical synthetic protocol and shows great potential in biological applications, as well as in drug discovery and development, which also opens an avenue for further exploration and utilization of DMSO in organic synthesis.

Professor Jiao concluded: "An efficient late-stage chlorination of complex natural products, drugs, and peptides has been developed by our group. The very common and humble DMSO was disclosed as an efficient Lewis base catalyst. The strong polarization of DMSO makes the activation of chlorocation possible, thus enabling the chlorination process. Accordingly, our mechanistic studies revealed that the active DMSO-Cl+ was the key intermediate of this protocol."







A117



About the authors



From left: S. Song, W. Wang, J. Wei, Y. Zhu, X. Shi, X. Zhang, N. Jiao

Song Song was born in Jiangsu, P. R. of China, in 1985. He received his B.Sc. degree in chemistry from Nankai University (P. R. of China) in 2007. He received his Ph.D. under the supervision of Prof. Qi-Lin Zhou at Nankai University in 2012 and then continued to work as a research assistant. His research area was asymmetric hydrogenation of unsaturated carboxylic acids. In 2013, he conducted his postdoc with Prof. Ning Jiao at Peking University (P. R. of China). From 2015, he worked at School of Pharmaceutical Sciences in Peking University, and from 2018 to the present, he has been an assistant professor. His research interests mainly focus on development of green methodologies for halogenation and thiolation.

Xinyao Li was born in Zhejiang, P. R. of China, in 1988. He studied organic chemistry under the supervision of Prof. Jiaxi Xu during his B.Sc. and M.Sc. at Beijing University of Chemical Technology (P. R. of China). He received his Ph.D. under the supervision of Prof. Ning Jiao at School of Pharmaceutical Sciences in Peking University (P. R. of China) in 2016 and then continued to work as a research assistant. In 2017, he was awarded a Humboldt Research Fellowship and conducted his postdoc with Prof. Thorsten Bach at the Technical University of Munich (Germany). His research interests mainly lie in transition-metal catalysis, radical chemistry, enantioselective photochemistry and DFT calculations for mechanistic studies.

Weijin Wang was born in Yunnan, P. R. of China, in 1998. He is currently a fourth-year undergraduate student at School of Pharmaceutical Sciences, Peking University (P. R. of China). Under the supervision of Prof. Ning Jiao, his research interests focus on developing green and efficient halogenation methodologies. **Jialiang Wei** was born in Hebei province, P. R. of China. He received his B.Sc. in macromolecular materials and engineering in 2016 from Sun Yat-sen University (P. R. of China). From 2013 to 2016, he was a member of the national "Top-Notch Undergraduate Program for Pure Science" under the supervision of Prof. Xiaodan Zhao doing research on organoselenium/sulfur catalysis. Currently, he is a Ph.D. candidate in Prof. Ning Jiao's group in Peking University (P. R. of China). His current research interests focus on the design and synthesis of novel and efficient catalysts for aerobic oxidation.

Yiqun Zhang was born in Tianjin, P. R. of China, in 1993. He studied aerobic oxidative dehydrogenative coupling under the supervision of Prof. Ning Jiao and Prof. Song Song during his B.Sc. and M.Sc. at Peking University (P. R. of China). In 2018, he moved to the Chinese University of Hong Kong (P. R. of China) and worked as a Ph.D. candidate under the supervision of Prof. Qian Miao. Now his research interest is covalent and non-covalent organic network construction based on negatively curved nanographene.

Lingsheng Ai was born in Beijing, P. R. of China, in 1994. He studied medicinal chemistry under the supervision of Prof. Ning Jiao during his B.Sc and M.Sc. at Peking University (P. R. of China). His research interests mainly lay in the green chemistry and synthetic methodology. After graduation in 2018, he works for New Oriental Group as a manager in Beijing (P. R. of China).

Yuchao Zhu was born in Shandong, P. R. of China, in 1990. He studied pharmaceutical sciences at Peking University (P. R. of China) during his B.Sc. In 2014, he moved to State Key Laboratory of Natural and Biomimetic Drugs in Beijing (P. R. of China) and worked with Prof. Ning Jiao on medicinal chemistry. He received his Ph.D. under the supervision of Prof. Ning Jiao at Peking University in 2019. His research interests mainly lie in developing efficient organic methodologies for drug synthesis.

Xiaomeng Shi received her Bachelor's degree in chemistry from Peking University (P. R. of China) in 2004. Then she went to Boston College in Boston, MA, USA and graduated with a PhD from Prof. Mary Roberts' lab in the field of mass spectrometry and protein science. After graduation, Xiaomeng worked in Northeastern University, Barnett Institute for Chemical & Biological Analysis (USA) as a postdoctoral research associate with Prof. John Engen. After two years of work experience in Tsinghua University (P. R. of China), Xiaomeng now is working as an associate research fellow in the State Key Laboratory of Natural and Biomimetic Drugs, Peking University. Her major research interests are proteomics, intact biomolecular mass spectrometry and hydrogen-deuterium exchange mass spectrometry.

A119

Xiaohui Zhang was born in Henan, P. R. of China, in 1990. He studied analytical chemistry under the supervision of Prof. Xinxiang Zhang during his BSc and PhD at Peking University (P. R. of China), mainly researching on capillary electrophoresis and tumor metabolism. Since 2018, he has been technologist-incharge at State Key Laboratory of Natural and Biomimetic Drugs at Peking University. He is responsible for the daily operation of the mass spectrometry platform and protein sample analysis.

Ning Jiao was born in Shandong, P. R. of China, in 1976. He received his B.Sc. degree at Shandong University (P. R. of Chi-

na, 1999), and Ph.D. (2004) (with Prof. Shengming Ma) at Shanghai Institute of Organic Chemistry (SIOC), Chinese Academy of Science (CAS, P. R. of China). He spent 2004–2006 as an Alexander von Humboldt postdoctoral fellow with Prof. Manfred T. Reetz at Max Planck Institute für Kohlenforschung (Germany). In 2007, he joined the faculty at Peking University (P. R. of China) as an Associate Professor and was promoted to Full Professor in 2010. His current research efforts are focused on: 1) new methodologies development in aerobic oxidation, oxygenation, nitrogenation, and halogenation reactions; 2) first-row-transition-metal catalysis and inert chemical bond activation; 3) drug discovery and development.

ethylation of a-Diazo

Enantioselective Three-Component Aminomethylation of α -Diazo Ketones with Alcohols and 1,3,5-Triazines

A120

Nat. Commun. 2020, DOI: 10.1038/s41467-020-15345-2

β-Amino carbonyl moieties are versatile synthetic building blocks that can be employed toward a wide variety of natural products and biologically active compounds. As an important research branch of the Mannich reaction. the aminomethylation reaction of ketones or aldehydes represents an efficient protocol for accessing β-amino carbonyl compounds. "Recent years have witnessed the application of a variety of aminomethylation reagents to the enantioselective variant of the reaction; however, asymmetric induction of these transformations is dominated by the use of chiral catalysts to activate the nucleophiles, i.e. chiral amines or Lewis acids with chiral ligands," said Professor Dong Xing from East China Normal University (P. R. of China), adding: "While this nucleophilebased activating strategy was feasible and showed good enantiocontrol, the nucleophiles were limited to inherently activated substrates such as unmodified ketones or 1,3-dicarbonyl compounds, thus the scope and application of this type of transformation were also significantly limited (Scheme 1, a)."

The groups of Professor Xing and Professor Wenhao Hu (Sun Yat-sen University, P. R. of China) have a long-standing common research interest in carbene-involved enantioselective multicomponent reactions (MCRs) via cooperative catalysis. "As part of our ongoing research, we designed the rhodium(II)/ chiral phosphoric acid (CPA) co-catalysed three-component reaction of α -diazo ketone, alcohol and 1,3,5-triaryl-1,3,5-triazine, with the hope that an electrophile activation strategy would be established (Scheme 1, b)," said Professor Hu, continuing: "However, due to the instability and low concentration properties of the formaldimine species generated in situ from 1,3,5-triazine, it remains uncertain whether such an electrophile activation mode is workable."

At the very beginning of their exploration, tremendous efforts were made to modify the reaction conditions with different α -carbonyl diazo compounds as the carbene precursor. "It finally turned out that the structure of the diazo compound has a significant impact on the outcome of the reaction," explained Professor Xing. He added: "We continually noticed that the structure of both the CPA and the rhodium(II) catalyst are both responsible for the stereoselective control of this transformation."

With the established optimized reaction conditions, a wide range of alcohols, including simple aliphatic alcohols,

allylic alcohol, propargyl alcohol, complex natural alcohols, and even water could all be applied to this three-component aminomethylation. "It is also impressive that this method can be used for the efficient linkage of two complicated drug candidates (Scheme 1, c)," said Professor Xing.

Professor Hu concluded: "This work offers an efficient electrophile-based asymmetric activation for aminomethylation with unstable formaldimine species. Further efforts to apply this protocol to the trapping of other types of active intermediates are ongoing in our labs."





A121



About the authors



Prof. W. Hu

Wenhao Hu received a B.S. in chemistry from Sichuan University (P. R. of China, 1987), M.S. from Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences (P. R. of China, 1990), and Ph.D. from Hong Kong Polytechnic University (P. R. of China) with Professor Albert S. C. Chan (1998). He became a postdoctoral fellow at the University of Arizona (USA) with Professor Michael P. Doyle. In 2002-2006, he worked as a

Dong Xing received his B.S. and M.S. degrees from East China Normal Uni-

versity (P. R. of China, 2003) and his Ph.D. from the University of Hong

Kong (P. R. of China) with Professor

Dan Yang in 2011. He then joined

the School of Chemistry and Molecu-

lar Engineering at East China Normal

University as an assistant professor

and Chenhui Scholar. He was then

promoted to associate professor in

medicinal chemist at Genesoft in South San Francisco (USA) and a process chemist at Bristol-Myers Squibb in New Jersey (USA). In 2006 he joined East China Normal University (P. R. of China) as a professor of Chemistry, and the head of Shanghai Engineering Research Center of Molecular Therapeutics and New Drug Development. In 2016, he moved to Sun Yat-sen University (P. R. of China) as the head of School of Pharmaceutical Science. He received National Natural Science Funds for Distinguished Young Scholar (2011). His research interests include synthetic methodology, MCRs, and medicinal and process chemistry.



Prof. D. Xing

2015. In 2016, he began his postdoctoral research with Professor Guangbin Dong at the University of Texas at Austin (USA) and then moved to the University of Chicago (USA) with the Dong laboratory. In 2018, he returned to East China Normal University, where his current research is focused on atom-economic reactions with novel cooperative catalytic system designs.



J. Che



L. Niu



Dr. S. Jia

Jiuwei Che received his B.S. degree in chemical engineering from Northeast Forestry University (P. R. of China) in 2012. He was granted his M.S. (2015) and Ph.D. (2019) degrees in chemistry from East China Normal University (P. R. of China) under the supervision of Prof. Wenhao Hu and Dr. Dong Xing. His research topic focuses on asymmetric multicomponent aminomethylation.

Li Niu received her B.S. degree in chemistry from East China University of Science and Technology (P. R. of China) in 2016. She then obtained her M.S. degree in medicinal chemistry from East China Normal University (P. R. of China) in 2019. Her research topic focuses on efficient synthesis and bioactivity evaluation of bioxindole compounds.

Shikun Jia received his B.S degree in chemistry from Zhengzhou University (P. R. of China) in 2012. He obtained his Ph.D. in chemistry from East China) Normal University (P. R. of China) under the supervision of Prof. Wenhao Hu in 2017. Then he moved to Sun Yat-sen University (P. R. of China) for postdoctoral research (2017–2019). In 2019, he began his independent career at Zhengzhou University (P. R. of China).

_

Coming soon

🗕 Editorial Board Focus

Prof. Xiaozhang Zhu (Institute of Chemistry, Chinese Academy of Sciences, P. R. of China)

Literature Coverage

Efficient Co-Catalyzed Double Hydroboration of Nitriles: Application to One-Pot Conversion of Nitriles into Aldimines

Literature Coverage

A General ¹¹C-Labeling Approach Enabled by Fluoride-Mediated Desilylation of Organosilanes

Further highlights

SynthesisReview: Recent Advances in DecarboxylativeReactions of Alkynoic Acids(by M. A. Idris and S. Lee)

Synlett Account: Selected Research Topics of the Dondoni Group over the Last Two Decades (2000–2020) (by A. Dondoni)

Synfacts Synfact of the Month in category "Flow Chemistry": Synthesis of Deuterodifluoromethylated Alcohols and *gem*-Difluoro Alkenes in Continuous Flow

Editor Matteo Zanda, Chair in Biomolecular Imaging, Centre for Imaging Science, Department of Chemistry, School of Science, Loughborough University, Leicestershire, LE11 3TU, UK and

C.N.R. – Istituto di Chimica del Riconoscimento Molecolare Via Mancinelli, 7, 20131 Milano, Italy Editorial Assistant: Alison M. Sage

synform@outlook.com; fax: +39 02 23993080

Editorial Office

- Managing Editor: Susanne Haak, susanne.haak@thieme.de, phone: +49 711 8931 786 Scientific Editors: Stefanie Baumann, stefanie.baumann@thieme.de, phone: +49 711 8931 776 Selena Boothroyd, selena.boothroyd@thieme.de
- Michael Binanzer, michael.binanzer@thieme.de, phone: +49 711 8931 768 Giuliana Rubulotta, giuliana.rubulotta@thieme.de, phone: +49 711 8931 183 Kathrin Ulbrich, kathrin.ulbrich@thieme.de, phone: +49 711 8931 785 Senior Production Editors:
- Thorsten Schön, thorsten.schoen@thieme.de, phone: +49 711 8931 778 Thorsten Schön, thorsten.schoen@thieme.de, phone: +49 711 8931 781 Draduction Managar Scabia Magart
- Production Manager: Sophia Hengst,
 sophia.hengst@thieme.de, phone: +49 711 8931 398
 Production Assistant: Tobias Brenner,
- Tobias.brenner@thieme.de, phone: +49 711 8931 769
- Editorial Assistant: Sabine Heller, sabine,heller@thieme.de, phone: +49 711 8931 744
- sabine.heller@thieme.de, phone: + Marketing Director: Julia Stötzner,
- julia.stoetzner@thieme.de, phone: +49 711 8931 771
- Postal Address: Chemistry Journals, Editorial Office, Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany,
 - Homepage: www.thieme-chemistry.com

Publication Information

Synform will be published 12 times in 2019 by Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany, and is an additional online service for Synthesis, Synlett and Synfacts.

Publication Policy

Product names which are in fact registered trademarks may not have been specifically designated as such in every case. Thus, in those cases where a product has been referred to by its registered trademark it cannot be concluded that the name used is public domain. The same applies as regards patents or registered designs.

Ordering Information for Print Subscriptions to Synthesis, Synlett and Synfacts

The Americas: Thieme Publishers New York, Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.

Via e-mail: customerservice@thieme.com Via website: www.thieme-chemistry.com Phone: +1 212 760 0888; Fax: +1 212 947 1112 Order toll-free within the USA: +1 800 782 3488

Europe, Africa, Asia, and Australia: Thieme Publishers Stuttgart, Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany. Via e-mail: customerservice@thieme.de

Via website: www.thieme-chemistry.com Phone: +49 711 8931 421; Fax: +49 711 8931 410

Current list prices are available through www.thieme-chemistry.com.

Online Access

The online versions of Synform as well Synthesis, Synlett, Synfacts and SynOpen are available through www.thiemeconnect.com/products/ejournals/journals) where it is also possible to register for a free trial account. For information on multi-site licenses and pricing for corporate customers as well as backfiles, please contact our regional offices:

The Americas: esales@thieme.com, phone: +1 212 584 4695 Europe, Africa, Asia, and Australia: eproducts@thieme.de, phone: +49 711 8931 407 India: eproducts@thieme.in, phone +91 120 45 56 600 Japan: brhosoya@poplar.ocn.ne.jp, phone +81 3 3358 0692

Manuscript Submission to Synthesis, Synlett, and SynOpen

Manuscript submissions will be processed exclusively online via http://mc.manuscriptcentral.com/synthesis, http://mc.manuscriptcentral.com/synlett and http://mc.manuscriptcentral.com/synopen, respectively. Please consult the Instructions for Authors before compiling a new manuscript. The current version and the Word template for manuscript preparation are available for download at www.thieme-chemistry.com.

Copyright

This publication, including all individual contributions and illustrations published therein, is legally protected by copyright for the duration of the copyright period. Any use, exploitation or commercialization outside the narrow limits set by copyright legislation, without the publisher's consent, is illegal and liable to criminal prosecution. This applies to translating, copying and reproduction in printed or electronic media forms (databases, online network systems, Internet, broadcasting, telecasting, CD-ROM, hard disk storage, microcopy edition, photomechanical and other reproduction methods) as well as making the material accessible to users of such media (e.g., as online or offline backfiles).

Copyright Permission for Users in the USA

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Georg Thieme Verlag KG Stuttgart - New York for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of USS 25.00 per copy of each article is paid directly to CCC, 22 Rosewood Drive, Danvers, MA 01923, USA, 0341-0501/02.

For current SYNFORM articles, please visit www.thieme-chemistry.com SYNFORM issue 2020/09 is available from August 18, 2020 at www.thieme-connect.com/ejournals