


Validation of the Six-item Female Sexual Function Index in Middle-Aged Brazilian Women

Validação do índice de funcionamento sexual feminino-6 em mulheres brasileiras de meia-idade

Mona Lúcia Dall'Agno¹  Charles Francisco Ferreira¹ Fernanda Vargas Ferreira¹
Faustino R. Pérez-López² Maria Celeste Osório Wender¹

¹Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil

²Department of Obstetrics and Gynecology, Lozano-Blesa University Hospital, Zaragoza, Spain

Address for correspondence: Mona Lúcia Dall'Agno, MSc, Rua Ramiro Barcelos, 2350, 90035-903, Largo Eduardo Faraco, Serviço de Ginecologia e Obstetrícia, Porto Alegre, RS, Brasil (e-mail: modallagno@hotmail.com).

Rev Bras Ginecol Obstet 2019;41:432–439.

Abstract

Objective To validate the six-item female sexual function index (FSFI-6) in middle-aged Brazilian women.


Methods Cross-sectional observational study, involving 737 (premenopausal $n = 117$, perimenopausal $n = 249$, postmenopausal $n = 371$) Brazilian sexually active women, aged between 40 and 55 years, not using hormonal contraceptive methods. The Brazilian FSFI-6 was developed from the translation and cultural adaptation of the Portuguese FSFI-6 version. The participants completed a general questionnaire, the FSFI-6, and the menopause rating scale (MRS). The validation was performed by AMOS 16.0 software (SPSS, Inc., Chicago, IL, USA) for a confirmatory factor analysis (CFA). The chi-square of degrees of freedom (χ^2/df), the comparative fit index (CFI), the Tucker-Lewis index (TLI) and the root-mean-square error of approximation (RMSEA) were used as indices of goodness of fit. Cronbach α coefficient was used for internal consistency.

Results The process of cultural adaptation has not altered the Brazilian FSFI-6, as compared with the original content. The CFA for the FSFI-6 score showed an acceptable fit ($\chi^2/df = 3.434$, CFI = 0.990, TLI = 0.980, RMSEA = 0.058, 90% confidence interval (90%CI) = 0.033–0.083, $p \leq 0.001$) and a good reliability was established in FSFI-6 and MRS (Cronbach $\alpha = 0.840$ and = 0.854, respectively). In addition, 53.5% of the sample had low sexual function.

Conclusion The FSFI-6 was translated and adapted to the Brazilian culture and is a consistent and reliable tool for female sexual dysfunction screening in Brazilian middle-aged women.

Keywords

- climacteric
- cross cultural comparison
- menopause
- test reliability
- female sexual health

 Mona Lúcia Dall'Agno's ORCID is <https://orcid.org/0000-0002-5424-8090>.

received
December 4, 2018
accepted
May 21, 2019

DOI <https://doi.org/10.1055/s-0039-1692694>.
ISSN 0100-7203.

Copyright © 2019 by Thieme Revinter
Publicações Ltda, Rio de Janeiro, Brazil

License terms



Resumo

Palavras-chave

- climatério
- comparação transcultural
- menopausa
- reprodutibilidade dos testes
- saúde sexual da mulher

Objetivo Validar o Índice de Função Sexual Feminina - 6 itens (FSFI-6, na sigla em inglês) para mulheres brasileiras de meia-idade.

Métodos Estudo transversal observacional que incluiu 737 (pré-menopausa $n = 117$, perimenopausa $n = 249$, pós-menopausa $n = 371$) mulheres brasileiras sexualmente ativas, entre 40 e 55 anos, sem métodos contraceptivos hormonais. A versão brasileira do FSFI-6 foi desenvolvida através da tradução e adaptação cultural da versão portuguesa do questionário. As participantes preencheram um questionário com dados gerais, o FSFI-6 e a escala de avaliação da menopausa (*menopause rating scale* [MRS]). A validação do instrumento se deu através de análise fatorial confirmatória (CFA, na sigla em inglês), realizada pelo software AMOS 16.0 (SPSS, Inc., Chicago, IL, EUA). Qui-quadrado sobre graus de liberdade (χ^2/df), índice de ajuste comparativo (CFI, na sigla em inglês), índice de Tucker-Lewis (TLI) e raiz média dos quadrados dos erros de aproximação (RMSEA, na sigla em inglês) foram utilizados como índices de adequação de ajustes. O coeficiente alfa de Cronbach foi utilizado para avaliar a consistência interna.

Resultados O processo de adaptação cultural não alterou a versão brasileira do FSFI-6, comparado ao conteúdo original. O CFA para o escore do FSFI-6 demonstrou ajuste aceitável ($\chi^2/df = 3,434$; CFI = 0,990; TLI = 0,980; RMSEA = 0,058; 90% IC = 0,033 a 0,083; $p \leq 0,001$). Demonstrou-se boa confiabilidade entre FSFI-6 e MRS (alfa de Cronbach = 0,840 e = 0,854, respectivamente). Do total, 53,5% da amostra apresentou baixa função sexual.

Conclusão O FSFI-6 foi traduzido e adaptado culturalmente, e é uma ferramenta consistente e confiável no rastreamento de disfunções sexuais em mulheres brasileiras de meia-idade.

Introduction

Sexual function (SF) is an important component of the quality of life (QoL) of peri- and postmenopausal women.¹ The climacteric population is progressively growing all over the world and is characterized by a range of signs and symptoms, with low sexual desire being a well reported one.²⁻⁴ Due to estrogen deficiency, menopause is related to other consequences in SF, as vaginal atrophy, reduction in lubrication and dyspareunia.⁵

The climacteric period impacts sexuality through the interaction between hormonal, biological, social, cultural and other individual characteristics, and can negatively impact the QoL.^{6,7} Although female sexual dysfunction (FSD) seems to increase with age,⁵ its actual prevalence is variable according to the literature.⁸ Cultural factors, physician-patient relationship, and lack of standardization of diagnosis are possibly determinants of this variation.⁸

Due to its complexity,⁹ questionnaires to assess female sexual health have been developed, providing a better understanding of subjective and objective aspects, allowing comparisons between individuals and populations.¹⁰ In addition, they are inexpensive, non-invasive, useful for health professionals and generally, self-fulfilling.

One of the most worldwide used instruments for assessment of FSD is the female sexual function index (FSFI),^{11,12} with reliability/validity demonstrated extensively in several studies for different populations.⁸

The original FSFI has 19 items and includes the 6 domains of FSD,¹¹ according to the International Classification of Diseases-10 (ICD-10) and the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV): desire, arousal, lubrication, orgasm, satisfaction, and pain.¹¹ The FSFI was validated for many languages and populations, including some Brazilian versions that have already been published.¹²⁻¹⁶

In an attempt to obtain a smaller and faster instrument, a short version of the FSFI was proposed, consisting of six items.¹⁷ The selected items refer to the six domains of the FSD (one item for each domain), maintaining the psychometric properties and reliability of the original tool.¹⁷ This shorter version was validated in Spanish,¹⁸⁻²¹ Korean,²² and Portuguese for a sample of Portuguese women.²³

Despite the importance of the SF for the QoL, it is rarely investigated in the women's health care.^{6,11,24} The minority of women with a sexual problem seek for professional help.²⁴ The use of an easier and faster questionnaire may be a relevant tool in the medical assistance of these women, thus allowing adequate assessment of SF.

Therefore, this research aimed to validate the FSFI-6, assessing middle-aged women's sexual function and associated variables in the Southern region of Brazil. It is suggested that the validation results in a reliable instrument applicable in research and clinical practice, as well as in the publication of important data on the female sexual health.

Methods

Study Design and Participants

This is a cross-sectional study, performed in areas of free access and transit of the population (e.g., parks, squares, streets, shopping) in the Southern region of Brazil, from January to October 2017, involving middle-aged women. The cities where the questionnaires were applied are located in the three states of the Southern region of Brazil, with populations varying between 47,000 and 1,484,000 inhabitants. Ethical approval was obtained from the Institutional Review Board of the Hospital de Clínicas de Porto Alegre (HCPA, Ref. N°. 16-0621). The study involved women in the community, aged between 40 and 55 years, classified as pre-, peri-, or postmenopausal, according to the Stages of Reproductive Aging Workshop +10 (STRAW +10),³ not using hormonal contraceptive methods (e.g., contraceptive use of levonorgestrel intrauterine device, combined hormonal contraceptives, progestin only pills or implants), who reported sexual activity in the past 4 weeks and that agreed to participate. Women unable to understand the survey or having incapacity imposing difficulties during the filling the questionnaire were excluded. After being informed of the study (e.g., objectives and tools used) and providing written consent, the surveyed women were requested to voluntarily fill out a general questionnaire containing health, habits and sociodemographic data, the FSFI-6 and the menopause rating scale (MRS), with the help of a drawing board. Sample size calculation was performed using the software WinPEPI, PEPI-for-Windows, version 11.65 based on the fact that 30 to 50% of middle-aged women would present lower sexual function (Blümel et al,⁵; Pérez-López et al¹⁹; Llaneza et al²⁵). Hence, a minimal sample of 715 participants was calculated, considering a 5% of losses, 5% desired precision, and a 99% confidence limit.

General Questionnaire

A questionnaire was developed by the researchers, and it contains female data, including age (years), partner status (yes/no), marital status, educational level (total years), sexual status in the past 4 weeks (active or inactive), parity, professional status, number of people living in the household, and family income (in minimum wages). Health data related to menopausal status (pre-, peri-, and postmenopausal), according to the STRAW + 10³ were collected, such as menopause age, type of menopause (natural or surgical), pharmacological treatment for menopausal symptoms (yes/no, type). General health and disease data, surgical procedures performed, psychological problems (depression, anxiety), urinary loss, anthropometric data (weight, height, and body mass index [BMI]), as well as life habits (smoke, alcohol and coffee consumption, physical activity) were assessed.

The FSFI-6

This instrument is composed of 6 questions based on existing items of the FSFI, each covering one of the original domains: desire, arousal, lubrication, orgasm, satisfaction, and pain.¹⁷

Each item can provide a score varying from 0 to 5, whose sum provides a total FSFI-6 score.¹⁷ This is a screening tool aiming to identify women at high risk of FSD.¹⁷

The MRS

The MRS scale is a valuable tool for assessing climacteric symptoms and health-related QoL of climacteric women through 11 items distributed in 3 domains: somatic, psychological, and urogenital symptoms.²⁶ The score for each question ranges from 0 (absence of symptoms) to 4 (very severe symptoms). The total score is obtained by adding the score of each domain. The higher the score reached, the worse the QoL related to the climacteric symptoms.^{27,28} In addition, symptom severity can be categorized for each domain: absent or occasional (0–4 points), mild (5–8 points), moderate (9–15 points) or severe (≥ 16 points) symptoms.²⁶ The Brazilian version of the MRS was already validated.²⁹

Translation and validation

Considering that Brazilian and Portuguese women do not share the exact same cultural and ethnical background, and therefore differences could exist in the understanding and the interpretation of the instrument questions, two stages were used for conducting this study. After obtaining the authorization of the author who validated the FSFI-6 to Portuguese language,²³ the first stage was performed and included translation and cultural adaptation of the original scale to our context. The instructions of the World Health Organization (WHO) were followed (e.g., translated by two independent and experienced native speakers of the target language translators; reconciliation by the researchers and the translators, who resolves the discrepancies of the forward translations into a single version; back translation by a different bilingual translator; and harmonization between the translations).³⁰ The cognitive debriefing was performed with 30 subjects from the study population (community women between 40 and 55 years, 10 each menopausal status) to assess the degree of understanding and testing for cognitive equivalence. At the end of the tool, the question “*Is there a word that has not been understood?*” detects words not understood by women in the early stages of cultural adaptation and validation. These subjects were not included in the analyzes of this study. Review of cognitive debriefing results for finalization and proofreading by the researchers and an expert committee (e.g., three gynecologists, one psychologist, and one biologist) was also performed.³⁰ Cultural equivalence was established according to the criteria by Guillemin et al:³¹ at least 85% of the subjects should not show any kind of difficulty to answer each question (e.g., no question should be considered incomprehensible by over 15% of the participants). The second stage was the instrument validation, corresponding to the statistical analysis of its psychometric properties. The validity occurred by the comparison with a specific reference tool (MRS), which has been previously validated for the Brazilian Portuguese.²⁹ The final Brazilian FSFI-6 questionnaire is shown in ► **Supplemental Material 1**.

Statistical Analysis

Regarding the data processing, database double entry, review and analysis were performed using the SPSS, version 18.0. (SPSS Inc., Chicago, IL, USA). Symmetric data was expressed as mean and standard error of mean (SEM), or by median and 25th to 75th percentiles (P25–P75). The Shapiro-Wilk test was used to determine the normality of data distribution. Categorical variables were described as absolute (n) and relative (n%) frequencies. According to this, differences in FSFI-6 domains and total scores were analyzed with the Kruskal-Wallis test (bivariate analysis, Dunn *post-hoc*) or the chi-squared test, with adjusted residual analysis for independent samples. To examine the construct validity of our model for mathematical FSFI-6, the factor loadings of the variables in each model was calculated with AMOS 16.0 software (SPSS, Inc., Chicago, IL, USA). A confirmatory factor analysis (CFA) was conducted. The chi-squared of degrees of freedom (χ^2/df), the comparative fit index (CFI), the Tucker-Lewis index (TLI), and the root-mean-square error of approximation (RMSEA) were used as indices of goodness of fit. The internal consistency (criterion validity) of FSFI-6 and MRS instruments was assessed using Cronbach α coefficients.³² Spearman ρ coefficients were estimated for determining the correlations between FSFI-6 total scores and variables, including scores obtained with the MRS.

The level of significance was set at 5% for all analyses.

Results

In total, 737 women were invited and met inclusion criteria of this study, being classified as pre- ($n = 117$), peri- ($n = 249$), or postmenopausal ($n = 371$) according to the STRAW + 10 criteria. They were included and answered a general questionnaire containing health, habits, and socio-demographic data; the Brazilian version of the FSFI-6; and the MRS. The general characteristics of the surveyed women are presented in ►Table 1. Briefly, most women were aged between 50 and 54 years (39.8%), multiparous (63.9%), married or living with partner (76.1%), currently having a sexual partner (96.3%), and with natural menopause (78.7%). The median (P25–P75) time since menopause onset was 5.00 (3.00–7.50) months, and most of them did not use medication for menopause symptoms (85.6%). Considering health and habits aspects, 116 (15.7%) were smokers, 130 (17.6%) had hypertensive disorder, 40 (5.4%) were diabetic, 193 (26.2%) had a psychiatric condition (e.g., depression symptoms and anxiety) and 116 (15.7%) were sedentary.

To achieve the cognitive debriefing, 30 climacteric women ($n = 10$ in each menopausal status) were included, and no difficulty was observed in the understanding of the proposed FSFI-6 version (data not shown). The Brazilian version of the questionnaire also underwent an assessment by an expert committee (e.g., three gynecologists, one psychologist, and one biologist). There were no suggestions for changes in this version of the FSFI-6. For the question added at the end of the questionnaire (e.g., “Is there a word that has not been understood?”), the answer “yes” was not marked by any of the

Table 1 Characteristics of all surveyed women

Female data	N = 737
Age (years)	
40–44	117 (15.9)
45–49	210 (28.5)
50–54	293 (39.8)
55–59	117 (15.9)
Parity	
0	78 (10.6)
1	188 (25.5)
≥ 2	471 (63.9)
Educational level (years)	
0–6	109 (14.8)
7–12	316 (42.9)
≥ 13	312 (42.3)
Marital status	
Married or living with partner	561 (76.1)
Divorced	98 (13.3)
Single	65 (8.8)
Widowed	13 (1.8)
Currently has partner	
Yes	710 (96.3)
No	27 (3.7)
Menopausal status	
Premenopausal	117 (15.9)
Perimenopausal	249 (33.8)
Postmenopausal	371 (50.3)
Natural menopause	292 (78.7)
Surgical menopause	79 (21.3)
Time since menopause onset (months)	5.00 [3.00–7.50]
[minimum–maximum]	[0.00–31.00]
Pharmacological treatment for menopause symptoms	
No	631 (85.6)
Systemic hormone therapy	44 (6.0)
Alternative therapies (herbal teas, phytoestrogens)	58 (7.9)
Psychotropics	1 (0.1)
Topic estrogen	3 (0.4)
Habits, lifestyle, health aspects and other issues	
Current smoking	116 (15.7)
Body mass index	26.29 [23.51–30.39]
[minimum–maximum]	[16.00–54.11]
Sedentary lifestyle	116 (15.7)
Hypertension	130 (17.6)
Diabetes	40 (5.4)
Psychiatric conditions	193 (26.2)

Abbreviations: (P25–P75): 25th–75th percentiles; SEM, standard error of the mean, n, absolute frequency; n%, relative frequency. Data presented as medians (P25–P75), means (\pm SEM) or frequencies [n(n%)].




Fig. 1 Statistic model—Factor loadings of the variables in each model (AMOS 16.0 software). Abbreviations: FSFI, female sexual function index; e, latent variable. Double arrow in latent variables: correlation between items. FSFI-1 and FSFI-4 correlation: -0.322. FSFI-3 and FSFI-6 correlation: 0.208.

participants in the validation stage (data not shown). The statistic model used is shown in ►Fig. 1. Each question of the FSFI-6 was considered a factor loading. The CFA showed an acceptable fit ($\chi^2/df = 3.434$, CFI = 0.990, TLI = 0.980, RMSEA = 0.058, CI 90% = 0.033–0.083, $p \leq 0.001$). Good values were evidenced in terms of both factorial weights, as well as regarding the multiple squared correlations (data not shown).

The FSFI-6 and MRS total scores are presented in ►Table 2, considering menopausal status (pre-, peri-, and postmenopausal) groups. A FSFI-6 total score ≤ 21 is consistent with a positive screening for FSD. The cut-off considered for the Brazilian version is the calculated median of the total scores for the sample (21 points). The frequency of positive screening for FSD, was greater in the postmenopausal group (61.2%)

in relation to pre- (43.6%) and perimenopausal (46.6%) women (Chi-Square test, $p \leq 0.0001$). Peri- and postmenopausal women presented higher menopausal symptoms when compared with the premenopausal group ($p = 0.001$). A good reliability was established in FSFI-6 and MRS (Cronbach α , $\alpha = 0.840$ and $= 0.854$, respectively).

The Spearman ρ coefficients between total FSFI-6 scores and variables are displayed in ►Table 3. Total FSFI-6 scores correlated positively with family income, parity, and educational level, and inversely with age, peri- and postmenopausal status, time of menopause onset, and total MRS score.

Discussion

In this study, a Brazilian version of a shorter and faster instrument for FSD screening was translated, culturally adapted and validated. Sexual function and related variables in middle-aged Brazilian women were also assessed.

A CFA and the assessment of the internal consistency were conducted. We found the validity of the FSFI-6 to be satisfactory in a group of 737 women, when compared with previous validation studies around the world.^{17,19–22,33} The FSFI-6 CFA showed an acceptable fit ($\chi^2/df = 3.434$, CFI = 0.990, TLI = 0.980, RMSEA = 0.058, CI 90% = 0.033–0.083, $p \leq 0.001$), and good values were evidenced in terms of both factorial weights, as well as regarding the multiple squared correlations.

Also, this study indicates that the Brazilian FSFI-6 has an excellent reliability (Cronbach $\alpha = 0.840$) as a screening instrument for FSD in middle-aged women, and the climacteric symptoms and domains measured by MRS reinforced these FSFI-6 properties.

Table 2 The six-item female sexual function index: Total and each domain scores

Items	Total N = 737	Premenopausal N = 117	Perimenopausal N = 249	Postmenopausal N = 371	*p-value	Cronbach alpha
Desire [minimum–maximum]	3.00 [2.00–3.00] [1.00–5.00]	3.00 [2.00–3.00] ^a [1.00–5.00]	3.00 [2.00–3.00] ^{ab} [1.00–5.00]	3.00 [2.00–3.00] ^b [1.00–5.00]	0.015	0.840
Arousal [minimum–maximum]	3.00 [3.00–4.00] [1.00–5.00]	3.00 [3.00–4.00] ^{ab} [1.00–5.00]	3.00 [3.00–4.00] ^a [1.00–5.00]	3.00 [2.00–3.00] ^b [1.00–5.00]	≤ 0.0001	
Lubrication [minimum–maximum]	4.00 [3.00–4.00] [1.00–5.00]	4.00 [3.00–5.00] ^a [1.00–5.00]	4.00 [3.00–4.00] ^a [1.00–5.00]	3.00 [2.00–4.00] ^b [1.00–5.00]	0.002	
Orgasm [minimum–maximum]	4.00 [3.00–4.00] [1.00–5.00]	4.00 [3.00–4.00] ^{ab} [1.00–5.00]	4.00 [3.00–5.00] ^a [1.00–5.00]	3.00 [3.00–4.00] ^b [1.00–5.00]	0.010	
Satisfaction [minimum–maximum]	4.00 [3.00–4.00] [1.00–5.00]	4.00 [3.00–4.00] [1.00–5.00]	4.00 [3.00–4.00] [1.00–5.00]	4.00 [3.00–4.00] [1.00–5.00]	0.871	
Pain [minimum–maximum]	4.00 [3.00–5.00] [0.00–5.00]	5.00 [4.00–5.00] ^a [0.00–5.00]	4.00 [3.00–5.00] ^{ab} [0.00–5.00]	4.00 [3.00–5.00] ^b [0.00–5.00]	0.005	
Total [minimum–maximum]	21.00 [17.00–24.00] [6.00–30.00]	22.00 [20.11–22.02] ^a [7.00–30.00]	22.00 [20.29–21.47] ^a [7.00–30.00]	20.00 [19.17–20.18] ^b [6.00–30.00]	≤ 0.0001	
FSFI total scores ≤ 21	394(53.5)	51(43.6)	116(46.6)	227(61.2)	≤ 0.0001	0.854
MRS total score [minimum–maximum]	13.00 [7.00–21.00] [0.00–41.00]	12.00 [5.00–17.00] ^a [0.00–38.00]	13.00 [7.00–22.00] ^b [0.00–41.00]	14.00 [8.00–21.00] ^b [0.00–40.00]	0.001	

Abbreviations: (P25–P75): 25th–75th percentiles; SEM, standard error of the mean; n, absolute frequency; n%, relative frequency; FSFI-6, 6-item female sexual function index; MRS, menopause rating scale; p, statistical significance.

Data presented as medians (P25–P75) or frequencies (n[n%]).

*Kruskal-Wallis (Dunn *posthoc*) and Chi-Square tests with adjusted residual analysis. Bold numbers: association by Chi-Square test with adjusted residual analysis.

^{ab}Different letters indicate statistical significance. Significance set at 5% for all analysis.

Table 3 Correlations between the six-item female sexual function index and its variables

Items	FSFI-6 total score (N = 737)	
	Coefficient	*p-value
Not having a sexual partner	-0.033	0.375
Single or not living with a partner	0.063	0.087
Female educational level	0.206	≤ 0.0001
Female age	-0.110	0.003
Family income	0.209	≤ 0.0001
BMI	0.038	0.300
Parity	0.097	0.009
Menopause (peri- and postmenopausal status)	-0.154	≤ 0.0001
Time of menopause	-0.107	0.039
Sedentary lifestyle	0.026	0.489
MRS total score	-0.375	≤ 0.0001

Abbreviations: BMI, body mass index (Kg/m²); MRS, menopause rating scale; FSFI-6, 6-item female sexual function index, p, statistical significance.

*Spearman correlations. Significance set at 5% for all analysis.

It is important to mention that the present study is the first to provide Brazilian data on the FSFI-6 and MRS. Only women who were sexually active in the last 4 weeks were included. High rates of positive screening for sexual dysfunction were found (53.5%), mostly in postmenopausal women (61.2%). Total FSFI-6 scores were positively correlated with family income, parity, and educational level, and inversely with age, menopause transition (peri- and postmenopausal status), time of menopause onset, and total MRS score.

The menopausal transition defines an impact on the QoL due to biological, social, cultural, physical and psychological aspects, which also affect sexual life.⁴ The FSFI is a well-known tool used to assess the female sexual function, which can be used among pre- and postmenopausal women, in different ethnical populations and medical conditions, with good reliability values.⁸ The FSFI-6 was developed as a shorter and faster alternative, with the same psychometric properties.¹⁷

Isidori et al¹⁷ pointed a cut-off value of 19 points for the FSFI-6, which demonstrates a high sensitivity, specificity and positive and negative predictive values for the identification of women with positive screening for FSD in a sample of Italian women. For Lee et al,²² the calculated cut-off point for Korean women was 21. Our population differs from the one to which the FSFI-6 was originally designed for in terms of age range, mixed ethnical background as well as different degrees of education and income. In contrast to European¹⁷ and South Korean²² homogeneity, our data was compared with Latin American populations. A large Ecuadorian study used as cut-off point the median calculated

from the FSFI-6 scores.²¹ In our study, the same method was used, resulting in a score of ≤ 21 displaying FSD. The computed Cronbach α for the FSFI-6 in this Brazilian research was high, indicating an adequate internal consistency, which was similar to that of the Ecuadorian study.²¹

In this study, 53.5% of the surveyed women displayed scores ≤ 21, suggesting higher risk for sexual dysfunction. This result is comparable to those of the Ecuadorian study, which used the same method for the cut-off value calculation.²¹ Most women with a positive screening for FSD were in the postmenopausal stage (61.2%), similarly to what was observed in previous studies.^{21,33} The worsening sexual function could be explained by postmenopausal status and its consequences, and by aging, once our participants were older than those in the Latin-American studies,^{21,33} and the prevalence of FSD increases with advancing age.³⁴ Additionally, Brazilian women had lower to moderate education, multiparity, non-hormone therapy use, and were overweight, a profile characteristic of developing countries.

Although postmenopausal women showed the highest prevalence of positive screening for FSD, perimenopausal women presented 43.6%, suggesting that oscillating levels of estradiol and aging may affect sexual satisfaction.³⁵ Besides, brain centers associated with sexual arousal in women, such as amygdale, anterior cingulate cortex (ACC), thalamus, hypothalamus, and insula, seem to exhibit decreased activation in menopause.³⁶ These areas are involved in the sexual drive.³⁶ Moreover, premenopausal women displayed higher FSFI-6 total scores, hence better sexual function, according to Latin-American studies.^{21,33}

The total FSFI-6 scores were positively correlated with family income, parity, and educational level, and inversely with age, peri-, and postmenopausal status, time of menopause onset, and total MRS score. In this research, higher parity was related to better sexual function, in opposition to the literature.^{37,38} The explanation to this discrepancy may rely on the fact that postmenopausal women have independent progeny at this time of life. Besides that, the correlation between parity and sexual function in the peri- and postmenopause had not yet been studied. We speculate that the number of children may confer emotional comfort, but other studies are necessary to verify which variables are associated with parity as a predictive factor of sexual function. Furthermore, high income and educational level can improve female sexual health status possibly due to higher self-care.

Our results agree with other previously published reporting in the literature^{8,19,37} that female aging and postmenopausal status apparently increase the risk for sexual dysfunction. Peri- and postmenopausal women presented higher total symptoms MRS score when compared with the premenopausal group, probably due to progressive estrogenic deficit.^{26,29}

As expected, the higher the MRS total score, the worst the female sexual function. The MRS total scores correlated inversely with the FSFI-6 total score, data similar to previously reported researches,^{19,37} demonstrating that multiple factors can contribute to female sexual function and well-being, such as frequency and intensity of menopausal

symptoms, social and cultural environment, and other psychological aspects. Besides, the Latin-American society is male-dominated, which can influence the female sexuality.³³

Therefore, our study confirmed that FSFI-6 is a short and rapid self-reporting instrument and can be combined with other instruments such as the MRS, which also displayed good internal consistencies. Considering these results, the Brazilian version of the FSFI-6 is a consistency tool for assessment of sexual function in middle-aged Brazilian women. In addition, we emphasize that this is a far-reaching study, involving 737 women, which consists in one of the largest populations for FSFI-6 validation up to the present moment.

Certain limitations should be considered. In general, information collected through self-reporting questionnaires lack diagnostic precision.³⁹ This may occur due to the difficulty of the subjects understanding the questions, although, as mentioned earlier, this was not the case in the present study (no difficulty was observed in the understanding of the proposed FSFI-6 version). Also, there is a difficulty measuring the extent to which symptoms leads to stress or suffering for each of the subjects. We believe that self-reporting instruments lead to more reliable answers, since the privacy of the participant is preserved when responding. Another important point is the lack of information on the sexual partners of the surveyed women, that should be addressed by a future research, in order to understand this factor involved in multimodal female sexual functioning.

Conclusion

The FSFI-6 was translated and adapted to the Brazilian culture and is a consistent tool for FSD screening in middle-aged women. Although there is no background data related to sexual function of Brazilian middle-aged women assessed with the FSFI-6, the results of our sample seem to indicate a high frequency of FSD. These results suggest that further investigations about prediction factors are needed, but it already demonstrates the need for more specific instruments for this evaluation in climacteric women.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Contributors

All authors listed above participated in the study to a significant extent. Mona Lúcia Dall'Agno, Charles Francisco Ferreira, Fernanda Vargas Ferreira, Faustino Ramón Pérez-López, and Maria Celeste Osório Wender worked on the conception, design, data collection, analysis and interpretation of the data, writing of the initial manuscript, and critical review of the final manuscript. All authors read and approved the submitted manuscript.

Acknowledgments

This work was supported by grants from the Fundo de Incentivo à Pesquisa e Eventos (FIPE—Hospital de Clínicas

de Porto Alegre (HCPA, Porto Alegre, RS, Brazil) and from the National Coordination for Improvement of Higher Education Personnel (CAPES, in the Portuguese acronym, Brazil).

The authors also thank all the researchers from the Climacteric and Menopause Research Group and Marco Aurélio Firmiano Scandalo, for conducting the English review of this manuscript. Additionally, we would like to express our deepest gratitude to the participants for their time and patience throughout this study.

References

- 1 Nappi RE, Lachowsky M. Menopause and sexuality: prevalence of symptoms and impact on quality of life. *Maturitas* 2009;63(02): 138–141. Doi: 10.1016/j.maturitas.2009.03.021
- 2 Higo M, Khan HTA. Global population aging: unequal distribution of risks in later life between developed and developing countries. *Glob Soc Policy* 2015;15:146–166. Doi: 10.1177/1468018114543157
- 3 Harlow SD, Gass M, Hall JE, et al; STRAW + 10 Collaborative Group. Executive summary of the Stages of Reproductive Aging Workshop + 10: addressing the unfinished agenda of staging reproductive aging. *Fertil Steril* 2012;97(04):843–851. Doi: 10.1016/j.fertnstert.2012.01.128
- 4 Avis NE, Brockwell S, Randolph JF Jr, et al. Longitudinal changes in sexual functioning as women transition through menopause: results from the Study of Women's Health Across the Nation. *Menopause* 2009;16(03):442–452. Doi: 10.1097/gme.0b013e3181948dd0
- 5 Blümel JE, Chedraui P, Baron G, et al; Collaborative Group for Research of the Climacteric in Latin America (REDLINC). Sexual dysfunction in middle-aged women: a multicenter Latin American study using the Female Sexual Function Index. *Menopause* 2009;16(06):1139–1148. Doi: 10.1097/gme.0b013e3181a4e317
- 6 Basson R. Women's sexual dysfunction: revised and expanded definitions. *CMAJ* 2005;172(10):1327–1333. Doi: 10.1503/cmaj.1020174
- 7 Nappi RE, Cucinella L, Martella S, Rossi M, Tiranini L, Martini E. Female sexual dysfunction (FSD): Prevalence and impact on quality of life (QoL). *Maturitas* 2016;94:87–91. Doi: 10.1016/j.maturitas.2016.09.013
- 8 Chedraui P, Pérez-López FR. Assessing sexual problems in women at midlife using the short version of the female sexual function index. *Maturitas* 2015;82(03):299–303. Doi: 10.1016/j.maturitas.2015.07.005
- 9 Basson R, Berman J, Burnett A, et al. Report of the international consensus development conference on female sexual dysfunction: definitions and classifications. *J Urol* 2000;163(03): 888–893. Doi: 10.1016/S0022-5347(05)67828-7
- 10 Carpenter JS, Jones SM, Studts CR, et al. Female Sexual Function Index short version: a MsFLASH item response analysis. *Arch Sex Behav* 2016;45(08):1897–1905. Doi: 10.1007/s10508-016-0804-5
- 11 Rosen R, Brown C, Heiman J, et al. The Female Sexual Function Index (FSFI): a multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Ther* 2000;26 (02):191–208. Doi: 10.1080/009262300278597
- 12 Hentschel H, Alberton DL, Capp E, Goldim JR, Passos EP. Validation of the female sexual function index (FSFI) for Portuguese language. *Rev HCPA*. 2007;27:10–14
- 13 Leite APL, Moura EA, Campos AAS, Mattar R, Souza E, Camano L. [Validation of the Female Sexual Function Index in Brazilian pregnant women]. *Rev Bras Ginecol Obstet* 2007;29:396–401. Doi: 10.1590/S0100-72032007000800003
- 14 Pacagnella RdeC, Vieira EM, Rodrigues OM Jr, Souza C. Adaptação transcultural do Female Sexual Function Index. *Cad Saude Publica* 2008;24(02):416–426. Doi: 10.1590/S0102-311 × 2008000200021

- 15 Pacagnella RdeC, Martinez EZ, Vieira EM. [Construct validity of a Portuguese version of the Female Sexual Function Index]. *Cad Saude Publica* 2009;25(11):2333-2344. Doi: 10.1590/S0102-311 x 2009001100004
- 16 Thiel RRC, Dambroz M, Palma PCR, Thiel M, Ricetto CLZ, Ramos Mdf. [Translation into Portuguese, cross-national adaptation and validation of the Female Sexual Function Index] *Rev Bras Ginecol Obstet* 2008;30:504-510. Doi: 10.1590/S0100-72032008001000005
- 17 Isidori AM, Pozza C, Esposito K, et al. Development and validation of a 6-item version of the female sexual function index (FSFI) as a diagnostic tool for female sexual dysfunction. *J Sex Med* 2010;7(03):1139-1146. Doi: 10.1111/j.1743-6109.2009.01635.x
- 18 Armeni A, Salazar-Pousada D, Andrade-Ponce SS, Tupacyupanqui-Mera JC, Pérez-López FR, Chedraui P. Reliability and validity of FSFI-6 in mid-aged Ecuadorian women. *Maturitas* 2017;103:93. Doi: 10.1016/j.maturitas.2017.06.016
- 19 Pérez-López FR, Fernández-Alonso AM, Trabalón-Pastor M, Vara C, Chedraui P; MenopAuse Risk Assessment (MARIA) Research Group. Assessment of sexual function and related factors in mid-aged sexually active Spanish women with the six-item Female Sex Function Index. *Menopause* 2012;19(11):1224-1230. Doi: 10.1097/gme.0b013e3182546242
- 20 Monterrosa-Castro A, Pérez-López FR, Ornat L. Sexual function assessment with the 6-item FSFI in postmenopausal Colombian women. *Maturitas* 2015;81:202. Doi: 10.1016/j.maturitas.2015.02.298
- 21 Chedraui P, Pérez-López FR, Sánchez H, et al. Assessment of sexual function of mid-aged Ecuadorian women with the 6-item Female Sexual Function Index. *Maturitas* 2012;71(04):407-412. Doi: 10.1016/j.maturitas.2012.01.013
- 22 Lee Y, Lim MC, Joo J, et al. Development and validation of the Korean version of the Female Sexual Function Index-6 (FSFI-6K). *Yonsei Med J* 2014;55(05):1442-1446. Doi: 10.3349/ymj.2014.55.5.1442
- 23 Pimenta F, Albergaria R, Gomes M, et al. Preliminary validating analysis of the Portuguese language 6-item Female Sexual Function Index (FSFI-6). *Maturitas* 2017;100:143-144. Doi: 10.1016/j.maturitas.2017.03.103
- 24 McCabe MP, Sharlip ID, Atalla E, et al. Definitions of Sexual Dysfunctions in Women and Men: A Consensus Statement From the Fourth International Consultation on Sexual Medicine 2015. *J Sex Med* 2016;13(02):135-143. Doi: 10.1016/j.jsxm.2015.12.019
- 25 Llana P, Fernández-Iñarrea JM, Arnott B, García-Portilla MP, Chedraui P, Pérez-López FR. Sexual function assessment in postmenopausal women with the 14-item changes in sexual functioning questionnaire. *J Sex Med* 2011;8(08):2144-2151. Doi: 10.1111/j.1743-6109.2011.02309.x
- 26 Heinemann K, Ruebig A, Potthoff P, et al. The Menopause Rating Scale (MRS) scale: a methodological review. *Health Qual Life Outcomes* 2004;2:45. Doi: 10.1186/1477-7525-2-45
- 27 Schneider HP, Heinemann LA, Rosemeier HP, Potthoff P, Behre HM. The Menopause Rating Scale (MRS): comparison with Kupperman index and quality-of-life scale SF-36. *Climacteric* 2000;3(01):50-58. Doi: 10.3109/13697130009167599
- 28 Schneider HP, Heinemann LA, Rosemeier HP, Potthoff P, Behre HM. The Menopause Rating Scale (MRS): reliability of scores of menopausal complaints. *Climacteric* 2000;3(01):59-64. Doi: 10.3109/13697130009167600
- 29 Heinemann LA, Potthoff P, Schneider HP. International versions of the Menopause Rating Scale (MRS). *Health Qual Life Outcomes* 2003;1:28. Doi: 10.1186/1477-7525-1-28
- 30 Wild D, Grove A, Martin M, et al; ISPOR Task Force for Translation and Cultural Adaptation. Principles of good practice for the translation and cultural adaptation process for Patient-Reported Outcomes (PRO) measures: report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value Health* 2005;8(02):94-104. Doi: 10.1111/j.1524-4733.2005.04054.x
- 31 Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol* 1993;46(12):1417-1432. Doi: 10.1016/0895-4356(93)90142-N
- 32 Byrne BM. *Structural Equation Modeling with AMOS: Basic Concepts, Applications, and Programming*. 3rd ed. London: Routledge; 2016
- 33 Sánchez SC, Chedraui P, Pérez-López FR, Ortiz-Benegas ME, Palacios-De Franco Y. Evaluation of sexuality in a Paraguayan mid-aged female urban population using the six-item Female Sexual Function Index. *Climacteric* 2016;19(03):256-260. Doi: 10.3109/13697137.2016.1151866
- 34 Wolpe RE, Zomkowski K, Silva FP, Queiroz APA, Sperandio FF. Prevalence of female sexual dysfunction in Brazil: A systematic review. *Eur J Obstet Gynecol Reprod Biol* 2017;211:26-32. Doi: 10.1016/j.ejogrb.2017.01.018
- 35 Clayton AH, Harsh V. Sexual function across aging. *Curr Psychiatry Rep* 2016;18(03):28. Doi: 10.1007/s11920-016-0661-x
- 36 Kim GW, Jeong GW. Menopause-related brain activation patterns during visual sexual arousal in menopausal women: An fMRI pilot study using time-course analysis. *Neuroscience* 2017;343:449-458. Doi: 10.1016/j.neuroscience.2016.12.010
- 37 Chedraui P, Pérez-López FR, Mezones-Holguin E, San Miguel G, Avila C; Collaborative Group for Research of the Climacteric in Latin America (REDLINC). Assessing predictors of sexual function in mid-aged sexually active women. *Maturitas* 2011;68(04):387-390. Doi: 10.1016/j.maturitas.2010.12.004
- 38 Hayes R, Dennerstein L. The impact of aging on sexual function and sexual dysfunction in women: a review of population-based studies. *J Sex Med* 2005;2(03):317-330. Doi: 10.1111/j.1743-6109.2005.20356.x
- 39 Abdo CH, Oliveira WM Jr, Moreira ED Jr, Fittipaldi JA. Prevalence of sexual dysfunctions and correlated conditions in a sample of Brazilian women—results of the Brazilian study on sexual behavior (BSSB). *Int J Impot Res* 2004;16(02):160-166. Doi: 10.1038/sj.jir.3901198