

KISS1/KISS1R and Breast Cancer: Metastasis Promoter

Stephania Guzman, PhD¹ Muriel Brackstone, MD, PhD² Frederic Wondisford, MD¹
 Andy V. Babwah, PhD^{3,4} Moshmi Bhattacharya, PhD^{1,4,5}

¹Department of Medicine, Robert Wood Johnson Medical School, Rutgers University, New Brunswick, New Jersey

²Department of Surgery, London Health Sciences Centre, London, Ontario, Canada

³Department of Pediatrics, Robert Wood Johnson Medical School, Rutgers University, New Brunswick, New Jersey

⁴Child Health Institute of New Jersey, New Brunswick, New Jersey

⁵Rutgers Cancer Institute of New Jersey, Rutgers University, New Brunswick, New Jersey

Address for correspondence Moshmi Bhattacharya, PhD, Department of Medicine, Robert Wood Johnson Medical School, Rutgers University, Clinical Academic Building, Room 7148, 125 Paterson Street, New Brunswick, NJ 08901 (e-mail: mb1722@rutgers.edu).

Semin Reprod Med 2019;37:197–206

Abstract

Kisspeptins (KPs), peptide products of the kisspeptin-1 (*KISS1*) gene, are the endogenous ligands for the KISS1 receptor, KISS1R, which is a G protein-coupled receptor. In many human tumors, *KISS1* functions as a metastasis-suppressor gene and KISS1/KISS1R signaling has antimetastatic and tumor-suppressor roles. On the contrary, emerging evidence indicates that the KP/KISS1R pathway plays detrimental roles in triple negative breast cancer (TNBC), the most difficult type of breast cancer to treat. TNBC patients initially respond to chemotherapy, but tumors acquire drug resistance and many patients relapse and die of metastases within a few years. In this review, we summarize recent developments in the understanding of the mechanisms by which KP/KISS1R signaling plays an adverse role in TNBC. This includes focusing on how KISS1R signaling regulates the cell cytoskeleton to induce tumor invadopodia formation and how KISS1R communicates with growth factor receptors such as the epidermal growth factor receptor, the receptor tyrosine kinase AXL, and transforming growth factor- β to promote cell invasion, metastasis, and drug resistance.

Keywords

- ▶ kisspeptin
- ▶ KISS1R
- ▶ triple negative breast cancer
- ▶ metastasis
- ▶ tumor invasion

Breast cancer is the most common noncutaneous malignancy among women worldwide. In 2018, over 2 million new cases and 627,000 deaths were reported globally with countries such as Belgium, France, Australia, and the United Kingdom having the highest rates.¹ In the United States in 2019, over 268,000 new cases have been reported with over 41,000 deaths.² Triple-negative breast cancer (TNBC) represents a particularly deadly form of breast cancer that occurs more commonly in women younger than 50 years, particularly those of African American and Hispanic descent.³ TNBC patients have poor prognosis and the tumors are typically high grade and metastatic. TNBC tumors are defined by a lack of estrogen receptor- α (ER α), progesterone receptor, and human epidermal growth factor 2 (HER2, also known as ErbB2).^{4,5} TNBC remains the most difficult breast cancer subtype to treat, as TNBC patients do not benefit from the

hormone receptor or HER2-targeting drugs used for other breast cancer subtypes. Additionally, these tumors become rapidly resistant to standard chemotherapies. Thus, an urgent need remains to identify new molecular targets for effective treatment of TNBC.

The *KISS1* gene is a metastasis-suppressor gene discovered in melanoma cells and its name reflects both its suppressor sequence and its place of discovery in Hershey, PA, the home of Hershey's Kisses chocolates.⁶ The *KISS1* gene encodes a 145 amino acid product that is proteolytically cleaved in the blood by furin and matrix metalloproteinases (MMPs) such as MMP-9 and the membrane type 1 matrix metalloprotease (MT1-MMP), into shorter kisspeptin peptides 10, 13, 14, or 54 amino acid in length.^{7,8} All kisspeptins have an amidated C-terminal end required for binding and activating the kisspeptin receptor (KISS1R), a G protein-

coupled receptor previously known as GPR54.⁸ KISS1R is a $G_{q/11}$ -coupled receptor, and upon ligand binding triggers phospholipase C activation, Ca^{2+} mobilization, and phosphorylation of mitogen-activated protein kinase (MAPK, ERK1/2) and p38.^{9–11} All KPs exhibit similar affinity for KISS1R; however, KP-10 is the agonist of choice for most studies.^{12–14} KISS1 and KISS1R are expressed in healthy breast tissue and play an essential role in mammary gland formation.^{9–11}

The KISS1/KISS1R signaling pathway has anti-metastasis/anti-tumorigenic roles in many cancers such as melanoma, pancreas, bladder, colorectal, ovary, and prostate, as previously reviewed.^{15–18} Mechanistically, the tumor-suppressive role has been linked to an inhibition of tumor invasion via a repression of MMP-9 expression and activity, and by inhibiting MAPK.^{19–22} Proteases such as MMP-9 promote tumor invasion by degrading the extracellular matrix (ECM). *KISS1* expression is lost in certain cancers such as colorectal and bladder cancer, due to epigenetic modifications of the *KISS1* promoter and this has been linked to poor patient survival.^{16,23,24}

In the last decade or so, several studies have shown that in marked contrast to its anti-tumor roles in most cancer, KISS1R signaling can promote cell migration and invasion in ER α -negative breast cancer such as TNBC. It is important to note that in the human carcinoma, MDA-MB-435 cells which were originally classified as a “breast cancer” line, *KISS1* was reported to act as a metastasis suppressor.²⁵ However, since then MDA-MB-435 cells have been found to express genes resembling melanoma cells and thus are no longer considered a robust breast cancer cell model.^{26–28} Here, we review the underlying mechanisms by which KISS1/KISS1R signal-

ing promotes metastasis in various breast cancer models such as established human breast cancer cell lines, animal models, and clinical samples. KISS1 appears to exhibit a dual role as an anti- and pro-tumor signaling molecule, as observed for c-MYC,²⁹ AMP-activated protein kinase (AMPK),³⁰ transforming growth factor- β (TGF- β),³¹ and NF κ B³² to name a few, emphasizing the importance of studying cancer in biological context.³³

Metastasis

Metastasis is the process by which cancer cells detach from the primary tumor, invade through the surrounding ECM, and migrate via the blood stream or lymphatics and colonize at secondary sites.³⁴ For breast tumors, these sites commonly include the brain, lungs, liver, and bone.³⁵ When tumors are confined to breast tissue, the 5-year survival rates exceed 90%.³⁶ However, once metastasis occurs, survival decreases rapidly depending on the extent and sites of tumor colonization.³⁷ Initial separation and spreading of cancer cells from the primary tumor involves a process known as epithelial-to-mesenchymal transition (EMT).^{38,39} Epithelial cells undergoing EMT lose their cell-to-cell adhesion and cell polarity as well as the expression of epithelial markers such as E-cadherin and gain the expression of mesenchymal markers such as MMP, N-cadherin, vimentin, and β -catenin (→Fig. 1). These cells also acquire a spindle-shaped morphology characteristic of fibroblasts and gain the ability to migrate and invade through the basement membrane into the surrounding ECM.³⁸ Cells which undergo EMT migrate away from primary tumors through cycling of adhesion/de-adhesion molecules regulated by




Fig. 1 Characteristics of epithelial cells undergoing epithelial-to-mesenchymal transition (EMT). Epithelial-like cells (shown on the left) have cell polarity, are nonmotile and noninvasive, and have cell-cell adhesion molecules such as E-cadherin. Mesenchymal-like cells (shown on the right) lack cell-cell adhesions and have a fibroblast-like morphology with front-rear polarity, increased motility, and invasiveness.




Fig. 2 Invadopodia formation by cancer cells. (a) Triple negative breast cancer (TNBC) cells form invadopodia, foot-like structures, upon stimulation by growth factors or kisspeptin. These are actin-rich membrane protrusions that stimulate cell invasion through the extracellular matrix (ECM) via accumulation of proteases such as membrane type 1 matrix metalloprotease (MT1-MMP). (b) Confocal image shown on the left showing the co-localization (yellow, arrows) of endogenous kisspeptin 1 receptor (KISS1R, red) and MT1-MMP (green) in TNBC MDA-MB-231 cell invadopodia. (c) KISS1R signaling promotes the activity of MT1-MMP via mitogen-activated protein kinase (MAPK, ERK1/2) to thereby promote ECM degradation. KISS1R via β -arrestin2 regulates the activity of invadopodia proteins; cortactin and cofilin regulate cytoskeleton.

integrins and by forming F-actin stress fibers through which myosin cycling initiates cell movement.⁴⁰

The ability of tumors to metastasize is dependent on the formation of invasive structures called invadopodia that mediate tumor cell entry into blood vessels (i.e., intravasate) or surrounding tissues (i.e., extravasate).⁴¹ Invadopodia are actin-rich, foot-like, adhesive membrane protrusions that form on tumor cell surfaces (—Fig. 2a). These are used to digest components of the ECM to create paths used for invading through tissues via the focal delivery of proteases such as MT1-MMP.⁴² Key invadopodia proteins such as cofilin and cortactin regulate invadopodia stabilization and maturation and the actin cytoskeleton for invadopodial function.⁴³ The capacity of tumors to form invadopodia is closely linked with the tumor's invasive and metastatic potential.^{44–46} Targeting pathways that regulate invadopodia formation are being tested clinically in many cancers, including TNBC, as a way of blocking metastasis.⁴⁷

TNBC: Current Treatment

The subtyping of breast cancers into molecular phenotype based on the expression of ER, progesterone receptor, and HER2 has transformed the way in which breast cancer is treated by permitting the tailoring of treatments specific to each intrinsic subtype.⁴⁸ Although the subtypes are defined by microarray-based gene expression, the clinical surrogate for

this is immunohistochemical testing of protein expression. In TNBC, the lack of ER, progesterone receptor, and HER2 expression limits therapeutic options to standard cytotoxic anthracycline and taxane-based chemotherapy with higher rates of treatment resistance and metastases. Thus, TNBC is the most challenging to treat and with the worst prognostic outcomes.⁵

With the absence of specific biomarkers in TNBC cells, researchers have been looking to exploit the rich epithelial-like and mesenchymal-like cancer stem cells by targeting pathways upregulated in these cells⁴⁹; however, these findings are in the early discovery phase. The main categories of therapies for TNBC currently being introduced into the clinical realm, albeit with limited data,⁵⁰ include (1) poly-ADP-ribosyl polymerase (PARP) inhibitors in the metastatic setting,⁵¹ given in combination therapy with DNA damaging alkylating chemotherapeutics (e.g., platinum) to patients with tumors lacking functional *BRCA1* or *BRCA2* genes (with resultant impaired ability to repair double-stranded DNA damage); (2) AKT inhibitor (ipatasertib), which has shown modest efficacy in progression-free survival⁵⁰; and (3) immunotherapy agents, for example, checkpoint inhibitors to programmed cell death protein 1 (PD-1) and programmed death-ligand 1 (PD-L1). PD-1 and PD-L1 bind to sites of DNA damage to target repair pathway activation. Immunotherapy drugs are given as single-agent therapies or combination therapies, with ongoing trials evaluating their efficacy in neoadjuvant, adjuvant, and metastatic settings.^{52,53} However, clinical trial results indicate

lower-than-expected impact on survival with checkpoint inhibitors in the metastatic setting and thus combination therapies with targets directed at EMT-related pathways are being explored.⁵⁴ Additionally, TNBC tumors exhibit extensive epigenetic alterations via DNA methylation and novel drug targets aimed at epigenetic re-programming such as BET bromo-domain inhibitors alone or in combination with immunotherapy are targeted for hypermethylated tumors and are also in clinical trials.^{5,55}

Molecular pathways that modulate tumorigenesis are deregulated in the aggressive TNBC phenotype. Although there appears to be some promising new therapies, there is a dire need for prognostic biomarkers and specific protein targets for drug development in TNBC. One significant potential target is KISS1R, and remainder of this review will highlight the role of KISS1R in regulating TNBC metastasis and chemotherapy resistance.

KISS1R: Metastasis Promoter

The first landmark study that revealed KISS1R promotes breast cancer metastasis came from Cho et al who used a mouse mammary tumor virus–polyoma virus middle T antigen (MMTV-PyMT) model of breast cancer metastasis.⁵⁶ In this model, the expression of the PyMT oncogene was driven by the MMTV promoter and transformation of the mammary epithelium led to the development of metastatic lesions in the lungs and lymph nodes, in addition to mammary adenocarcinomas.^{57,58} The study showed that *Kiss1r* haploinsufficiency (*Kiss1r*^{+/-}) led to delayed breast tumor initiation, growth, and metastasis. To show a direct effect on tumorigenesis, the authors conducted orthotopic injections of isolated mouse primary breast cancer MMTV-PyMT/*Kiss1r*^{+/-} cells into the mammary fat pads of mice and observed a reduction in primary tumor growth compared with mice injected with MMTV-PyMT/*Kiss1r*^{+/+} cells. Thus, loss of *Kiss1r* expression reduced tumor growth. Importantly, the authors verified that there were no pubertal defects of mammary gland development in the *Kiss1r* heterozygotes that would have resulted in impaired mammary tumor development. The authors also isolated breast tumor cells from primary tumors of PyMT/*Kiss1r*^{+/+} and PyMT/*Kiss1r*^{+/-} mice and found that *Kiss1r* heterozygosity negatively regulated tumor cell proliferation, cell motility, and invasion. Mechanistically, the study found that *Kiss1r* heterozygous (PyMT/*Kiss1r*^{+/-}) tumors displayed a significant reduction in *MMP-9* mRNA compared with wild-type (PyMT/*Kiss1r*^{+/+}) tumors. Additionally, KISS1R signaling was found to activate the small G protein, RhoA, a key regulator of the cytoskeleton necessary for cell migration and invasion downstream of Gαq activation, through Gαq-p63RhoGEF-RhoA signaling pathway (► Fig. 3).

KISS1/KISS1R: Regulation by ERα

Estrogen signaling via ERα critically regulates the development of the mammary ducts during puberty.⁵⁹ ERα signaling is an important regulator of breast carcinogenesis,⁶⁰ and silencing ERα has been shown to induce EMT.⁶¹ It is well

known that estradiol (E2) directly regulates *KISS1* expression in the hypothalamus. Specifically, in the arcuate neurons, ERα negatively regulates *KISS1*.⁶² Similarly, ERα has been shown to regulate KISS1 and KISS1R in breast carcinoma. Marot and colleagues showed that treatment of ERα-positive luminal breast cancer cells T47D and MCF7 with Tamoxifen, an ERα antagonist, stimulated the expression of *KISS1/KISS1R* mRNA.⁶³ Furthermore, women with ERα-positive breast tumors treated with Tamoxifen had high *KISS1* and *KISS1R* mRNA levels, which is associated with poor prognosis.⁶³ Clinically, *KISS1* and *KISS1R* levels are higher in ERα breast tumors compared with ERα+ tumors and this correlates with poor patient outcome.^{37,64} Thus, ERα signaling downregulates the expression of *KISS1/KISS1R* and when ERα is lacking, this results in an upregulation of *KISS1/KISS1R* in breast carcinoma. This was also observed to occur in TNBC cells. When ERα was overexpressed in TNBC MDA-MB-231 breast cancer cells, treatment of cells with E2 resulted in a decrease in *KISS1/KISS1R* mRNA and protein levels, compared with controls.^{19,63} In support of this, *KISS1/KISS1R* mRNA and protein levels were found to be upregulated in primary tumor biopsies compared with healthy breast tissue and KISS1 and KISS1R protein was found to be immunolocalized in invasive ductal carcinoma in TNBC tumors.⁶⁵ Taken together, the presence of ERα in breast epithelia appears to critically regulate the expression of KISS1/KISS1R.

KISS1/KISS1R: Malignant Transformation

The normal (nonmalignant) human MCF10A breast epithelial cells are a widely used model to study breast cell function as well as cell transformation.^{66,67} These cells express mammary gland-specific markers resembling normal human breast and form mammary acini that are capable of secreting milk.⁶⁸ Although MCF10A cells express KISS1R endogenously,¹⁹ they lack ERα.⁶⁶ This cell model is used commonly to study EMT-like processes in human breast.⁶⁹⁻⁷¹ KP-10 treatment or stable overexpression of KISS1R in MCF10A cells promoted an EMT-like event, resulting in the loss of E-cadherin from cell–cell junctions, the acquisition of mesenchymal markers such as N-cadherin and Snail/Slug and cells become migratory and invasive.¹⁹ This divulged that KISS1R signaling in breast epithelia lacking ERα can promote malignant transformation (► Fig. 3).

Metastatic breast cancer cells such as TNBC cells, MDA-MB-231, and Hs578T exhibit high expression of KISS1 and KISS1R in contrast to weakly invasive, ERα-positive breast cancer cells (T47D, MCF7) or nonmalignant breast MCF10A cells.⁶⁵ Depletion of KISS1R in TNBC cells using RNA interference reduced the expression of mesenchymal markers and stress-fiber formation and impaired invadopodia formation concurrent with their ability to migrate or invade.⁷²

Normal cells, in contrast to transformed cells, depend on cell to ECM contact to be able to grow and divide. In the absence of a substrate to attach to, normal cells undergo apoptosis.⁷³ In contrast, transformed cells such as tumors have the capability to grow and divide without the need of a substrate and this is a hallmark of carcinogenesis.⁷⁴ Thus,




Fig. 3 Kisspeptin/KISS1R signaling pathways in triple negative breast cancer (TNBC). In breast epithelia lacking estrogen receptor ($ER\alpha$) which negatively regulates KISS1/KISS1R expression, KISS1R promotes malignant transformation by inducing epithelial-to-mesenchymal transition (EMT), resulting in a decrease in E-cadherin expression and acquisition of mesenchymal markers and promoting stress fiber formation. Activation of KISS1R stimulates TNBC invasion by stimulating invadopodia formation and by activating the epidermal growth factor (EGFR) via β -arrestin2 and ERK/MMP-9-dependent pathways. KISS1R can also stimulate invasion via fibulin-3, which regulates MMP-9 activity. Lastly, KISS1R can signal via RhoA to regulate the cytoskeleton. Blue solid lines represent KISS1R interacting proteins; black dashed lines represent EGFR interacting proteins. DAG, diacylglycerol; ERK, extracellular signal-regulated kinase; IP_3 , inositol-(1,4,5)-trisphosphate; KP, kisspeptin; MMP-9, matrix metalloproteinase-9; PLC, phospholipase C; PKC, protein kinase C.

tumor cells can form colonies in an anchorage-independent manner and this ability can be evaluated using an anchorage-independent growth assay, also known as soft agar colony formation assay. Loss of KISS1R expression in TNBC cells reduced the ability of TNBC cells to form colonies on soft agar, further implicating a role for KISS1R in malignant transformation of breast epithelia.⁷²

KISS1/KISS1R: Tumor Cell Invasion

Using several TNBC cell lines (MDA-MB-231, Hs578t, SCP2, and SUM159) and $ER\alpha$ -negative cells (MCF10A, SKBR3),^{19,65,72,75–77} KISS1R signaling has been shown to promote cell migration and invasion, vital processes for metastasis. These have been observed in response to stimulating cells with KP-10. Using a *gain-of-function* model, the overexpression of human KISS1R in $ER\alpha$ -negative cells, normal MCF10A cells, and SKBR3 breast cancer cells (which have low levels of endogenous KISS1R) also induced cell migration and invasion.^{19,65} Furthermore, KISS1R overexpression in SKBR3 cells triggered tumor cells extravasation using the chick chorioallantoic membrane assay. KP-10

treatment of cells further increased the number of tumor cells that extravasated into the stroma, whereas a KISS1R antagonist (P-234) blocked the KP-10-mediated effect. This provided the first evidence that human KISS1R signaling regulates breast cancer invasion in an *in vivo* context.

In human TNBC cells, KISS1R signaling promotes cell invasion by several mechanisms (–Figs. 2 and 3). First, KISS1R stimulates invadopodia formation by activating key invadopodia proteins, cofilin, and MT1-MMP, through a β -arrestin2 and ERK1/2-dependent mechanisms.⁷⁸ Second, KISS1R activates the tyrosine kinase receptor epidermal growth factor receptor (EGFR or ErbB1) and stimulates TNBC invasion via β -arrestin2 and MMP-9.^{19,79} In fact, KISS1R directly binds EGFR in TNBC cells and KP-10 enhances this interaction.⁷⁹ Treatment of TNBC cells with the KISS1R antagonist, P-234, inhibited KP-10-induced invadopodia formation, cell invasion, and EGFR activation, implicating a role for KISS1R signaling in these processes.¹⁹ EGFR signaling promotes cell growth and survival⁸⁰ and TNBC frequently (50–70%) overexpresses EGFR.⁸¹ This overexpression correlates with a loss of estrogen responsiveness and a poor patient prognosis.⁸² EGFR expression itself is negatively

regulated by ER α .⁸³ Unfortunately, anti-EGFR therapies alone have not been effective in treating TNBC.⁸⁴

KISS1R is localized to the leading edge in lamellipodia of migratory breast cancer cells, co-localizing the actin-scaffolding protein, IQGAP1.¹⁹ KISS1R binding to IQGAP1 is necessary for EGFR activation that promotes TNBC invasion. However, KP-10 failed to activate EGFR or stimulate cell invasion of ER α -positive MCF7 and T47D cells, or upon expression of ER α in TNBC cells.¹⁹ This implies that ER α signaling keeps KISS1/KISS1R function in check, in addition to negatively regulating their expression.

Another way by which KISS1R signaling stimulates cell invasion is via the secreted ECM protein, fibulin-3, also known as epidermal growth factor (EGF)-containing fibulin-like extracellular matrix protein 1 (EFEMP1). Fibulin-3 is a secreted glycoprotein found in the ECM, known to regulate cell-matrix interactions, tissue remodeling, cell morphology, cell adhesion, and motility.⁸⁵ A recent study found using transcriptome analysis that fibulin-3 was expressed in cancer exosomes, which are extracellular vesicles released by cancer cells known to regulate organ-specific metastasis.⁸⁶ The fibulin-3 gene, *EFEMP1*, was found to be amplified in TNBC in contrast to other breast subtypes and plasma fibulin-3 levels were found to be elevated in TNBC patients compared with healthy subjects.⁷⁵ Furthermore, KISS1R signaling was found to induce fibulin-3 mRNA and protein expression as well as fibulin-3 secretion by TNBC cells. Downregulation of fibulin-3 in TNBC cells impaired KP-10-induced cell migration and invasion. Mechanistically, fibulin-3 was found to regulate TNBC cell invasion via ERK1/2 and MMP-9-dependent pathway, downstream of KISS1R activation (\rightarrow Fig. 3).

It is well known that *KISS1* mRNA levels are highly expressed in the placenta and plasma kisspeptin levels soar during pregnancy and remain elevated until parturition.⁸⁷⁻⁸⁹ Rasoulzadeh and colleagues examined the effect of placental-derived kisspeptin on breast cancer cell adhesion and invasion.⁷⁷ They collected conditioned media from human placental cells obtained from 11 placenta from healthy women with term pregnancy and undergoing Cesarean delivery. They found that placental-derived kisspeptins had a differential effect on TNBC (MDA-MB-231) cells compared with ER α -positive breast cancer (MCF-7) cells. In MDA-MB-231 cells, placental kisspeptins selectively reduced cell adhesion and promoted cell migration and invasion, which was inhibited upon treatment of cells with the KISS1R antagonist, P-234. In contrast, treatment of MCF-7 cells with placental kisspeptins had no effect on cell adhesion or invasion. Mechanistically, the study showed that placental kisspeptins increased MMP-9 expression and activity in MDA-MB-231 cells to thereby regulate cell invasion. This corroborates findings from previous studies that kisspeptins can differentially modulate invasiveness of breast cancer cells, depending on the ER α status.

Interestingly, KISS1 has been shown to function downstream of TGF- β to promote cell invasion in TNBC⁷⁶ (\rightarrow Fig. 4). Tian et al showed that TGF- β signaling via Smad2/p21 induced high expression of KISS1 in TNBC cells (MDA-MB-231 and SCP2) in contrast to ER α -positive (MCF-7) breast




Fig. 4 Transforming growth factor- β (TGF- β) promotes TNBC invasion via KISS1. TGF- β signaling via Smad2/p21 induces KISS1 expression, to promote TNBC invasion in an ERK/MMP-9-dependent manner. ERK, extracellular signal-regulated kinase; MMP-9, matrix metalloproteinase-9.

cancer cells.⁷⁶ Downregulation of KISS1 blocked TGF- β -induced TNBC invasion by stimulating MMP-9 expression and activity. The authors examined *KISS1* gene expression in 1,215 human breast tumors classified by ER status using the Cancer Genome Atlas (TCGA) database and observed higher *KISS1* expression in the aggressive basal-type, ER α -negative tumors versus the less aggressive ER α -positive tumors. Furthermore, using immunohistochemical analysis of KISS1 protein expression in a tumor microarray composed of normal breast tissue and breast carcinoma tissue from 48 patients, the study reported that positive lymph node status is associated with a higher KISS1 levels. Further studies are required to determine how this links with survival outcome and by what mechanisms TGF- β regulates KISS1 expression and whether KISS1R is involved in this pathway.

KISS1/KISS1R: Drug Resistance

TNBC patients have a lower survival rate following initial response to chemotherapy because tumors tend to develop chemoresistance and this limits the available options for previously treated patients.^{90,91} KISS1R signaling has been

shown to induce chemoresistance in TNBC cells and multiple ER α -negative breast cancer cells by two mechanisms.⁶⁵ First, KISS1R modulates the expression of the ATP-binding cassette drug efflux transporter, breast cancer resistance protein (BCRP), a major multidrug-resistant transporter in TNBC.^{92,93} Second, KP-10 treatment of cells activates the tyrosine kinase, AXL,⁶⁵ a binding partner of EGFR that is highly expressed in TNBC.^{94,95} AXL signaling is known to induce EMT as well as drug resistance^{96,97} and AXL inhibitors are currently in clinical trial in various cancers.^{98–100} KISS1R induces AXL transcription via enhanced binding of Sp1 to the AXL promoter. Knockdown of AXL using siRNA decreased the expression of EMT markers, but did not decrease BCRP expression, suggesting that KISS1R promotes drug resistance via two independent mechanisms. KISS1R overexpressing cells displayed reduced accumulation of the chemotherapeutic drug, doxorubicin, and displayed resistance to apoptosis as evidenced by a reduction of cleaved PARP, which occurs in the onset of apoptosis. In fact, KISS1R signaling promoted tumor cell survival by enhancing the expression of cell survival molecules such as AKT, ERK, and the antiapoptotic protein, survivin.⁶⁵ Depletion of AXL or antagonizing KISS1R signaling resensitized tumor cells to chemotherapy. Although the mechanisms by which KISS1R activates AXL are to be determined, this study revealed that KISS1R is a key regulator of TNBC drug resistance.

In a study of human head and neck squamous cell carcinoma (HNSCC), KISS1 has been shown to regulate drug resistance to the chemotherapeutic agent, cisplatin.¹⁰¹ In this cancer, *KISS1* functions as a metastasis suppressor and thus, *KISS1* expression is lost in patients with metastatic tumors compared with nonmetastatic tumors. The authors observed a profound decrease in *KISS1* mRNA expression in the cisplatin-resistant cell lines compared with the parental cells, but there was no change in KISS1R expression. Depletion of *KISS1* by siRNA in the parental HNSCC cells induced resistance to cisplatin. Conversely, overexpression of *KISS1* in HNSCC cells increased sensitivity of cisplatin-resistant cells to the chemotherapeutic in addition to inhibiting cell migration and proliferation. Importantly, using two preclinical xenograft models where human cells were injected into immunocompromised mice, the study showed that reexpression of *KISS1* in human cisplatin-resistant HNSCC cells decreased metastasis. Although the authors do not address the mechanism regulating the loss of *KISS1* in HNSCC, this study clearly revealed a role for *KISS1* in inhibiting cisplatin drug resistance as well as metastasis in HNSCC.

Conclusions and Future Perspective

Metastasis, the leading cause of cancer-related death, remains a major impediment for the treatment of cancer¹⁰² and several new therapeutic approaches are being investigated including targeting the tumor microenvironment, mitophagy, and immune therapy.^{103–105} In contrast to its original classification as a metastasis-suppressor gene in melanoma, this review provides information as to the state of our current knowledge on the proinvasive roles of *KISS1* in breast cancer.

Understanding the role of the KISS1/KISS1R pathway in regulating TNBC metastasis and drug resistance is still in its early period. Studies using preclinical xenograft models as well as patient-derived xenografts are required for a better understanding of the underlying molecular mechanisms by which KISS1R regulates cell invasion and whether targeting this pathway inhibits TNBC metastasis and drug resistance in vivo. Questions that remain to be answered include how does KISS1R communicate with other receptors such as growth factor receptors (e.g., TGF- β and AXL) and what role does KISS1R signaling play in other ER α -negative breast cancer subtypes (e.g., HER2-positive tumors).

Traditionally, the epigenetic regulation of gene expression in cancer is considered to be mediated via the inactivation of expression of tumor suppressor genes. Indeed, in colorectal and bladder cancer where *KISS1* functions as a tumor suppressor, *KISS1* expression is decreased in primary tumors due to epigenetic modifications.¹⁶ Recently, it is surfacing that epigenetic de-repression mechanisms such as DNA and histone demethylation are at play resulting in the overexpression of some oncogenes or cancer promoting genes in cancer and these can initiate cancer development and make tumors more aggressive and resistant to treatment.^{106–108} *KISS1* expression is elevated in patient TNBC primary tumors; however, whether *KISS1* is epigenetically regulated or undergoes miRNA regulation that results in changes in regulatory elements in TNBC needs further investigation.

In summary, biomarker identification to guide treatment decisions in TNBC is an unmet need, and an improved understanding of the molecular pathways that are deregulated in the early stages of TNBC can provide opportunities for targeted drug therapeutic intervention. TNBC cells overexpress KISS1 and secrete kisspeptin; however, it is yet to be determined whether plasma kisspeptin levels are altered in TNBC patients and whether this could serve as a potential biomarker for monitoring metastasis and disease recurrence. Despite the need for further investigations, it remains abundantly clear that KISS1/KISS1R plays important roles in TNBC invasion, metastasis, and drug resistance and this pathway represents a novel drug target.

Conflict of Interest

The authors declare no potential conflict of interest.

Acknowledgments

This work was supported in part with funds from the Canadian Institutes of Health Research (CIHR; grant PJT-14882, Rutgers Grant transfer agreement SDG) awarded to Moshmi Bhattacharya and Muriel Brackstone.

References

- 1 World Research Cancer Fund International. Available at: <https://www.wcrf.org/>. Accessed December 13, 2019
- 2 American Cancer Society. Available at: <https://www.cancer.org/>. Accessed December 13, 2019
- 3 Waks AG, Winer EP. Breast cancer treatment: a review. *JAMA* 2019;321(03):288–300
- 4 Foulkes WD, Smith IE, Reis-Filho JS. Triple-negative breast cancer. *N Engl J Med* 2010;363(20):1938–1948

- 5 Garrido-Castro AC, Lin NU, Polyak K. Insights into molecular classifications of triple-negative breast cancer: improving patient selection for treatment. *Cancer Discov* 2019;9(02):176–198
- 6 Lee JH, Miele ME, Hicks DJ, et al. KiSS-1, a novel human malignant melanoma metastasis-suppressor gene. *J Natl Cancer Inst* 1996;88(23):1731–1737
- 7 Biran J, Ben-Dor S, Levavi-Sivan B. Molecular identification and functional characterization of the kisspeptin/kisspeptin receptor system in lower vertebrates. *Biol Reprod* 2008;79(04):776–786
- 8 Kotani M, Detheux M, Vandenbogaerde A, et al. The metastasis suppressor gene KiSS-1 encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor GPR54. *J Biol Chem* 2001;276(37):34631–34636
- 9 Funes S, Hedrick JA, Vassileva G, et al. The KiSS-1 receptor GPR54 is essential for the development of the murine reproductive system. *Biochem Biophys Res Commun* 2003;312(04):1357–1363
- 10 Seminara SB, Messenger S, Chatzidaki EE, et al. The GPR54 gene as a regulator of puberty. *N Engl J Med* 2003;349(17):1614–1627
- 11 Kirby HR, Maguire JJ, Colledge WH, Davenport AP. International Union of Basic and Clinical Pharmacology. LXXVII. Kisspeptin receptor nomenclature, distribution, and function. *Pharmacol Rev* 2010;62(04):565–578
- 12 George JT, Veldhuis JD, Roseweir AK, et al. Kisspeptin-10 is a potent stimulator of LH and increases pulse frequency in men. *J Clin Endocrinol Metab* 2011;96(08):E1228–E1236
- 13 Messenger S, Chatzidaki EE, Ma D, et al. Kisspeptin directly stimulates gonadotropin-releasing hormone release via G protein-coupled receptor 54. *Proc Natl Acad Sci U S A* 2005;102(05):1761–1766
- 14 Min L, Soltis K, Reis AC, et al. Dynamic kisspeptin receptor trafficking modulates kisspeptin-mediated calcium signaling. *Mol Endocrinol* 2014;28(01):16–27
- 15 Cvetković D, Babwah AV, Bhattacharya M. Kisspeptin/KISS1R system in breast cancer. *J Cancer* 2013;4(08):653–661
- 16 Guzman S, Brackstone M, Radovick S, Babwah AV, Bhattacharya MM. KISS1/KISS1R in cancer: friend or foe? *Front Endocrinol (Lausanne)* 2018;9:437
- 17 Cho SG, Li D, Tan K, Siwko SK, Liu M. KiSS1 and its G-protein-coupled receptor GPR54 in cancer development and metastasis. *Cancer Metastasis Rev* 2012;31(3–4):585–591
- 18 Corno C, Perego P. KiSS1 in regulation of metastasis and response to antitumor drugs. *Drug Resist Updat* 2019;42:12–21
- 19 Cvetkovic D, Dragan M, Leith SJ, et al. KISS1R induces invasiveness of estrogen receptor-negative human mammary epithelial and breast cancer cells. *Endocrinology* 2013;154(06):1999–2014
- 20 Chen S, Chen W, Zhang X, Lin S, Chen Z. Overexpression of KiSS-1 reduces colorectal cancer cell invasion by downregulating MMP-9 via blocking PI3K/Akt/NF- κ B signal pathway. *Int J Oncol* 2016;48(04):1391–1398
- 21 Li N, Wang HX, Zhang J, Ye YP, He GY. KiSS-1 inhibits the proliferation and invasion of gastric carcinoma cells. *World J Gastroenterol* 2012;18(15):1827–1833
- 22 Zhang Y, Tang YJ, Li ZH, Pan F, Huang K, Xu GH. KiSS1 inhibits growth and invasion of osteosarcoma cells through inhibition of the MAPK pathway. *Eur J Histochem* 2013;57(04):e30
- 23 Cebrian V, Fierro M, Orenes-Piñero E, et al. KISS1 methylation and expression as tumor stratification biomarkers and clinical outcome prognosticators for bladder cancer patients. *Am J Pathol* 2011;179(02):540–546
- 24 Chen SQ, Chen ZH, Lin SY, Dai QB, Fu LX, Chen RQ. KISS1 methylation and expression as predictors of disease progression in colorectal cancer patients. *World J Gastroenterol* 2014;20(29):10071–10081
- 25 Lee JH, Welch DR. Suppression of metastasis in human breast carcinoma MDA-MB-435 cells after transfection with the metastasis suppressor gene, KiSS-1. *Cancer Res* 1997;57(12):2384–2387
- 26 Ross DT, Scherf U, Eisen MB, et al. Systematic variation in gene expression patterns in human cancer cell lines. *Nat Genet* 2000;24(03):227–235
- 27 Ellison G, Klinowska T, Westwood RF, Docter E, French T, Fox JC. Further evidence to support the melanocytic origin of MDA-MB-435. *Mol Pathol* 2002;55(05):294–299
- 28 Rae JM, Creighton CJ, Meck JM, Haddad BR, Johnson MD. MDA-MB-435 cells are derived from M14 melanoma cells—a loss for breast cancer, but a boon for melanoma research. *Breast Cancer Res Treat* 2007;104(01):13–19
- 29 Uribealago I, Benitah SA, Di Croce L. From oncogene to tumor suppressor: the dual role of Myc in leukemia. *Cell Cycle* 2012;11(09):1757–1764
- 30 Zadra G, Batista JL, Loda M. Dissecting the dual role of AMPK in cancer: from experimental to human studies. *Mol Cancer Res* 2015;13(07):1059–1072
- 31 Bachman KE, Park BH. Duel nature of TGF- β signaling: tumor suppressor vs. tumor promoter. *Curr Opin Oncol* 2005;17(01):49–54
- 32 Perkins ND. NF- κ B: tumor promoter or suppressor? *Trends Cell Biol* 2004;14(02):64–69
- 33 Bissell MJ. Context matters. *Trends Cancer* 2015;1(01):6–8
- 34 Mehlen P, Puisieux A. Metastasis: a question of life or death. *Nat Rev Cancer* 2006;6(06):449–458
- 35 Patanaphan V, Salazar OM, Risco R. Breast cancer: metastatic patterns and their prognosis. *South Med J* 1988;81(09):1109–1112
- 36 Welch DR, Steeg PS, Rinker-Schaeffer CW. Molecular biology of breast cancer metastasis. Genetic regulation of human breast carcinoma metastasis. *Breast Cancer Res* 2000;2(06):408–416
- 37 Martin TA, Watkins G, Jiang WG. KiSS-1 expression in human breast cancer. *Clin Exp Metastasis* 2005;22(06):503–511
- 38 Micalizzi DS, Farabaugh SM, Ford HL. Epithelial-mesenchymal transition in cancer: parallels between normal development and tumor progression. *J Mammary Gland Biol Neoplasia* 2010;15(02):117–134
- 39 Perl AK, Wilgenbus P, Dahl U, Semb H, Christofori G. A causal role for E-cadherin in the transition from adenoma to carcinoma. *Nature* 1998;392(6672):190–193
- 40 Hunter KW, Crawford NP, Alsarraj J. Mechanisms of metastasis. *Breast Cancer Res* 2008;10(Suppl 1):S2
- 41 Katt ME, Wong AD, Searson PC. Dissemination from a solid tumor: examining the multiple parallel pathways. *Trends Cancer* 2018;4(01):20–37
- 42 Buccione R, Caldieri G, Ayala I. Invadopodia: specialized tumor cell structures for the focal degradation of the extracellular matrix. *Cancer Metastasis Rev* 2009;28(1–2):137–149
- 43 Oser M, Condeelis J. The cofilin activity cycle in lamellipodia and invadopodia. *J Cell Biochem* 2009;108(06):1252–1262
- 44 Yamaguchi H. Pathological roles of invadopodia in cancer invasion and metastasis. *Eur J Cell Biol* 2012;91(11–12):902–907
- 45 Eckert MA, Lwin TM, Chang AT, et al. Twist1-induced invadopodia formation promotes tumor metastasis. *Cancer Cell* 2011;19(03):372–386
- 46 Gligorijevic B, Wyckoff J, Yamaguchi H, Wang Y, Rousos ET, Condeelis J. N-WASP-mediated invadopodium formation is involved in intravasation and lung metastasis of mammary tumors. *J Cell Sci* 2012;125(Pt 3):724–734
- 47 Meirson T, Gil-Henn H. Targeting invadopodia for blocking breast cancer metastasis. *Drug Resist Updat* 2018;39:1–17
- 48 Perou CM, Sørlie T, Eisen MB, et al. Molecular portraits of human breast tumours. *Nature* 2000;406(6797):747–752
- 49 Sulaiman A, McGarry S, Han X, Liu S, Wang L. CSCs in breast cancer—one size does not fit all: therapeutic advances in targeting heterogeneous epithelial and mesenchymal CSCs. *Cancers (Basel)* 2019;11(08):E1128
- 50 McCann KE, Hurvitz SA, McAndrew N. Advances in targeted therapies for triple-negative breast cancer. *Drugs* 2019;79(11):1217–1230

- 51 Robson M, Im SA, Senkus E, et al. Olaparib for metastatic breast cancer in patients with a germline BRCA mutation. *N Engl J Med* 2017;377(06):523–533
- 52 Schmid P, Adams S, Rugo HS, et al; IMpassion130 Trial Investigators. Atezolizumab and Nab-Paclitaxel in advanced triple-negative breast cancer. *N Engl J Med* 2018;379(22):2108–2121
- 53 Planes-Laine G, Rochigneux P, Bertucci F, et al. PD-1/PD-L1 targeting in breast cancer: the first clinical evidences are emerging. a literature review. *Cancers (Basel)* 2019;11(07):E1033
- 54 Soundararajan R, Fradette JJ, Konen JM, et al. Targeting the interplay between epithelial-to-mesenchymal-transition and the immune system for effective immunotherapy. *Cancers (Basel)* 2019;11(05):E714
- 55 Mohamad Hanif EA, Shah SA. Overview on epigenetic re-programming: a potential therapeutic intervention in triple negative breast cancers. *Asian Pac J Cancer Prev* 2018;19(12):3341–3351
- 56 Cho SG, Wang Y, Rodriguez M, et al. Haploinsufficiency in the prometastasis Kiss1 receptor Gpr54 delays breast tumor initiation, progression, and lung metastasis. *Cancer Res* 2011;71(20):6535–6546
- 57 Fantozzi A, Christofori G. Mouse models of breast cancer metastasis. *Breast Cancer Res* 2006;8(04):212
- 58 Lin EY, Jones JG, Li P, et al. Progression to malignancy in the polyoma middle T oncoprotein mouse breast cancer model provides a reliable model for human diseases. *Am J Pathol* 2003;163(05):2113–2126
- 59 Stingl J. Estrogen and progesterone in normal mammary gland development and in cancer. *Horm Cancer* 2011;2(02):85–90
- 60 Clarke RB, Anderson E, Howell A. Steroid receptors in human breast cancer. *Trends Endocrinol Metab* 2004;15(07):316–323
- 61 Al Saleh S, Al Mulla F, Luqmani YA. Estrogen receptor silencing induces epithelial to mesenchymal transition in human breast cancer cells. *PLoS One* 2011;6(06):e20610
- 62 Smith JT, Cunningham MJ, Rissman EF, Clifton DK, Steiner RA. Regulation of Kiss1 gene expression in the brain of the female mouse. *Endocrinology* 2005;146(09):3686–3692
- 63 Marot D, Bieche I, Aumas C, et al. High tumoral levels of Kiss1 and G-protein-coupled receptor 54 expression are correlated with poor prognosis of estrogen receptor-positive breast tumors. *Endocr Relat Cancer* 2007;14(03):691–702
- 64 Papaioconomou E, Lymperi M, Petraki C, et al. Kiss-1/GPR54 protein expression in breast cancer. *Anticancer Res* 2014;34(03):1401–1407
- 65 Blake A, Dragan M, Tirona RG, et al. G protein-coupled KISS1 receptor is overexpressed in triple negative breast cancer and promotes drug resistance. *Sci Rep* 2017;7:46525
- 66 Debnath J, Mills KR, Collins NL, Reginato MJ, Muthuswamy SK, Brugge JS. The role of apoptosis in creating and maintaining luminal space within normal and oncogene-expressing mammary acini. *Cell* 2002;111(01):29–40
- 67 Soule HD, Maloney TM, Wolman SR, et al. Isolation and characterization of a spontaneously immortalized human breast epithelial cell line, MCF-10. *Cancer Res* 1990;50(18):6075–6086
- 68 Underwood JM, Imbalzano KM, Weaver VM, Fischer AH, Imbalzano AN, Nickerson JA. The ultrastructure of MCF-10A acini. *J Cell Physiol* 2006;208(01):141–148
- 69 Rodriguez-Monterrosas C, Díaz-Aragon R, Leal-Orta E, Cortes-Reynosa P, Perez Salazar E. Insulin induces an EMT-like process in mammary epithelial cells MCF10A. *J Cell Biochem* 2018;119(05):4061–4071
- 70 Zhang J, Tian XJ, Zhang H, et al. TGF- β -induced epithelial-to-mesenchymal transition proceeds through stepwise activation of multiple feedback loops. *Sci Signal* 2014;7(345):ra91
- 71 Li S, Zhou J, Wu H, et al. Oncogenic transformation of normal breast epithelial cells co-cultured with cancer cells. *Cell Cycle* 2018;17(16):2027–2040
- 72 Goertzen CG, Dragan M, Turley E, Babwah AV, Bhattacharya M. KISS1R signaling promotes invadopodia formation in human breast cancer cell via β -arrestin2/ERK. *Cell Signal* 2016;28(03):165–176
- 73 Taddei ML, Giannoni E, Fiaschi T, Chiarugi P. Anoikis: an emerging hallmark in health and diseases. *J Pathol* 2012;226(02):380–393
- 74 Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011;144(05):646–674
- 75 Noonan MM, Dragan M, Mehta MM, et al. The matrix protein Fibulin-3 promotes KISS1R induced triple negative breast cancer cell invasion. *Oncotarget* 2018;9(53):30034–30052
- 76 Tian J, Al-Odaini AA, Wang Y, et al. KISS1 gene as a novel mediator of TGF β -mediated cell invasion in triple negative breast cancer. *Cell Signal* 2018;42:1–10
- 77 Rasoulzadeh Z, Ghods R, Kazemi T, et al. Placental kisspeptins differentially modulate vital parameters of estrogen receptor-positive and -negative breast cancer cells. *PLoS One* 2016;11(04):e0153684
- 78 Goertzen CF, Dragan M, Turley E, Bhattacharya M. KISS1R signaling promotes human triple negative breast cancer tumorigenesis and invadopodia formation via beta-arrestin2/ERK Cellular signalling. *Cell Signal* 2016;28(03):165–176
- 79 Zajac M, Law J, Cvetkovic DD, et al. GPR54 (KISS1R) transactivates EGFR to promote breast cancer cell invasiveness. *PLoS One* 2011;6(06):e21599
- 80 Jorissen RN, Walker F, Pouliot N, Garrett TP, Ward CW, Burgess AW. Epidermal growth factor receptor: mechanisms of activation and signalling. *Exp Cell Res* 2003;284(01):31–53
- 81 de Ruijter TC, Veeck J, de Hoon JP, van Engeland M, Tjan-Heijnen VC. Characteristics of triple-negative breast cancer. *J Cancer Res Clin Oncol* 2011;137(02):183–192
- 82 Toi M, Osaki A, Yamada H, Toge T. Epidermal growth factor receptor expression as a prognostic indicator in breast cancer. *Eur J Cancer* 1991;27(08):977–980
- 83 Flågåeng MH, Knappskog S, Haynes BP, Lønning PE, Mellgren G. Inverse regulation of EGFR/HER1 and HER2-4 in normal and malignant human breast tissue. *PLoS One* 2013;8(08):e74618
- 84 Sohn J, Liu S, Parinyanitikul N, et al. cMET activation and EGFR-directed therapy resistance in triple-negative breast cancer. *J Cancer* 2014;5(09):745–753
- 85 Timpl R, Sasaki T, Kostka G, Chu ML. Fibulins: a versatile family of extracellular matrix proteins. *Nat Rev Mol Cell Biol* 2003;4(06):479–489
- 86 Qadir F, Aziz MA, Sari CP, et al. Transcriptome reprogramming by cancer exosomes: identification of novel molecular targets in matrix and immune modulation. *Mol Cancer* 2018;17(01):97
- 87 Horikoshi Y, Matsumoto H, Takatsu Y, et al. Dramatic elevation of plasma metastin concentrations in human pregnancy: metastin as a novel placenta-derived hormone in humans. *J Clin Endocrinol Metab* 2003;88(02):914–919
- 88 Dhillon WS, Murphy KG, Bloom SR. The neuroendocrine physiology of kisspeptin in the human. *Rev Endocr Metab Disord* 2007;8(01):41–46
- 89 Bhattacharya M, Babwah AV. Kisspeptin: beyond the brain. *Endocrinology* 2015;156(04):1218–1227
- 90 Gu G, Dustin D, Fuqua SA. Targeted therapy for breast cancer and molecular mechanisms of resistance to treatment. *Curr Opin Pharmacol* 2016;31:97–103
- 91 Crown J, O'Shaughnessy J, Gullo G. Emerging targeted therapies in triple-negative breast cancer. *Ann Oncol* 2012;23(Suppl 6):vi56–vi65
- 92 Britton KM, Eyre R, Harvey IJ, et al. Breast cancer, side population cells and ABCG2 expression. *Cancer Lett* 2012;323(01):97–105
- 93 Natarajan K, Xie Y, Baer MR, Ross DD. Role of breast cancer resistance protein (BCRP/ABCG2) in cancer drug resistance. *Biochem Pharmacol* 2012;83(08):1084–1103

- 94 Meyer AS, Miller MA, Gertler FB, Lauffenburger DA. The receptor AXL diversifies EGFR signaling and limits the response to EGFR-targeted inhibitors in triple-negative breast cancer cells. *Sci Signal* 2013;6(287):ra66
- 95 Wu X, Zahari MS, Ma B, et al. Global phosphotyrosine survey in triple-negative breast cancer reveals activation of multiple tyrosine kinase signaling pathways. *Oncotarget* 2015;6(30):29143–29160
- 96 Antony J, Huang RY. AXL-driven EMT state as a targetable conduit in cancer. *Cancer Res* 2017;77(14):3725–3732
- 97 Wu X, Liu X, Koul S, Lee CY, Zhang Z, Halmos B. AXL kinase as a novel target for cancer therapy. *Oncotarget* 2014;5(20):9546–9563
- 98 Gorcea CM, Burthem J, Tholouli E. ASP2215 in the treatment of relapsed/refractory acute myeloid leukemia with FLT3 mutation: background and design of the ADMIRAL trial. *Future Oncol* 2018;14(20):1995–2004
- 99 Chia SK, Ellard SL, Mates M, et al. A phase-I study of lapatinib in combination with foretinib, a c-MET, AXL and vascular endothelial growth factor receptor inhibitor, in human epidermal growth factor receptor 2 (HER-2)-positive metastatic breast cancer. *Breast Cancer Res* 2017;19(01):54
- 100 Wakelee HA, Gettinger S, Engelman J, et al. A phase Ib/II study of cabozantinib (XL184) with or without erlotinib in patients with non-small cell lung cancer. *Cancer Chemother Pharmacol* 2017;79(05):923–932
- 101 Jiffar T, Yilmaz T, Lee J, et al. KiSS1 mediates platinum sensitivity and metastasis suppression in head and neck squamous cell carcinoma. *Oncogene* 2011;30(28):3163–3173
- 102 Akram M, Iqbal M, Daniyal M, Khan AU. Awareness and current knowledge of breast cancer. *Biol Res* 2017;50(01):33
- 103 Hinshaw DC, Shevde LA. The tumor microenvironment innately modulates cancer progression. *Cancer Res* 2019;79(18):4557–4566
- 104 Panigrahi DP, Praharaj PP, Bhol CS, et al. The emerging, multifaceted role of mitophagy in cancer and cancer therapeutics. *Semin Cancer Biol* 2019:S1044–579X(19)30115–4
- 105 Voorwerk L, Slagter M, Horlings HM, et al. Immune induction strategies in metastatic triple-negative breast cancer to enhance the sensitivity to PD-1 blockade: the TONIC trial. *Nat Med* 2019;25(06):920–928
- 106 Ehrlich M. DNA hypomethylation in cancer cells. *Epigenomics* 2009;1(02):239–259
- 107 Saito Y, Jones PA. Epigenetic activation of tumor suppressor microRNAs in human cancer cells. *Cell Cycle* 2006;5(19):2220–2222
- 108 Jeong HM, Kwon MJ, Shin YK. Overexpression of cancer-associated genes via epigenetic derepression mechanisms in gynecologic cancer. *Front Oncol* 2014;4:12