



# Basilar Invagination: Transoral Microsurgical Endoscopically-Controlled Odontoidectomy without Palatotomy in Extreme form of Basilar Invagination

Hischam Bassiouni<sup>1</sup>

<sup>1</sup>Departments of Neurosurgery, Klinikum Amberg and Klinikum Weiden, Bavaria, Germany

J Neurol Surg B 2021;82(suppl S1):S4–S5.

Address for correspondence Hischam Bassiouni, MD, PhD, Departments of Neurosurgery, Klinikum Amberg and Klinikum Weiden, Mariahilfbergweg 7, 92224 Amberg, Bavaria, Germany (e-mail: hibassiouni@yahoo.de).

## Abstract

**Objective** Transoral odontoidectomy in the treatment of basilar invagination is surgically challenging. Incision of the soft palate significantly increases rostral exposure of the clivus but is associated with a high incidence of speech and swallowing difficulties after surgery. We present a patient suffering from severe compression of the medulla oblongata due to an extreme form of basilar invagination treated successfully with the resection of dens via a transoral nasopharyngeal approach without palatotomy.

**Setting** Microsurgical endoscopic-assisted odontoidectomy through a transoral epipharyngeal approach was performed with subsequent craniocervical stabilization in a 21-year-old patient suffering from progressive myelopathy due to compression of the medulla oblongata and associated progressive syringomyelia.

**Results** The 21-year-old man was initially treated with suboccipital craniotomy and duroplasty in another institution. After initial improval he subsequently developed progressive ataxia, dysphagia, a bulbar speech, and weakness of the extremities. Beside ventral compression, he developed a secondary Chiari's malformation and a holospinal syringomyelia. Resection of the dens was successfully accomplished via a microsurgical transoral epipharyngeal endoscopic-controlled odontoidectomy without palatotomy. One week after odontoidectomy, posterior craniocervical stabilization was performed. All preoperative symptoms and signs improved significantly and the patient leads an independent life 4 years after odontoidectomy. On follow-up magnetic resonance imaging (MRI), the syringomyelia completely resolved.

**Conclusions** Palatotomy with its potential adverse effects can usually be avoided even for the treatment of extreme forms of basilar invagination.

The link to the video can be found at: <https://youtu.be/CBKE4n94W4g>.

## Keywords

- ▶ basilar invagination
- ▶ skull base
- ▶ transoral approach
- ▶ endoscopy
- ▶ odontoidectomy



## Conflict of Interest

None declared.

[www.thieme.com/skullbasevideos](http://www.thieme.com/skullbasevideos)

[www.thieme.com/jnlsbvideos](http://www.thieme.com/jnlsbvideos)


received  
March 24, 2019  
accepted after revision  
September 29, 2019  
published online  
March 20, 2020

DOI <https://doi.org/10.1055/s-0040-1701216>.  
ISSN 2193-6331.


© 2020. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



**Fig. 1** Upper row: initial MRI displaying basilar invagination with lower brainstem compression; MRI after suboccipital decompression in another institution showing secondary syringomyelia; and MRI after odontoidectomy with brainstem decompression and craniocervical stabilization. Lower row: bone window CT scan before and after odontoidectomy. MRI, magnetic resonance imaging.



**Fig. 2** Endoscopic intraoperative control demonstrating transoral epipharyngeal approach with soft palatal preservation and decompression of the brainstem after odontoidectomy.