

Acupuncture and electroacupuncture in the treatment of carpal tunnel syndrome: Systematic review

Marc Badia¹ Manel M. Santafé¹

¹Unit of Histology and Neurobiology, Department of Basic Medical Sciences, Faculty of Medicine and Health Sciences, Universitat Rovira i Virgili, Reus, Spain

Rev Fisioter Invasiva 2020;3:13–25.

Address for correspondence Manel M. Santafé, MD, PhD, Unidad de Histología y Neurobiología, Departamento de Ciencias Médicas Básicas, Facultad de Medicina y Ciencias de la Salud, Universitat Rovira i Virgili, Carrer de St Llorenç, 21, 43201, Reus, Tarragona, Spain (e-mail: manuel.santafe@urv.es).

Abstract

Introduction Carpal tunnel syndrome (CTS) is the most diagnosed compression neuropathy of the upper limb. In mild and moderate cases, the treatment is extensive and controversial, whereas severe cases receive surgical intervention.

Objective To analyze the scientific evidence on the effectiveness of treatment of CTS using acupuncture and electroacupuncture.

Methods The PRISMA declaration was followed. A literature search was performed using the following databases: Clinical Evidence BMJ, NICE, Cochrane Library, PubMed, PEDro, Science Direct, SciELO, Google Scholar. The search strategy used the terms “acupuncture”, “electroacupuncture” “carpal tunnel syndrome”. The inclusion criteria limited studies to English publications. The risk of bias was assessed for each study using the Cochrane scale. The level of evidence and the level of recommendation was determined using the SIGN scale.

Results The search retrieved 698 articles in total. After applying the inclusion and exclusion criteria, 21 articles were included. The level of evidence of all the articles was medium-high. The level of recommendation was medium, and the risk of bias was neutral, with a tendency towards low bias. The articles included revealed symptomatic and neurophysiological improvements, both peripheral as well as central, due to the cerebral response that occurs associated with the function of the median nerve.

Conclusion Acupuncture and electroacupuncture are a therapeutic option for mild to moderate CTS with a medium level of scientific evidence, tending towards a high level, and with a medium level of recommendation.

Keywords

- ▶ magnetic resonance
- ▶ median nerve
- ▶ neurography
- ▶ neuronal plasticity
- ▶ pain

Introduction

Carpal tunnel syndrome (CTS) is a neuropathy caused by the compression and traction of the median nerve in the carpal tunnel, delimited by the carpal bones and the transverse carpal ligament.¹ The clinical symptoms of CTS are syndromic, including paresthesia and nocturnal pain, normally related with poor positioning of the radius.² This syndrome is very common (4–5% of the general population), especially in people between 40–60 years old. Furthermore, it is more frequent in women (9.2%) than in men (6%).^{2–4}

The carpal tunnel is an osteofibrous outlet between the flexor retinaculum and the carpal bones. The roof of this tunnel is the fibrose carpal ligament, located deep in the intermediate region of the flexor retinaculum. In this space, nine tendons are found with their sheath and the median nerve. The nerve enters the tunnel in the midline or slightly radial to the same.⁵ The median nerve has sensory branches to supply the three radial digits and the radial half of the fourth finger, as well as the palmar cutaneous sensory branches, which innervate the skin of the palm,⁶ and the recurrent thenar branch. The sensitive area of the median

received
November 1, 2019
accepted
November 22, 2019

DOI <https://doi.org/10.1055/s-0040-1701432>.
ISSN 2386-4591.

Copyright © 2020 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms



nerve comprises the palmar aspect of the three radial fingers, and the radial half of the fourth finger, the dorsal aspect of the last two phalanges of the three first fingers, and the external half of the fourth finger. The motor branch passes by the distal aspect of the ligament, although in 20% of cases it is transligamentary, and because of this anatomic variant, an isolated compression exits at the level of the ligamentous outlet. The motor branch innervates the abductor pollicis brevis and the first lumbrical muscles.

Anatomically, two points of compression exist for the median nerve, one at the proximal border of the carpal tunnel, caused by wrist flexion and due to the change of thickness and stiffness between the forearm fascia and the proximal portion of the flexor retinaculum, and a second in the narrowest portion of the tunnel, at the level of the hook of the hamate. Pathological changes are thought to occur in the ligaments that surround the nerves, including alterations in the quantity and flexibility of the connective tissue as the source of the increase in pressure. This increase in pressure of the carpal tunnel causes the ischemic compression of the median nerve.⁷ In CTS, the nocturnal increase in pressure at the level of the tunnel may be due to several factors, such as the redistribution of fluids of the upper limbs in supine position, the lack of muscle pumping mechanisms which contribute towards draining the interstitial liquid in the carpal tunnel, the tendency to place the wrist in flexion, increasing pressure within the canal, an increased blood pressure during the second half of the night, and a drop in the level of cortisol.⁸

Median nerve entrapment can cause disorders of intraneural microcirculation, and injuries to the myelin sheath and the axon (altered function and dysfunction or damage from the point of compression in the distal direction).⁶ Additionally, there may be decomposition of the hematoencephalic barrier of the median nerve, formed by perineural cells and endothelial cells of endoneural capillaries.⁵ Together, these events can provoke an increased pressure of the endoneural liquid and the development of an intrafascicular edema,⁹ provoking an early impairment of the perineural and endoneural microcirculation, with fibrosis and secondary axonal demyelination. Demyelination is responsible for blocking the nerve transmission (neuropaxia) recorded under electromyography, and, if the compression persists, it can lead to Wallerian degeneration.¹ The rapid development of edema with epineural predominance leads to the inflammation of the nerve, limiting its ability to slide during movements. In response to this injury, there is an increased density of fibroblasts, of the size of the collagen fibers, an increase of type-III collagen, and vascular proliferation in the subsynovial connective tissue.¹⁰ This scar tissue is formed around the median nerve,¹¹ which, in turn, contributes to the nerve entrapment.

Treatment of CTS may be conservative or surgical. Conservative treatment may be active or passive. Passive treatments are based on the use of a wrist splint and infiltration of corticosteroids. Immobilization of the wrist in a neutral position of 0° of extension can provide symptom relief.¹² Furthermore, infiltration with corticosteroids is indicated in patients with mild to moderate clinical symptoms, offering a

marked relief mainly for pain.^{13,14} Active treatments are applied to patients with mild and moderate symptoms, and they include physical therapy, thermotherapy, electrotherapy, ultrasound therapy, laser, magnetotherapy, and manual techniques of mobilization of the carpal bones.

Acupuncture is reported to modulate the immune response and decrease inflammatory mediators (IL1,2 and FNT) which induce inducible cyclooxygenase (COX2). COX2 has an anti-inflammatory effect as it decreases the synthesis of prostaglandins.¹⁵ Furthermore, endogenous opioid activation has been suggested.¹⁶ In addition, electroacupuncture can develop a powerful inhibitory effect upon hyperalgesia in the short term, as well as an anti-inflammatory effect. It appears that the effects of electroacupuncture on hyperalgesia and inflammation can involve both the endocrine system and the nervous system.^{17,18} From the endocrine point of view, the participation of the suprarenal system has been suggested, therefore, producing an anti-inflammatory effect.^{19,20} In addition, at the spinal level, electroacupuncture inhibits the expression of Fos (transcription factor)^{19,20} as well as inducing an increased expression of opioid receptors mu and delta.^{21,22}

In the treatment of CTS, acupuncture is an easily accessed resource, with a low cost and without secondary effects, especially compared to more conventional treatments, such as infiltration of corticosteroids or surgical intervention. Thus, the aim of this systematic review was to evaluate the published evidence on the use of acupuncture for the treatment of CTS, and to assess the effectiveness of this treatment approach to therefore validate its use.

Methods

This systematic review has followed the criteria of the PRISMA declaration.

Selection criteria. The search was limited to:

- Case series, case studies, clinical trials, case-control studies, randomized controlled trials.
- Publications in English.
- Subjects with a medical diagnosis of carpal tunnel syndrome.
- Studies in which the intervention/therapy is acupuncture in carpal tunnel syndrome.

Studies which did not fulfill the previously described requisites were excluded, together with systematic reviews, as the aims of these were different to that of the present review.

Search strategy. A literature search was performed in the following databases: Cochrane Library, PubMed, Clinical Evidence BMJ, NICE, PEDro, Science Direct, SciELO, and Google Scholar. The last search was performed on April 21, 2019.

The search strategy used was based on the terms “acupuncture” and “electroacupuncture” and “carpal tunnel syndrome” as the key word related to CTS, to locate articles within the objectives of the review. Subsequently, the articles were selected according to eligibility criteria. Finally, systematic reviews were excluded. In all of the databases, the following terms were used: (acupuncture[MeSH Terms] OR

acupuncture[All Fields] OR *acupuncture therapy*[MeSH Terms] OR (*acupuncture*[All Fields] AND *therapy*[All Fields]) OR *acupuncture therapy*[All Fields]) AND (*carpal tunnel syndrome*[MeSH Terms] OR (*carpal*[All Fields] AND *tunnel*[All Fields] AND *syndrome*[All Fields]) OR *carpal tunnel syndrome*[All Fields]) AND (*electroacupuncture*[MeSH Terms] OR *electroacupuncture*[All Fields]) AND (*carpal tunnel syndrome*[MeSH Terms] OR (*carpal*[All Fields] AND *tunnel*[All Fields] AND *syndrome*[All Fields]) OR *carpal tunnel syndrome*[All Fields])).

Study selection. After performing the database search and downloading the total results, firstly, the title and abstract of all articles were read to exclude the articles which failed to fulfill the previously described eligibility criteria. Thereafter, records which were duplicated among the different databases were removed. The resulting articles were then read in-depth, to determine whether they fulfilled the eligibility criteria and to further screen those to be used for the data analysis. Subsequently, the selected studies were analytically read for data collection. A results table was created to determine the inclusion and exclusion criteria according to the specified eligibility criteria.

Risk of bias of individual studies. –Table 1 displays the risk of bias of each study, using the bias risk scale by Cochrane.

Level of evidence and degree of recommendation. The level of evidence and the degree of recommendation of each individual study is shown in –Table 2, based on the scale proposed by the Scottish Intercollegiate Guidelines Network (SIGN).

Synthesis of the results. To unify the results of the selected studies for the performance of the systematic review, a table was created, (–Table 3) featuring the study results, based on eight items: author and year, design, subjects and stage of illness, intervention, duration or follow-up, outcome measures, relevant findings, conclusions or commentaries. The results were then analyzed and interpreted as presented in the discussion.

Results

The aim of this review was to demonstrate the scientific evidence of the effects of acupuncture and electroacupuncture on the evolution and treatment of CTS.

Eight databases were consulted in this review, retrieving 698 articles, of which 20 were from the Clinical Evidence BMJ search engine, 45 were from NICE, 3 were retrieved from the Cochrane Library, 59 were from PubMed, 25 were from PEDro, 437 were from Science Direct, 1 was from SciELO, and, 108 were retrieved from Google Scholar.

Table 1 Evaluation of risk of bias of individual studies. Analysis of risk of bias of the studies included, using the COCHRANE tool

Article	Item 1	Item 2	Item 3	Item 4	Item 5	Item 6	Item 7
Schulman et al, 2008 ²³	(–)	(–)	(–)	(–)	(–)	(+)	(–)
Shokunbi, 2014 ²⁴	(–)	(–)	(–)	(–)	(–)	(–)	(–)
Napadow et al, 2004 ²⁵	(–)	(–)	(–)	(+)	(+)	(+)	(+)
Zailaa, 2010 ²⁶	(+)	(+)	(–)	(+)	(+)	(+)	(+)
Ding and Shen, 2013 ²⁷	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Wong et al, 2016 ²⁸	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Dimitrova et al, 2019 ²⁹	(+)	(+)	(–)	(+)	(+)	(+)	(–)
Chen et al, 2017 ³⁰	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Ural and Öztürk, 2017 ³¹	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Maeda et al, 2017. ³²	(–)	(–)	(–)	(+)	(+)	(+)	(+)
Hadianfard et al, 2015 ³³	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Maeda et al, 2013a ³⁴	(+)	(+)	(–)	(+)	(–)	(+)	(–)
Ho et al, 2014 ³⁵	(–)	(–)	(–)	(+)	(+)	(+)	(–)
Maeda et al, 2013b ³⁶	(+)	(+)	(–)	(–)	(+)	(+)	(–)
Khosrawi et al, 2012 ³⁷	(+)	(+)	(–)	(–)	(–)	(+)	(+)
Kummerdee and Kaewtong, 2010 ³⁸	(+)	(+)	(–)	(–)	(+)	(+)	(+)
Yang et al, 2011 ³⁹	(+)	(+)	(–)	(–)	(–)	(+)	(–)
Yang et al, 2009 ⁴⁰	(+)	(+)	(–)	(–)	(–)	(+)	(–)
Napadow et al, 2007a ⁴¹	(+)	(+)	(+)	(–)	(+)	(+)	(–)
Napadow et al., 2007b ⁴²	(+)	(+)	(+)	(–)	(+)	(+)	(–)
Yao et al, 2012 ⁴³	(+)	(+)	(–)	(+)	(+)	(–)	(–)

Notes: (+), low bias; (–), high bias. Item 1: generation of the random sequence. Item 2: concealed allocation. Item 3: type of blinding of participants and researchers. Item 4: type of blinding of evaluators. Item 5: follow-up and conclusions. Item 6: selective report of results. Item 7: other biases.

Table 2 Level of evidence

Database	Reference	Number of patients	Level of evidence	Level of recommendation
Google Scholar	Schulman et al, 2008 ²³	17	3	D
	Shokunbi, 2014 ²⁴	3	3	D
	Napadow, et al, 2004 ²⁵	11	2+	C
	Zailaa, 2010 ²⁶	22	2++	B
	Ding and Shen, 2013 ²⁷	38	2-	C
ScienceDirect	Wong et al., 2016 ²⁸	181	2+	C
PubMed	Dimitrova et al, 2019 ²⁹	60	2++	B
	Chen et al, 2017 ³⁰	60	2+	C
	Ural and Öztürk, 2017 ³¹	27	2+	C
	Maeda et al, 2017 ³²	80	2+	C
	Hadianfard et al, 2015 ³³	50	2+	C
	Maeda et al, 2013a ³⁴	67	2++	B
	Ho et al, 2014 ³⁵	26	2+	C
	Maeda et al, 2013b ³⁶	59	2++	B
	Khosrawi et al, 2012 ³⁷	72	2+	C
	Kumnerddee and Kaewtong, 2010 ³⁸	61	2+	C
	Yang et al, 2011 ³⁹	77	2+	C
	Yang et al, 2009 ⁴⁰	77	2+	C
	Napadow et al, 2007 ⁴¹	25	2++	B
Napadow et al, 2006 ⁴²	25	2++	B	
Yao et al, 2012 ⁴³	41	2+	C	

Legend: 1++ (meta-analysis of high quality, systematic review of clinical trials or high quality clinical trials with very low risk of bias), 1+ (well performed meta-analyses, systematic reviews of clinical trials or well performed clinical trials with a low risk of bias), 1- (meta-analysis, systematic reviews of clinical trials or clinical trials with a high risk of bias), 2++ (high quality systematic reviews of cohort studies or of cases and controls. Cohort studies or of cases and controls with a very low risk of bias and a high probability of establishing a causal relationship), 2+ (cohort or case and control studies which are well performed with low risk of bias and with a moderate probability of establishing a causal relation), 2- (cohort or case and control studies with a high risk of bias and a significant risk for a non-causal relation), 3 (non-analytic studies, such as case reports and case series), 4 (expert opinion), A (at least one meta-analysis, systematic review or clinical trial classified as 1++ and directly applicable to the target population of the guide; or a volume of scientific evidence comprised of studies classified as 1+ and with large consistence among them), B (a volume of scientific evidence comprised of studies classified as 2++, directly applicable to the target population of the guide and which demonstrate a great consistency among them; or scientific evidence extrapolated from studies classified as 1++ or 1+), C (a volume of scientific evidence comprised of studies classified as 2+, directly applicable to the target population of the guide and which demonstrate great consistency between them; or scientific evidence extrapolated from studies classified as 2++), D (level 3 or 4 scientific evidence; or scientific evidence extrapolated from studies classified as 2+), ✓ (recommended practice based on clinical experience and consensus of the research team).

As shown on ► **Figure 1**, in a first analysis, 622 articles were discarded by their title and abstract, failing to fulfill the inclusion criteria for this review, because of different aims or treatments. Of the 76 remaining studies, 39 were discarded (34 were duplicated in several databases, and 5 were discarded as these were systematic reviews). Of the 37 remaining articles selected, after an in-depth reading, 16 papers were discarded for the following reasons: 7 performed the study together with other therapies or drugs, 5 because of stimulation of the acupuncture point or the needle with other techniques that do not concern this study (laser, moxibustion, acupressure, cupping therapy), 3 did not fulfill the study aims because of performing other analyses, 1 performed the treatment by medical decision external to the study. Finally, 21 articles were selected for inclusion in this systematic review. The selected studies are analyzed in ► **Table 3**.

► **Table 2** shows that the level of evidence is medium with a tendency to be high and the level of recommendation is medium. Regarding the individual bias of the articles gathered in this systematic review (► **Table 1**; ► **Figure 2**), in general, there is a risk of neutral bias, with a tendency for low bias. To a large extent, this validates the results obtained in this study.

Of the articles gathered in this review, over 95% report favorable changes in response to the intervention with acupuncture and electroacupuncture, showing improvements in CTS (► **Table 3**).²³⁻⁴² The only study introduced which did not show favorable changes was the study by Yao et al (► **Table 3**).⁴³ These authors found that the group receiving intervention with acupuncture improved at 3 months in an identical manner to the placebo group. Several studies^{27,28,32,36,43} show that changes experienced by subjects via the application of acupuncture and electroacupuncture are maintained during long

Table 3 Analysis of results, key concepts and brief description of the articles

Author, year	Type of study	Number (N) of subjects and stage of evolution	Intervention	Duration of follow-up	Outcome measures	Relevant results	Conclusion
Schulman et al, 2008 ²³	Case series	N = 17. Nerve conduction impairment: mild, moderate and severe.	Needling for 20 minutes with electrical stimulation at 4 Hz.	(-)	Patient verbalization regarding the resolution of symptoms: complete, partial or without resolution.	Of 14 patients treated with acupuncture, 12 reported a partial or complete resolution of symptoms, and 2 did not obtain a resolution of symptoms.	Early intervention for CTS in patients with mild to moderate impairment provides better results than those of advanced cases.
Shokunbi 2014 ²⁴	Case study	3 subjects with over 3 months of evolution and refractory to other non-surgical treatments.	16 sessions of acupuncture lasting 30 minutes, twice a week during 8 weeks. Needling and ergonomic changes at work.	8 weeks.	SSS test at baseline and 8 weeks after.	SSS test: improvement. The symptoms of waking up in the middle of the night were eliminated. Reduction of the intensity of pain: 80% in 2 cases. Reduction of 50% of: paresthesia, numbness, and weakness.	Acupuncture treatment with ergonomic workplace changes improves the subjective symptoms of CTS.
Napadow et al, 2004 ²⁵	Case-control study	N = 11 subjects with positive mild to moderate CTS. Intervention group, N = 6; control group, N = 5.	5-week acupuncture protocol.	Assessment before the session, at 2 weeks and after 5 weeks.	Nerve conduction somatosensory evoked potential, and functional magnetic resonance before, during and after treatment. Boston questionnaire. Physical exam Phalen test, Tinel sign, and grip strength.	Functional magnetic resonance: activation of the postcentral gyrus of the contralateral fingers. For patients with CTS, activation after 5 weeks of acupuncture changed more for the 2nd and 3rd fingers than for the 5th finger. Boston questionnaire: decreasing the mean severity of the neuropathic symptoms of CTS.	CTS improved with acupuncture. Real acupuncture was superior to simulated acupuncture in peripheral and cerebral neurophysiology. Functional magnetic resonance can be used to monitor the cortical activation associated with a chronic neuropathy due to entrapment.
Zailaa, 2010 ²⁶	Case-control study	N = 22 subjects with documented history of CTS, assessed using magnetic resonance.	Application of a small dose of neutral liquid on acupuncture points.	The following six hours after the intervention with functional magnetic resonance assessment.	Structural images of the wrist in high resolution 4T of the median nerve. Grouped functional magnetic resonance data, BOLD-contrast imaging response.	All of the subjects reported a moderate sensation of electric tingling. Functional resonance; no significant BOLD-contrast imaging response was found to the stimulations on both points. Median nerve without structural changes.	Statistically significant reduction in the intensity of the signal and the inflammation within the median nerve after acupuncture.
Ding and Shen, 2013 ²⁷	Randomized controlled trial	N = 38 Intervention group: N = 19 Control group.	Group C: moxibustion performed on the Yangchi point. Group I: electric acupuncture on the Yangchi point.	3 months.	VAS assessments, SSS before and after treatment	VAS, SSS assessments: improvements compared to the same group before treatment in favor of the acupuncture group. 3 months later: the VAS scores did not show differences between both groups; the SSS scores showed differences for the	Acupuncture improves pain, numbness and motor activity in patients with CTS.

(Continued)

Table 3 (Continued)

Author, year	Type of study	Number (N) of subjects and stage of evolution	Intervention	Duration of follow-up	Outcome measures	Relevant results	Conclusion
						acupuncture group.	
Wong et al, 2016 ²⁸	Randomized controlled trial	N = 181. Electroacupuncture + nocturnal splinting; N = 90. Control group: N = 91; only nocturnal splinting.	Electroacupuncture (10–20 mA, 20–40 Hz, continuous wave) on the affected side in 20-minute sessions during 17 weeks, with a total of 13 sessions.	17 weeks.	Boston questionnaire; DASH questionnaire. Pain intensity, strength, fingertip pinch measured with a dynamometer of therapeutic pinch; Sensation of pain; Semmes-Weinstein monofilament examination; Dexterity: Moberg pick-up test (modified by Dellon).	Boston and DASH questionnaires: improvement. For pain intensity: reduction of pain. Dexterity, strength and sensation of pain: greater proportion of patients with improvement.	Electroacupuncture improves the symptoms of the disability function, dexterity and strength in a 17-week assessment, in combination with nocturnal splinting.
Dimitrova et al, 2019 ²⁹	Mechanical pilot study	N = 60. Mild to moderate CTS based on EMG. Group 1: manual acupuncture. Group 2: low frequency electroacupuncture. Group 3: high frequency electroacupuncture.	Needling on points of the pericardial nerve and heart associated with the median and ulnar nerves respectively. Two treatments: acupuncture on the median nerve-pericardium meridian and ulnar nerve-heart meridian, with one week in between.	2 weeks.	Assessment of sensory nerve conduction and qualitative sensory test (assessment of cold and vibration threshold); post-intervention in the median and ulnar nerve compared to preintervention measures.	Group 1: improvement of the threshold for detection of cold in the median nerve, but not in the healthy ulnar nerve. Group 2: recovery of action potentials of the sensory median nerve. Group 3: improvement of the sensation of cold of the median nerve and vibration with increased conduction velocity of the sensory and motor median nerves.	Acupuncture has a specific effect on the median and ulnar nerves of the forearm, which can be measured with a nerve conduction test and quantitative sensory test.
Chen et al, 2017 ³⁰	Case-control study	N = 60 subjects with mild to moderate CTS. Control group: N = 30; Case group: N = 30.	Case group: needling on the contralateral side and then needling on the affected side. Control group: acupuncture on the points of the affected side. 10-day treatment, with acupuncture once a day for 30 minutes.	10 days.	3 sessions. Electrophysiology of the median nerve. CTS questionnaire by Levine. Wrist assessment classified by the patients.	Effectiveness of 90% in the observation group, and of 70% in the control group. The velocity of the sensory conduction and the amplitude of the median nerve increased in both groups.	The combined treatment with the technique of homo and contralateral acupuncture achieves therapeutic effects in cases of mild to moderate CTS, which are superior to those of the regular needling technique.
Ural and Öztürk, 2017 ³¹	Case-control study	N = 27 women diagnosed with CTS by EMG. Control group: treatment with nocturnal splinting. Case group: treatment with nocturnal splinting plus acupuncture.	Acupuncture was performed during 10 sessions lasting 25 minutes each; 2 or 3 sessions per week.	4 weeks	Visual scale for the severity of the symptoms. Duruoz Hand Index and DASH questionnaire for hand function and disability. Electrophysiology: muscle action potential, speed of motor and sensory nerve conduction,	All of the parameters improved in both groups. The diameter of the median nerve decreased in the acupuncture group. The changes were greater in the acupuncture group than in the control group.	The diameter of the median nerve decreased after treatment with acupuncture, as well as the severity of the symptoms, the functions of the hand and the electromyographic measurements, which were more significant in the

Table 3 (Continued)

Author, year	Type of study	Number (N) of subjects and stage of evolution	Intervention	Duration of follow-up	Outcome measures	Relevant results	Conclusion
					distal motor latency. Ultrasound: diameter of the median nerve.		acupuncture group.
Maeda et al, 2017 ³²	Randomized clinical trial	N = 80: group of real local electroacupuncture on the affected hand; group of real electroacupuncture distal in the ankle contralateral to the affected hand; and group of simulated electroacupuncture with a non-penetrating sham needle.	The 3 groups underwent 16 sessions of electroacupuncture during 8 weeks.	3 months.	Boston questionnaire: scoring of pain and paresthesia at the beginning of the study, after treatment, and during the 3 months of follow-up. Determination of the sensory latency of the nerve. Functional magnetic resonance of the somatosensory cortex at the beginning of the study and after therapy.	The three groups displayed an improvement in the severity of symptoms. Neurophysiology: real local and distal electroacupuncture was superior to simulated electroacupuncture. The greatest improvement in the distance of cortical separation occurred between digits 2 and 3 in real electroacupuncture at 3 months of follow-up.	Acupuncture on local sites versus distal sites can improve the function of the median nerve at the wrist by somatotopic neuroplasticity, different at the level of the primary somatosensory cortex after therapy.
Hadianfard et al, 2015 ³³	Case-control study	N = 50 subjects with mild to moderate CTS. Ibuprofen group, N = 25. Acupuncture group, N = 25.	Both groups were submitted to nocturnal splinting. Acupuncture group: 8 sessions for 20 minutes twice a week during 4 weeks. The ibuprofen group received 400 mg of ibuprofen 3 times a day during 10 days.	1 month	VAS score, Boston questionnaire, BCQT FUNCT and BCQT SYMPT. Electrophysiology at the beginning and end of the treatment.	Significant improvements in both groups with greater significance in the acupuncture group. The scores on the BCTQ, FUNCT, global SYMPT, VAS, and electrophysiological findings were superior in the acupuncture group compared with the ibuprofen group.	Acupuncture could be an effective treatment for CTS.
Maeda et al, 2013a ³⁴	Case-control study	N = 67. CTS group, N = 37; group of healthy subjects N = 30.	Electroacupuncture at 2 Hz on the affected wrist and dominant hand in healthy subjects. Distal acupuncture on the contralateral leg in relation to the most affected hand with CTS or dominant in healthy subjects.	(-)	Data of structural image via weighted pulse sequence of multiple captures.	The brain response in both groups and acupuncture points included the activation of the contralesional somatosensory cortex. Difference between points of local and distal acupuncture for healthy subjects, but not for subjects with CTS. No correlation was found between distal acupuncture points for any of the groups.	The brain response to electroacupuncture differs among subjects with CTS and healthy subjects for the local stimulation of the acupuncture point.
Ho et al, 2014 ³⁵	Case-control study	N = 26. Acupuncture group, N = 15; electroacupuncture group, N = 11.	24 sessions of acupuncture for 15 minutes during 6 weeks.	6 weeks.	Short clinical questionnaire by LO and Chiang; electrophysiological	Improved scores for symptoms after treatment with electroacupuncture.	Electroacupuncture can improve symptoms.

(Continued)

Table 3 (Continued)

Author, year	Type of study	Number (N) of subjects and stage of evolution	Intervention	Duration of follow-up	Outcome measures	Relevant results	Conclusion
					assessment; Tinel sign.		
Khosrawi et al, 2012 ³⁶	Case-control study	N = 72. Control group (N = 32): nocturnal splinting, vitamins B1 and B6; Case group (N = 32): acupuncture.	Control group: intervention with nocturnal splinting, vitamin B1 and B6 and simulated acupuncture during 5 weeks. Case group: 8 sessions of acupuncture and nocturnal splinting during 4 weeks.	4 weeks.	Global score of clinical symptoms; SSS; electrophysiology.	Case group: improvement in the SSS and electrophysiology.	Acupuncture improves the symptoms of CTS.
Kumnerdee et al, 2010 ³⁸	Randomized clinical trial	N = 61 subjects with mild to moderate CTS. Acupuncture group; nocturnal splinting group.	Acupuncture group: 10 sessions of electroacupuncture twice a week; nocturnal splinting group: neutral wrist splint worn each night.	5 weeks.	BCTQ, SSS, FSS, VAS.	VAS score: decreased in the acupuncture group compared with the nocturnal splinting. There were no differences in the SSS and FSS between the groups.	Electroacupuncture reduced pain more than nocturnal splinting in cases of mild and moderate CTS.
Yang et al, 2011 ³⁹	Randomized controlled trial	N = 77. Mild to moderate idiopathic CTS confirmed with electrophysiology. Prednisone group, n = 39; acupuncture group, n = 38.	Prednisone group: 2 weeks with 20 mg of prednisone a day followed by 2 weeks of 10 mg of prednisone a day during 4 weeks; acupuncture group: 8 sessions during 4 weeks.	13 months.	Follow-up at 7 and 13 months using the assessment of global symptoms, SSS, repeated nerve conduction tests.	Acupuncture group compared to the prednisone group: improvement in SSS, distal motor latencies and distal sensory latencies. A significant correlation was observed between the SSS changed from month 13 to the baseline and all of the electrophysiological parameters, except the amplitude of the composed muscle action potential	Treatment with acupuncture in the short term improves mild to moderate idiopathic CTS in the long term.
Yang et al, 2009 ⁴⁰	Randomized controlled trial	N = 77. Patients with mild to moderate idiopathic CTS diagnosed with studies of nerve conduction. Acupuncture group: N = 38; prednisone group: N = 39.	Acupuncture group: 8 sessions during 4 weeks. Prednisone group: 20 mg a day during 2 weeks, and 10 mg a day during the 2 following weeks.	4 weeks.	SSS questionnaire on weeks 2 and 4. Neurophysiology at the end of the study.	SSS: a high percentage of improvement in both groups; however, without differences between both groups. The acupuncture group compared to the prednisone group: decrease in distal motor latency in week 4.	Treatment with acupuncture in the short term is as effective as prednisone in the short term for the treatment of mild to moderate CTS.
Napadow et al, 2007a ⁴¹	Case-control study	N = 25. CTS group: N = 13; Group of	Electroacupuncture during	5 weeks.	Functional magnetic resonance.	Real acupuncture activates the hypothalamus and	Patients with chronic pain respond to

Table 3 (Continued)

Author, year	Type of study	Number (N) of subjects and stage of evolution	Intervention	Duration of follow-up	Outcome measures	Relevant results	Conclusion
		healthy subjects: N = 12.	10 minutes at 2 Hz. Treatment 3 times a week during 3 weeks and 2 times a week during the last 2 weeks. Electroacupuncture in patients with CTS in the affected hand. Electroacupuncture in the dominant hand in healthy subjects.			deactivates the amygdala. Acupuncture inversely activates the hypothalamus and the amygdala. The hypothalamus response was positively correlated with the level of adapted cortical plasticity in patients with CTS.	acupuncture in a different way compared to healthy subjects via the limbic system, which includes the hypothalamus and the amygdala.
Napadow et al, 2007b ⁴²	Case-control study	N = 25. CTS group: N = 13; group of healthy subjects: N = 12.	Sensory stimulation during the imaging test. Electroacupuncture at 2 Hz during 10 minutes, 3 times a week, for the first three weeks, and twice a week during the remaining 2 weeks.	5 weeks	Functional magnetic resonance. Electrophysiology of the ulnar and median nerve. Grip strength, sensory threshold testing with Semmes-Weinstein monofilament examination and Phalen and Tinel tests. CTS test BCTQ.	Acupuncture improved all of the subjective and objective measures. Upon functional magnetic resonance, it was observed that acupuncture activated the cortical areas of the median nerve, but decreased the activation in the contralateral precentral and postcentral gyrus, in the prefrontal parietal cortex and the inferior dorso-lateral cortex.	Electroacupuncture provokes a stimulus of somatosensory conditioning.
Yao et al, 2012 ⁴³	Randomized controlled trial	N = 41. Acupuncture group: N = 21; simulated acupuncture (placebo): N = 20.	Nocturnal splints for both groups. 6 weekly sessions of 20 minutes of acupuncture on the affected and contralateral side.	3 months	Symptoms and function domains of the self-assessment scale of the carpal tunnel CTSAQ. Pinch strength. Combined sensory index.	Acupuncture group and placebo group: improvement in symptom domain 3 months after the last treatment.	Acupuncture is not a better treatment than simulated acupuncture when used with the splint for patients with mild to moderate CTS.

Abbreviations: BCTQ, Boston carpal tunnel syndrome questionnaire; BCQT FUNCT, functional status score; BCQT SYMPT, and symptom severity score; BOLD, blood-oxygen-level-dependent; CTS, carpal tunnel syndrome; CTSAQ, carpal tunnel syndrome assessment questionnaire; DASH, Disabilities of the Arm Shoulder and Hand questionnaire; EMG, electromyography; FSS, Functional Status Scale; FUNCT, functional status; SSS, Symptom Severity Scale; SYMPT, symptom severity; VAS, Visual Analog Scale.

Note: (-), not available.

periods of time after acupuncture, registering benefits up to 12 weeks later.

From the methodological point of view, over 50% of the studies showed improvements that were proven with electrophysiology (► **Table 3**).^{25,26,29–37,39,40,42} Hadianfar et al (► **Table 3**),³³ compared treatment with acupuncture and another with ibuprofen, obtaining a decrease of DSL (*distal sensory latency*) in the acupuncture group, whereas

DML (*distal motor latency*) significantly decreased in both groups. Similarly, Yang et al (► **Table 3**)⁴⁰ also compared acupuncture treatment with treatment using prednisone and in the group with acupuncture, a decrease in distal motor latency was obtained. Occasionally, electrophysiological studies of nerve conduction were combined with other imaging studies, for example the study by Napadow et al. (► **Table 3**)^{25,42} combined electrophysiological methods




Fig. 1 Study selection process. This figure represents the study selection process, including the selection criteria and reasons for exclusion.




Fig. 2 Graph of risk of bias. Review of the arguments of the authors of the studies included in this systematic review on each item of risk of bias, presented as percentages. Item 1: Generation of the random sequence. Item 2: Concealed allocation. Item 3: Type of blinding of participants and researchers. Item 4: Type of blinding of evaluators. Item 5: Follow-up and exclusions. Item 6: Selective report of results. Item 7: other biases.

(nerve conduction) with functional magnetic resonance, obtaining improved neurophysiological results, both peripheral as well as cerebral after treatment with acupuncture.

Over 25% of the articles included in this review performed a cerebral assessment using functional magnetic resonance, finding changes which underline the interaction of the effect of acupuncture and electroacupuncture in the improvement

of symptoms of patients with CTS (► **Table 3**).^{25,32,34,36,41,42} In addition, most of these studies also described differences regarding the group of real-simulated acupuncture, local-distal points, symptomatic-asymptomatic side, as well as changes in control subjects at the level of the brain (► **Table 3**).^{25,32,34,36,41,42} For example, Maeda et al (► **Table 3**)^{32,34} performed a functional magnetic resonance

during the intervention, showing that local electroacupuncture produced an activation of brain areas related with pain modulation, both directly and indirectly, and, however, simulated electroacupuncture did not show any response.³²

Discussion

Most articles gathered in this review reveal favorable changes in response to acupuncture and electroacupuncture in CTS. Several studies report that the changes experienced by subjects via the application of acupuncture and electroacupuncture are maintained during long periods of time after receiving acupuncture.

The maintained improvement of symptoms after real acupuncture found in the present systematic review is in line with the results of previous clinical trials⁴³ that relate this effect with the plasticity of the primary somatosensory cortex (S1).

Moreover, the reduction of the symptoms of CTS has been found to persist during months after the cessation of therapy with acupuncture.³⁹ These improvements are shown in studies measuring the latency of nerve conduction and cerebral neuroplasticity. It is known that intraneural blood flow is controlled by the sympathetic system;⁴⁴ therefore, it has been proposed that acupuncture may be able to modulate the blood flow of the vasa nervorum of the median nerve via antidromic vasodilation after stimulation of the dorsal spinal roots. Furthermore, using neuroimaging techniques, the S1 has been shown to be a part of the cerebral regions that regulate the outflow of the sympathetic system,⁴⁵ providing feedback to the process. A recent study in patients with chronic pain found an increased connectivity of the S1 and the cortex of the anterior/middle insula that was associated with reduced cardiovascular modulation.⁴⁶ On the peripheral level, this vascular effect could be mediated by the calcitonin gene-related peptide (CGRP).^{47,48} Ultrasound has been used to explore the vascularization of the median nerve trapped in the CTS,^{49,50} observing hypervascularization. This particular vascular situation has been interpreted as a compensatory response to ischemia within the tunnel. This compensation can be due to the decrease in the peripheral afferent, which triggers an increased central processing; therefore, the brain tries to compensate the slower signaling in the periphery by amplifying the signal in regions of somatosensory cerebral processing. A greater activation of the S1 cortex has been described and visualized using functional magnetic resonance⁵¹ or contralateral electrophysiological amplification⁵² after stimulation of the side affected by CTS, which confirms this hypothesis. In addition, it is known that the activity of the S1 cortex reflects discriminatory sensory information, such as the intensity of the sensation.^{53,54}

In several of the assessed studies simulated acupuncture was performed as a placebo in the control group. This indeed is a controversial subject. A previous systematic review affirmed that the evidence of the benefits of acupuncture in the treatment of CTS was unsatisfactory due to the poor methodological quality of the study.⁵⁵ In other words, simulated

procedures are not physiologically inert.^{56,57} Indeed, simulated acupuncture has been observed to reduce aversive symptoms more easily than a placebo pill.⁵⁸ Interestingly, analgesia can be obtained with both real and simulated electroacupuncture. The brain mechanisms that support this analgesia can differ substantially. One study by Maeda et al,³⁶ for example, described that the brain response in the combined-acupuncture group was more intense than that of the simulated-acupuncture group. The authors described an activity in specific brain regions in response to combined electroacupuncture that may be related to the reduction of pain. One study⁵⁹ also found that this cerebral response is also proposed as an objective marker to discriminate simulated acupuncture from real acupuncture in a more reliable manner than the subjective pain reports made by the patients.

The physiological mechanism of the action of electroacupuncture on CTS pain has also been proposed to occur via the central nervous system.^{60,61} Several studies^{62,63} demonstrate that electroacupuncture induces a large cerebral response in healthy adults. The study by Maeda et al³⁶ demonstrated that electroacupuncture was able to deactivate brain areas of the network in limbic mode in patients with CTS and in healthy subjects on a subthreshold level. Thus, this study by Maeda et al³⁶ differs from previous publications on this subject, possibly because different intensities were used, activation thresholds.

The reduction of pain after real combined electroacupuncture has been related with the activation of the supplementary motor area (SMA). The SMA is a cortical region that modulates the communication between the somatosensory and motor systems, and it has been found to be activated by painful stimulation, as seen on functional magnetic resonance,⁶⁴ and it also contributes to pain control.⁶⁵ A greater activation after combined electroacupuncture can reflect a greater transfer of somatosensory entrances induced by electroacupuncture to the motor system,⁶⁶ promoting a more normalized sensorimotor communication, compared with the sporadic input of diffuse paresthesia. Therefore, electroacupuncture, either local or distal, can also reduce pain by providing a regulated somatosensory entrance to the brain. Therefore, studies^{66,67} involving magnetic resonance have demonstrated that treatment with acupuncture can alter the brain activity and regulate the activity of the limbic system in patients with CTS.

Conclusions

The studies featured in the present systematic review have a medium to high level of evidence, with a neutral to low risk of bias, which, together, largely validate the present work. Most studies evaluated nerve conduction and functional magnetic resonance imaging, demonstrating beneficial effects of acupuncture and electroacupuncture in the evolution of CTS. Thus, acupuncture and electroacupuncture are treatment options for CTS that may provide clinical and neurophysiological improvements (peripheral and cerebral) that may be superior to those obtained through treatment merely based on the use of nocturnal splints and nonsteroidal anti-inflammatory drugs (NSAIDs).

Conflict of Interests

The authors have no conflict interests to declare.

References

- Alfonso C, Jann S, Massa R, Torreggiani A. Diagnosis, treatment and follow-up of the carpal tunnel syndrome: a review. *Neurol Sci* 2010;31(03):243–252
- Stevens JC; American Association of Electrodiagnostic Medicine. AAEM minimonograph #26: the electrodiagnosis of carpal tunnel syndrome. *Muscle Nerve* 1997;20(12):1477–1486
- Aroori S, Spence RA. Carpal tunnel syndrome. *Ulster Med J* 2008; 77(01):6–17
- Jablecki CK, Andary MT, So YT, Wilkins DE, Williams FH; AAEM Quality Assurance Committee. Literature review of the usefulness of nerve conduction studies and electromyography for the evaluation of patients with carpal tunnel syndrome. *Muscle Nerve* 1993;16(12):1392–1414
- MacDermid JC, Doherty T. Clinical and electrodiagnostic testing of carpal tunnel syndrome: a narrative review. *J Orthop Sports Phys Ther* 2004;34(10):565–588
- Dorwart BB. Carpal tunnel syndrome: a review. *Semin Arthritis Rheum* 1984;14(02):134–140
- Viikari-Juntura E, Silverstein B. Role of physical load factors in carpal tunnel syndrome. *Scand J Work Environ Health* 1999;25 (03):163–185
- Chammas M. Carpal tunnel syndrome. *Chir Main* 2014;33(02):75–94
- Millesi H, Zöch G, Rath T. The gliding apparatus of peripheral nerve and its clinical significance. *Ann Chir Main Memb Super* 1990;9(02):87–97
- Lundborg G. Intraneural microcirculation. *Orthop Clin North Am* 1988;19(01):1–12
- Ettema AM, Amadio PC, Zhao C, Wold LE, An KN. A histological and immunohistochemical study of the subsynovial connective tissue in idiopathic carpal tunnel syndrome. *J Bone Joint Surg Am* 2004; 86(07):1458–1466
- Lundborg G. Nerve entrapment. In: Lundborg G, editor. *Nerve injury and repair*. Philadelphia (PA): Churchill Livingstone; 1988: 102–148
- Page MJ, Massy-Westropp N, O'Connor D, Pitt V. Splinting for carpal tunnel syndrome. *Cochrane Database Syst Rev* 2012;7(07): CD010003
- Marshall S, Tardif G, Ashworth N. Local corticosteroid injection for carpal tunnel syndrome. *Cochrane Database Syst Rev* 2007;2(02): CD001554
- Boyer MI. Corticosteroid injection for carpal tunnel syndrome. *J Hand Surg Am* 2008;33(08):1414–1416
- Gong HS, Baek GH, Oh JH, Lee YH, Jeon SH, Chung MS. Factors affecting willingness to undergo carpal tunnel release. *J Bone Joint Surg Am* 2009;91(09):2130–2136
- Leyva Glez M, Palomina Carmenate E, Barrios Hdez J, Cruz Font J, Romero Hdez L. Acupuntura del Microsistema Su Jok y quiropraxia en el Síndrome del Túnel Carpiano. *Correo Científico Méd Holguín* 2011;15(01):
- Lao L, Zhang G, Wei F, Berman BM, Ren K. Electro-acupuncture attenuates behavioral hyperalgesia and selectively reduces spinal Fos protein expression in rats with persistent inflammation. *J Pain* 2001;2(02):111–117
- Lao L, Zhang RX, Zhang G, Wang X, Berman BM, Ren K. A parametric study of electroacupuncture on persistent hyperalgesia and Fos protein expression in rats. *Brain Res* 2004;1020(1-2):18–29
- Li A, Zhang RX, Wang Y, et al. Corticosterone mediates electroacupuncture-produced anti-edema in a rat model of inflammation. *BMC Complement Altern Med* 2007;7:27
- Zhang RX, Lao L, Wang X, et al. Electroacupuncture attenuates inflammation in a rat model. *J Altern Complement Med* 2005;11 (01):135–142
- Zhang RX, Lao L, Wang L, Liu B, Wang X, Ren K, Berman BM. Involvement of opioid receptors in electroacupuncture-produced anti-hyperalgesia in rats with peripheral inflammation. *Brain Res* 2004;1020(1-2):12–17
- Schulman RA, Liem B, Moroz A. Treatment of carpal tunnel syndrome with medical acupuncture. *Med Acupunct* 2008;20 (03):163–167
- Shokunbi GO. Acupuncture and Ergonomic interventions in the management of carpal tunnel syndrome among industrial workers-Case studies. *Nigerian J Med Rehabil* 2014;17(01):
- Napadow V, Liu J, Ryan A, et al. Effect of Acupuncture on Carpal Tunnel Syndrome with fMRI Assessment of Cortical Somatotopy. *Int Soc Magn Resonance Med* 2004;11(01):11–23
- Zailaa A. Magnetic Resonance Imaging Evaluation of Acupuncture Effects on Carpal Tunnel Syndrome. 2010
- Ding Q, Shen F. Effects of acupuncture and moxibustion of Yangchi point on carpal tunnel syndrome. *Int J Traditional Chin Med* 2013; 2:117–119
- Wong CHL, Wu XY, Wong W. Is adding electroacupuncture on top of nocturnal splinting an alternative treatment option for carpal tunnel syndrome patients? *AIMED* 2016;3(01):31–32
- Dimitrova A, Murchison C, Oken B. Local effects of acupuncture on the median and ulnar nerves in patients with carpal tunnel syndrome: a pilot mechanistic study protocol. *Trials* 2019;20(01):8
- Chen L, Xue L, Li S, Kang T, Chen H, Hou C. [Clinical research on mild and moderate carpal tunnel syndrome treated with contralateral needling technique at distal acupoints and acupuncture at local acupoints]. *Zhongguo Zhenjiu* 2017;37(05):479–482
- Ural FG, Öztürk GT. The Acupuncture Effect on Median Nerve Morphology in Patients with Carpal Tunnel Syndrome: An Ultrasonographic Study. *Evid Based Complement Alternat Med* 2017; 2017:7420648
- Maeda Y, Kim H, Kettner N, et al. Rewiring the primary somatosensory cortex in carpal tunnel syndrome with acupuncture. *Brain* 2017;140(04):914–927
- Hadianfard M, Bazrafshan E, Momeninejad H, Jahani N. Efficacies of Acupuncture and Anti-inflammatory Treatment for Carpal Tunnel Syndrome. *J Acupunct Meridian Stud* 2015;8(05): 229–235
- Maeda Y, Kettner N, Lee J, et al. Acupuncture Evoked Response in Contralateral Somatosensory Cortex Reflects Peripheral Nerve Pathology of Carpal Tunnel Syndrome. *Med Acupunct* 2013a25 (04):275–284
- Ho CY, Lin HC, Lee YC, et al. Clinical effectiveness of acupuncture for carpal tunnel syndrome. *Am J Chin Med* 2014;42(02):303–314
- Maeda Y, Kettner N, Lee J, et al. Acupuncture-evoked response in somatosensory and prefrontal cortices predicts immediate pain reduction in carpal tunnel syndrome. *Evid Based Complement Alternat Med* 2013b2013:795906
- Khosrawi S, Moghtaderi A, Haghight S. Acupuncture in treatment of carpal tunnel syndrome: A randomized controlled trial study. *J Res Med Sci* 2012;17(01):1–7
- Kumnerdee W, Kaewtong A. Efficacy of acupuncture versus night splinting for carpal tunnel syndrome: a randomized clinical trial. *J Med Assoc Thai* 2010;93(12):1463–1469
- Yang CP, Wang NH, Li TC, et al. A randomized clinical trial of acupuncture versus oral steroids for carpal tunnel syndrome: a long-term follow-up. *J Pain* 2011;12(02):272–279
- Yang CP, Hsieh CL, Wang NH, et al. Acupuncture in patients with carpal tunnel syndrome: A randomized controlled trial. *Clin J Pain* 2009;25(04):327–333
- Napadow V, Kettner N, Liu J, et al. Hypothalamus and amygdala response to acupuncture stimuli in Carpal Tunnel Syndrome. *Pain* 2007a130(03):254–266
- Napadow V, Liu J, Li M, et al. Somatosensory cortical plasticity in carpal tunnel syndrome treated by acupuncture. *Hum Brain Mapp* 2007b28(03):159–171

- 43 Yao E, Gerritz PK, Henricson E, et al. Randomized controlled trial comparing acupuncture with placebo acupuncture for the treatment of carpal tunnel syndrome. *PM R* 2012;4(05):367–373
- 44 Beissner F, Meissner K, Bär KJ, Napadow V. The autonomic brain: an activation likelihood estimation meta-analysis for central processing of autonomic function. *J Neurosci* 2013;33(25):10503–10511
- 45 Kim J, Loggia ML, Cahalan CM, et al. The somatosensory link in fibromyalgia: functional connectivity of the primary somatosensory cortex is altered by sustained pain and is associated with clinical/autonomic dysfunction. *Arthritis Rheumatol* 2015;67(05):1395–1405
- 46 Padua L, Coraci D, Erra C, Pazzaglia C, Paolasso I, Loreti C, Caliandro P, Hobson-Webb LD. Carpal tunnel syndrome: clinical features, diagnosis, and management. *Lancet Neurol* 2016;15:1273–1284
- 47 Sato A, Sato Y, Uchida S. Blood flow in the sciatic nerve is regulated by vasoconstrictive and vasodilative nerve fibers originating from the ventral and dorsal roots of the spinal nerves. *Neurosci Res* 1994;21(02):125–133
- 48 Ooi CC, Wong SK, Tan AB, et al. Diagnostic criteria of carpal tunnel syndrome using high-resolution ultrasonography: correlation with nerve conduction studies. *Skeletal Radiol* 2014;43(10):1387–1394
- 49 Ghasemi-Esfe AR, Khalilzadeh O, Mazloumi M, et al. Combination of high-resolution and color Doppler ultrasound in diagnosis of carpal tunnel syndrome. *Acta Radiol* 2011;52(02):191–197
- 50 Napadow V, Dhond R, Kennedy D, Hui KK, Makris N. Automated brainstem co-registration (ABC) for MRI. *Neuroimage* 2006;32(03):1113–1119
- 51 Tecchio F, Padua L, Aprile I, Rossini PM. Carpal tunnel syndrome modifies sensory hand cortical somatotopy: a MEG study. *Hum Brain Mapp* 2002;17(01):28–36
- 52 Coghill RC, Sang CN, Maisog JM, Iadarola MJ. Pain intensity processing within the human brain: a bilateral, distributed mechanism. *J Neurophysiol* 1999;82(04):1934–1943
- 53 Derbyshire SW, Jones AK, Gyulai F, Clark S, Townsend D, Firestone LL. Pain processing during three levels of noxious stimulation produces differential patterns of central activity. *Pain* 1997;73(03):431–445
- 54 Sim H, Shin BC, Lee MS, Jung A, Lee H, Ernst E. Acupuncture for carpal tunnel syndrome: a systematic review of randomized controlled trials. *J Pain* 2011;12(03):307–314
- 55 Lund I, Lundeberg T. Are minimal, superficial or sham acupuncture procedures acceptable as inert placebo controls? *Acupunct Med* 2006;24(01):13–15
- 56 Linde K, Niemann K, Schneider A, Meissner K. How large are the nonspecific effects of acupuncture? A meta-analysis of randomized controlled trials. *BMC Med* 2010;8:75
- 57 Kaptchuk TJ, Stason WB, Davis RB, et al. Sham device v inert pill: randomised controlled trial of two placebo treatments. *BMJ* 2006;332(7538):391–397
- 58 Harris RE, Zubieta JK, Scott DJ, Napadow V, Gracely RH, Clauw DJ. Traditional Chinese acupuncture and placebo (sham) acupuncture are differentiated by their effects on mu-opioid receptors (MORs). *Neuroimage* 2009;47(03):1077–1085
- 59 Napadow V, Lee J, Kim J, et al. Brain correlates of phasic autonomic response to acupuncture stimulation: an event-related fMRI study. *Hum Brain Mapp* 2013;34(10):2592–2606
- 60 Claunch JD, Chan ST, Nixon EE, et al. Commonality and specificity of acupuncture action at three acupoints as evidenced by fMRI. *Am J Chin Med* 2012;40(04):695–712
- 61 Yan B, Li K, Xu J, et al. Acupoint-specific fMRI patterns in human brain. *Neurosci Lett* 2005;383(03):236–240
- 62 Zhang WT, Jin Z, Luo F, Zhang L, Zeng YW, Han JS. Evidence from brain imaging with fMRI supporting functional specificity of acupoints in humans. *Neurosci Lett* 2004;354(01):50–53
- 63 Hui KK, Liu J, Makris N, et al. Acupuncture modulates the limbic system and subcortical gray structures of the human brain: evidence from fMRI studies in normal subjects. *Hum Brain Mapp* 2000;9(01):13–25
- 64 Huang W, Pach D, Napadow V, et al. Characterizing acupuncture stimuli using brain imaging with fMRI—a systematic review and meta-analysis of the literature. *PLoS One* 2012;7(04):e32960
- 65 Peyron R, Laurent B, García-Larrea L. Functional imaging of brain responses to pain. A review and meta-analysis (2000). *Neurophysiol Clin* 2000;30(05):263–288
- 66 Shinoura N, Suzuki Y, Yamada R, Kodama T, Takahashi M, Yagi K. Fibers connecting the primary motor and sensory areas play a role in grasp stability of the hand. *Neuroimage* 2005;25(03):936–941
- 67 Asghar AU, Green G, Lythgoe MF, Lewith G, MacPherson H. Acupuncture needling sensation: the neural correlates of deqi using fMRI. *Brain Res* 2010;1315:111–118