

## Case Report

# MANAGEMENT OF ACUTE CORNEAL HYDROPS SECONDARY TO KERATOCONUS BY DESCEMETOPEXY USING INTRACAMERAL PERFLUOROPROPANE (C<sub>3</sub>F<sub>8</sub>) – A CASE REPORT

Vijay Pai<sup>1</sup>, Jayaram Shetty<sup>2</sup>, Hrishikesh Amin<sup>3</sup>, S. Bhat<sup>4</sup>, Divya Lakshmi<sup>5</sup>  
<sup>1</sup>Professor, <sup>2</sup>Prof. & Head, <sup>3</sup>Professor, <sup>4</sup>Assoc. Professor, <sup>5</sup>Sr. Resident,  
 Dept. of Ophthalmology, K.S. Hegde Medical Academy, Deralakatte, Mangalore - 575 018.

Correspondence:

Vijay Pai,

Professor, Dept. of Ophthalmology, K.S. Hegde Medical Academy, Deralakatte, Mangalore – 575018.

E-mail : kshemaophthal@gmail.com, eyevijay@yahoo.com

## Abstract :

Keratoconus is a clinical term used to describe bilateral non-inflammatory corneal ectasia in its axial part due to which cornea assumes a conical shape<sup>1</sup>. The onset of keratoconus is generally at the age of puberty, and progresses over a period of 10-20 years<sup>2,3</sup>. The treatment of Keratoconus is rarely an emergency, with the exception of corneal hydrops resulting from rupture of the Descemet's membrane. This may be the common mode of presentation in patients with associated developmental delay, probably related to habitual ocular massage<sup>4,5</sup>.

Keywords : Keratoconus, acute hydrops, descemetopexy, C<sub>3</sub>F<sub>8</sub>

## Case Report :

A 21 year old girl presented to the cornea services of Justice K.S. Hegde Charitable Hospital, Mangalore with complaints of sudden decrease in vision in the right eye since one week. She was a known case of developmental delay and had a past history of seizures. On examination, her visual acuity in the right eye was limited to light perception, and that in the left eye was counting fingers at 3 metres (-5.0sph/-2.0cyl 20°). Slit-lamp examination of the right eye revealed central corneal ectasia with dense stromal oedema involving 8mm of the cornea, with a stromal cyst [Fig 1(a) & 1(b)] ; Rest of the anterior segment could not be visualised. The slit-lamp examination of the left eye showed central corneal ectasia with Vogt's striae, with the rest of the anterior segment being unremarkable [Fig 1(c)]. The patient was not co-operative for keratometry, pachymetry and intra-ocular pressure measurements. B-scan of the right eye showed a normal posterior segment, whereas dilated fundus examination of the left eye was within normal limits.

Based on the above clinical findings, a diagnosis of bilateral keratoconus with right eye acute hydrops was made.

On the subsequent day, the patient was taken up for

Descemetopexy under intravenous sedation, after obtaining an informed consent. Preoperatively, the pupil was constricted using 2% Pilocarpine eye drops, 1 drop every 15 min, 1 hour prior to the surgery, not only to avoid injury to the lens, but also to prevent the gas from escaping behind the iris. Anterior chamber paracentesis was done following which 0.1ml of gaseous cocktail (3 parts of C<sub>3</sub>F<sub>8</sub> and 1 part of filtered air) was injected [Fig 2].

Post-operatively, the patient was advised to remain in supine position. On the 1<sup>st</sup> post-operative day, approximately 2/3<sup>rd</sup> of the anterior chamber contained the gas and the stromal oedema had decreased considerably [Fig 3(a)]. She was started on antibiotic-steroid combination 4 times a day and aqueous suppressants.

On the 3<sup>rd</sup> day, the gas had expanded to fill the anterior chamber entirely. The stromal oedema now, was approximately 5mm in diameter [Fig 3(b)]. The intra-ocular pressure was normal and there were no signs of pupillary block.

At the end of one week, the stromal oedema was now, only about 3mm, however, intrastromal cyst persisted [Fig 3(c)]. There were still no signs of pupillary block and the patient



Figure 1(a)



Figure 1(b)



Figure 1(c)



Figure 2



Figure 3(a)



Figure 3(b)



Figure 3(c)

was asked to follow-up after one week.

Discussion :

Corneal ectatic disorders like keratoconus<sup>6</sup>, keratoglobus<sup>7</sup> and pellucid marginal degeneration<sup>8</sup> can be complicated with the occurrence of acute hydrops. Incidence of acute hydrops can vary from 2.8%<sup>9</sup> to 3.15%<sup>10</sup> in keratoconus. In most instances, treatment of acute hydrops has been conservative and non-specific such as use of topical antibiotic-steroids, cycloplegics and hypertonic saline, wherein the average time of resolution varies from 5-36

weeks<sup>11</sup>.

Miyata et al<sup>12</sup> described intracameral air injections for resolution of acute hydrops wherein resolution of took about 20.1±9.0 days with an average of 2.4±1.3 injections compared to those who received conventional treatment which took 64.7±34.6 days. Panda et al<sup>13</sup> reported the use of SF<sub>6</sub> instead of air in their series of 9 cases with the need for repeat injections in 6 cases. Sayan Basu et al<sup>10</sup> have used non-expansile C<sub>3</sub>F<sub>8</sub> for the same purpose in their retrospective study wherein resolution of hydrops took 78.7±53.2 days versus the conventional treatment that took 117.9±68.2 days.

In our case, we have used C<sub>3</sub>F<sub>8</sub> along with air in the ratio of 3:1, filling 2/3<sup>rd</sup> of the anterior chamber with an intention to reduce the concentration of the gas and also allow its expansion. C<sub>3</sub>F<sub>8</sub> has a distinct advantage over air and SF<sub>6</sub> as it is retained in anterior chamber for a longer time and

therefore, repeated injections may not be needed<sup>14</sup>. The gas acts as a mechanical barrier by preventing entry of aqueous into the stroma and at the same time, provides the tamponade necessary to repose the torn edges of Descemet's membrane, due to which there is a faster resolution of stromal oedema.

C<sub>3</sub>F<sub>8</sub> has also been used to treat acute hydrops following keratoglobus<sup>10</sup>, pellucid marginal degeneration<sup>15</sup> and Descemet membrane detachments following viscocanaliculostomy<sup>16,17</sup> with varying degrees of success. Endothelial toxicity of C<sub>3</sub>F<sub>8</sub> has been a concern, but its use is described as safe and effective in re-attachment of Descemet's membrane after a complicated cataract surgery<sup>18</sup>. The stromal cyst that has been described in our case was left alone due to the existence of a possible communication with the anterior chamber<sup>19</sup>. Therefore, Descemetopexy using C<sub>3</sub>F<sub>8</sub> in the management of acute corneal hydrops can be an effective and safe procedure for the early resolution.

#### References:

1. Krachmer, Mannis, Holland. Textbook of Cornea, 2005, 2<sup>nd</sup> Edition, Vol. 1 Pg. 955-956.
2. Krachmer JH, Feder RS, Belin MW. Keratoconus and related non-inflammatory corneal thinning disorders. *Surv Ophthalmol.* Jan-Feb 1984;28(4):293-322.
3. Rabinowitz YS. Keratoconus. *Surv Ophthalmol.* Jan-Feb 1998;42(4):297-319.
4. Haugen OH. Keratoconus in mentally retarded, *Acta Ophthalmol (Copenh).* 1992;70:111-114.
5. Karseras AG, Ruben M. Aetiology of Keratoconus. *Brit J Ophthalmol.* 1976;60:522-525.
6. Tuft SJ, Gregory WM, Buckley RJ. Acute Corneal Hydrops in Keratoconus. *Ophthalmology.* 1994;101:1738-44.
7. Gupta VP, Jain RK, Angra SK. Acute Hydrops in Keratoglobus with Vernal Keratoconjunctivitis. *Indian J Ophthalmol.* 1985;33:121-3.
8. Carter JB, Jones DB, Wilhelmus KR. Acute hydrops in Pellucid Marginal Degeneration. *Am J Ophthalmol.* 1989;107:167-70.
9. Wolter JR, Menderson JW, Clahessey EG. Ruptures of Descemet's Membrane in Keratoconus causing Acute Hydrops and Posterior Keratoconus. *Am J Ophthalmol.* 1967;63:1689-92.
10. Sagar Basu, Pravin K Vaddavalli, Muralidhar Ramappa et al. Intracameral Perfluoropropane gas in treatment of acute corneal hydrops. *Ophthalmology.* 2011;118:934-939.
11. Grewal S, Laibson PR, Cohen EJ, Rapnano CJ. Acute hydrops in corneal ectasias: Associated factors and outcomes. *Trans Am Ophthalmol Soc.* 1999;97:187-98.
12. Miyata K, Tsuji H, Tanabe T et al. Intracameral air injection for Acute Hydrops in Keratoconus. *Am J Ophthalmol.* 2002;133:750-2.
13. Panda A, Aggarwal A, Madhavi P et al. Management of Acute Corneal Hydrops secondary to Keratoconus with intracameral injection of SF<sub>6</sub>. *Cornea.* 2007;26:1067-9.
14. Shah SG, Sridhar MS, Sangwan VS. Acute Corneal Hydrops treated by intracameral C<sub>3</sub>F<sub>8</sub>. *Am J Ophthalmol.* 2005;139:368-70.
15. Kaushal S, Sharma N, Vajpayee RB. Treatment of Acute Corneal Hydrops with intracameral C<sub>3</sub>F<sub>8</sub> in a patient of Pellucid Marginal Degeneration with Keratoglobus. *Clin Experiment Ophthalmol.* 2007;35:697-9.
16. Kaan Unlu, Ahmet Aksunger. Descemet's Membrane detachment after Viscocanaliculostomy. *Am J Ophthalmol.* 2000;130:833-834.
17. Hiroko Fujimoto, Takanori Mizoguchi, Shinichiro Kuroda, Makoto Nagata. Intracorneal Hematoma with Descemet's Membrane detachment after Viscocanaliculostomy. *Am J Ophthalmol.* 2004;137:195-196.
18. Kim T, Hasan SA. A new technique for repairing Descemet's membrane detachments using intracameral gas injections. *Arch Ophthalmol.* 2002;120:181-3.
19. Curtis E Margo, Mathew W Mosteller. Corneal Pseudocyst following acute hydrops. *Brit J Ophthalmol.* 1987;71:359-360.