

Confronting COVID-19: Issues in Hemophilia and Congenital Bleeding Disorders

Antonio Coppola, MD¹ Annarita Tagliaferri, MD¹ Gianna Franca Rivolta, MD¹
Gabriele Quintavalle, MD¹ Massimo Franchini, MD²

¹Regional Reference Center for Inherited Bleeding Disorders
University, Hospital of Parma, Parma, Italy

²Department of Hematology and Transfusion Medicine, Carlo Poma
Hospital, Mantova, Italy

Address for correspondence Antonio Coppola, MD, Regional
Reference Center for Inherited Bleeding Disorders, University Hospital
of Parma, Via Gramsci, 14-43126 Parma, Italy
(e-mail: ancoppola@ao.pr.it).

Semin Thromb Hemost 2020;46:819-822.

The Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), emerging toward the end of 2019 in an epidemic focused in Wuhan, China, is responsible for Coronavirus Disease 2019 (COVID-19), which spread quickly throughout China and the neighboring Asian countries, and immediately after to virtually all countries around the world,¹ with ~ 3,000,000 confirmed infections and more than 200,000 deaths, so far.² Flu-like symptoms are the most common clinical presentation; however, 15 to 20% of patients develop a severe form of interstitial pneumonia and respiratory insufficiency, up to need for intensive care and mechanical ventilation, with significant fatal outcome, particularly in those with advanced age and comorbidities.¹

The unexpectedly large numbers of severely affected patients have overwhelmed health care systems, with most human and economic resources devoted to the critical epidemic emergency. Moreover, the need for mobility restrictions is limiting access to usual standard of medical care, which is likely harmful for patients with chronic diseases and/or requiring regular clinical follow-up. This is the case for congenital bleeding disorders (CBD), especially the most symptomatic forms, such as severe hemophilia. The challenges of health care management in the age of the COVID-19 pandemic, with focus on persons with hemophilia (PWH), have been recently highlighted³ and the World Federation of Hemophilia (WFH) issued specific recommendations for patients, including hygienic measures and information to enable reducing their exposure to SARS-CoV-2 and the associated risks of inadequate access to, or conduct of, treatment.⁴ Shortage of replacement product supply due to problems or restrictions in air transport can occur in low-income countries receiving aid from WFH and other humanitarian programs.³ In high-income countries, no relevant immediate impact in product delivery is expected. However, regular medical and laboratory follow-up is likely to

be hampered at specialized hemophilia treatment centers (HTC), otherwise useful to enable optimized outcomes in patients on personalized regimens of primary or secondary prophylaxis to minimize bleeding and its deleterious effects, particularly on the musculoskeletal system.^{5,6} Relevant advances in therapeutic products, that is, extended half-life (EHL) factor VIII and IX concentrates⁶ or the FVIII-mimetic bispecific monoclonal antibody emicizumab, recently introduced for prophylaxis in hemophilia A patients with and without inhibitors,⁷ may provide benefits in this emergency context, like higher bleeding protection and reduced need for treatment administration and supply.⁸ However, switching products is not generally advised,⁴ being practically impossible to carefully follow-up patients during treatment transitions.⁹ Moreover, some patients are on investigational treatments, including non-replacement agents and gene therapy, or are scheduled to be enrolled in clinical trials, the conduct of which may be interrupted or necessarily adapted to the evolving situations.³ Overall, HTCs are closely meeting the specific needs for diagnostic or therapeutic consultations and follow-up of PWH and CBD patients, also using telemedicine and social media communication, and relying on the long-established trustworthy relationships with patients, their families, and key associations.³

The hemophilia community is actively confronting challenges of management of CBD in the pandemic era, but issues related to the specific impact of COVID-19 in CBD patients are largely unknown. To date a single case report has been published describing a patient with severe hemophilia A in Wuhan, China, with mild symptoms due to confirmed SARS-CoV-2 pneumonia.¹⁰ National and international initiatives are ongoing to collect information about CBD patients with confirmed or clinically suspected COVID-19. Considering the numbers of COVID-19 infections reported in Italy¹¹ and the

published online
June 8, 2020

Issue Theme Maintaining Hemostasis
and Preventing Thrombosis in COVID-19
—Part I; Guest Editors: Emmanuel J.
Favaloro, PhD, FFS (RCPA), and Giuseppe
Lippi, MD.

Copyright © 2020 by Thieme Medical
Publishers, Inc., 333 Seventh Avenue,
New York, NY 10001, USA.
Tel: +1(212) 760-0888.

DOI <https://doi.org/10.1055/s-0040-1712961>.
ISSN 0094-6176.

incidence of CBD in our country,¹² at least 30 CBD patients should have been diagnosed with COVID-19, 6 with severe or critical symptoms. Currently, there is no reason to think that susceptibility to SARS-CoV-2 infection or clinical course should differ in CBD patients from the general population.¹³ Male and hypertensive patients more frequently experience severe up to fatal COVID-19.¹ These clinical features could negatively affect PWH, almost all men and, according to some studies, suffering from hypertension with higher prevalence than age-matched subjects.¹⁴ Comorbidities in PWH possibly associated with higher risk for severe COVID-19 are human immunodeficiency (HIV) and hepatitis B and C (HBV and HCV) virus infections, frequently found in those born before mid-1980s,¹² (i.e., before the introduction of virus-inactivated plasma-derived factor concentrates). However, most patients are not immunocompromised or show hepatic decompensation, thanks to currently available treatments.

An intriguing issue is represented by the possible influence of the coagulation defect in the clinical outcome of COVID-19 in CBD patients. Severe COVID-19 is associated with abnormalities of coagulation tests, reflecting a hypercoagulable state and hyperfibrinolysis, with mildly prolonged prothrombin time (PT) and reduced platelet count in most patients and, in particular, highly increased D-dimer, correlated with the severity of disease and risk of death.¹⁵⁻¹⁷ After pulmonary viral replication and localized inflammation, the acute lung injury and hypoxia trigger a systemic hyperinflammatory state, due to activation of endothelia and cells of the monocyte/macrophage lineage, thus causing the 'cytokine storm' of the second stage of disease.¹⁸ A vicious cycle of inflammation and coagulation activation is likely to be responsible for lung microvascular thrombosis and systemic venous thromboembolism (VTE), increasingly reported in COVID-19.¹⁹⁻²¹ Therefore, thromboprophylaxis with low-molecular weight or unfractionated heparin or fondaparinux at least at standard doses is advised in COVID-19 patients,²² but heparin intensified-dose

regimens up to therapeutic anticoagulation are being investigated in clinical trials or used according to pragmatic risk-adjusted protocols,²³ aiming at modulating thrombo-inflammatory mechanisms.

On this basis, could the underlying abnormality of coagulation mitigate the prothrombotic state and the clinical course of COVID-19 in CBD patients? This question correlates with the possible protection of CBD patients from cardiovascular and thromboembolic diseases,^{24,25} keenly debated in recent years, in the light of increased patients' life expectancy and the consequent higher burden of cardiovascular risk factors. Similarly, indications and conduct of antithrombotic treatments in this setting should consider the inherent bleeding risk and the need for concomitant replacement treatment, according to the severity of coagulation defect. No evidence-based guidelines are available; however, standard prophylactic doses of heparin are considered safe in PWH with (baseline or therapeutically achieved) factor levels > 5%. Higher factor levels are needed if increased doses are used, up to > 20 to 30% for full anticoagulation, as in the case of thromboembolic complications.²⁴⁻²⁶

Antithrombotic prophylaxis/treatment is challenging in the management of CBD patients with COVID-19, as well all conditions inducing increased bleeding risk, due to the disease or to its management and treatment.¹³ Therefore, adequate replacement treatment should be considered, aiming to achieve protective factor levels, according to the severity of risk and of the specific CBD (► **Table 1**). The only reported PWH with COVID-19 did not experience bleeding complications, related to the disease or to treatment, performed at home with antiviral (oseltamivir) and antibiotic (cefdinir) drugs.¹⁰ Bleeding symptoms are not frequent in COVID-19. Hemoptysis, due to mucosal congestion and lung injury, is reported in 0.9% of patients, but up to 3% in severe cases.¹ Nasal and throat bleeding, and rarely gastrointestinal hemorrhage also occur. Frequency and severity of symptoms are expected to be higher

Table 1 Bleeding risk in the clinical course and management of COVID-19 in patients with hemophilia and congenital bleeding disorders

Condition	Setting	Low-moderate risk	High risk
Disease-related	Bleeding symptoms - Tissue injury and congestion	Nasal and throat bleeding	Hemoptysis GI bleeding
	- Trauma and predisposing conditions	Subcutaneous and muscle hematoma	Intracranial bleeding
Management and treatment-related	Invasive procedures	Arterial puncture for blood gas analysis	ECMO
		PICC or CICC insertion	
		Invasive ventilation (intubation, tracheostomy)	
		Renal replacement (dialysis and ultrafiltration)	
	Pharmacological treatment	Antiviral drugs	Corticosteroids
		Immunomodulatory agents	NSAIDs
Antithrombotic agents at prophylactic or weight-adjusted doses		Antithrombotic agents at therapeutic doses	

Abbreviations: CICC, centrally inserted central catheters; ECMO, extracorporeal membrane oxygenation; GI, gastrointestinal; NSAID, non-steroidal anti-inflammatory drugs; PICC, peripherally inserted central catheters.

in CBD patients, who may experience bleeding even due to trauma or other predisposing conditions (for example, prolonged axillary and neck compression by continuous positive airway pressure helmet and straps, or persistent cough attacks or nose obstruction and vigorous blowing, causing increase of blood or abdominal pressure).⁴ Hospitalized patients are advised to carefully inform medical COVID-19 team about their CBD and replacement treatment and to provide contacts of the reference HTC. Prophylaxis regimens should be continued or implemented in all patients with severe or moderately severe CBD, aiming at maintaining factor levels higher than those in the routine practice.⁴ This will help minimize bleeding risk, even due to the frequent arterial punctures for blood gas monitoring and other invasive procedures, including those needed for vascular access, mechanical ventilation or renal replacement (► **Table 1**). For patients requiring extracorporeal membrane oxygenation, at highest bleeding risk, achieving virtually normal factor levels has been suggested.

Bleeding threats are also related to pharmacological treatments (► **Table 1**). Non-steroidal anti-inflammatory drugs (NSAIDs) should be avoided, because of inhibition of platelet function with increased bleeding tendency, in particular upper gastrointestinal hemorrhage.²⁷ Anti-platelet effects are negligible with selective cyclo-oxygenase 2 inhibitors and corticosteroids. The latter, frequently used in the management of the second phase of COVID-19,²⁸ mainly in severe cases, are known to be associated with significant risk of gastrointestinal bleeding, in particular in hospitalized patients.²⁹ These risks, in addition to that of stress ulcer in this setting of critically-ill patients, highlight the role of prophylaxis with proton pump inhibitors or histamine-2 receptor antagonists.³⁰

Although a specific treatment for COVID-19 is currently unavailable, an increasing series of drugs, previously used for treating other viral diseases or investigational agents with antiviral targets or immunomodulatory effects, are being evaluated in clinical practice.²⁸ Many of these drugs may increase bleeding tendency because of toxicity at gastrointestinal (i.e., lopinavir/ritonavir, umifenovir) or hematologic (i.e., the interleukin-6 receptor inhibitor tocilizumab and the RNA-polymerase inhibitor favipiravir) level. In this respect, reduction in platelet count should be carefully considered, because of bleeding risk, but also due to the debated association with mortality in COVID-19 patients.³¹

As mentioned earlier, coagulation laboratory testing provides crucial prognostic markers, that is, PT and, particularly, D-dimer.¹⁵⁻¹⁷ Their abnormalities reflect the hypercoagulable state in COVID-19 and require monitoring, as some of most severe patients may develop sepsis and overt disseminated intravascular coagulopathy (DIC).¹ If coagulation laboratory abnormalities in COVID-19 are not associated with bleeding tendency, in CBD patients additional issues should be considered. The massive inflammatory response leads to increase of reactive proteins, including factor VIII and von Willebrand factor.³² This may mask mild cases of hemophilia A and von Willebrand disease. Other CBDs with normal testing include FXIII deficiency and platelet functional defects; thus, undiagnosed patients could be missed, partic-

ularly in cases on non-referral to HTC. Moreover, in patients with hemophilia A currently on prophylaxis with emicizumab, activated partial thromboplastin time (APTT) and all clotting tests based on APTT are not reliable, overestimating coagulation and thus masking the underlying severe CBD.^{3,4} Due to the prolonged half-life of emicizumab, these effects may persist for months after drug discontinuation. Patients on this and other investigational non-replacement agents (i.e., fitusiran and anti-TFPI) should be carefully monitored during COVID-19, because the risk of thrombotic complications is currently unknown.¹⁶ However, treatment should not be discontinued and caution is needed if additional hemostatic therapy is administered, in particular activated prothrombin complex concentrate (APCC) in inhibitor patients.³³

In conclusion, HTCs and the hemophilia community are defining safe modalities for facing the COVID-19 pandemic. Clinical data in CBD patients with COVID-19 are lacking; however, they should undergo all treatment modalities available, including invasive approaches, provided that their bleeding risk is covered by adequate hemostatic prophylaxis. As our knowledge and clinical experience develop, we will learn whether the coagulation defects present in PWH and CBD may influence the epidemiologic impact and severity of COVID-19.

Conflict of Interest

Dr. Tagliaferri reports personal fees from Bayer, Novo Nordisk, and Roche, outside the submitted work. Dr. Coppola reports personal fees from Bayer, Novo Nordisk, and Werfen, outside the submitted work. Dr. Quintavalle reports personal fees from Bayer, outside the submitted work.

References

- 1 Guan WJ, Ni ZY, Hu Y, et al; China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382(18):1708-1720
- 2 World Health Organization. Coronavirus disease (COVID-19) outbreak. Available at <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>. Accessed April 28, 2020
- 3 Hermans C, Weill A, Pierce GF. The COVID-19 pandemic: New global challenges for the haemophilia community. *Haemophilia* 2020. [Epub ahead of print]. Doi: 10.1111/hae.14001
- 4 World Federation of Haemophilia. COVID-19: Practical recommendations for people with hemophilia. Available at <https://news.wfh.org/covid-19-coronavirus-disease-2019-pandemic-caused-by-sars-cov-2-practical-recommendations-for-hemophilia-patients/>. Accessed April 28, 2020
- 5 Coppola A, Tagliaferri A, Di Capua M, Franchini M. Prophylaxis in children with hemophilia: evidence-based achievements, old and new challenges. *Semin Thromb Hemost* 2012;38(01):79-94
- 6 Berntorp E, Andersson NG. Prophylaxis for hemophilia in the era of extended half-life factor VIII/factor IX products. *Semin Thromb Hemost* 2016;42(05):518-525
- 7 Franchini M, Marano G, Pati I, et al. Emicizumab for the treatment of haemophilia A: a narrative review. *Blood Transfus* 2019;17(03):223-228
- 8 Hermans C, Lambert C. Impact of the COVID-19 pandemic on therapeutic choices in thrombosis-hemostasis. *J Thromb Haemost* 2020. [Epub ahead of print]. Doi: 10.1111/jth.14845

- 9 Tagliaferri A, Matichecchia A, Rivolta GF, et al. Optimising prophylaxis outcomes and costs in haemophilia patients switching to recombinant FVIII-Fc: a single-centre real-world experience. *Blood Transfus* 2019. [Epub ahead of print] . Doi: 10.2450/2019.0220-19
- 10 Cui D, Zhang A, Liu A, Hu Q. Clinical findings in a patient with haemophilia A affected by COVID-19. *Haemophilia* 2020. [Epub ahead of print] . Doi: 10.1111/hae.14000
- 11 National Institute of Health. COVID-19 integrated surveillance in Italy. Key national data. April 27, 2020 update. Available at <https://www.epicentro.iss.it/en/coronavirus/sars-cov-2-integrated-surveillance-data>. Accessed April 28, 2020
- 12 Abbonizio F, Hassan HJ, Riccioni R, Santagostino E, Arcieri R, Giampaolo A; Italian Association of Haemophilia Centres (see Appendix). New data from the Italian National Register of Congenital Coagulopathies, 2016 Annual Survey. *Blood Transfus* 2020;18(01):58–66
- 13 World Federation of Haemophilia. Specific risks of COVID-19 to the bleeding disorders community. Available at <https://news.wfh.org/specific-risks-of-covid-19-to-the-bleeding-disorders-community/>. Accessed April 28, 2020
- 14 Fransen van de Putte DE, Fischer K, Makris M, et al. Increased prevalence of hypertension in haemophilia patients. *Thromb Haemost* 2012;108(04):750–755
- 15 Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost* 2020;18(04):844–847
- 16 Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020;18(05):1023–1026
- 17 Favaloro EJ, Lippi G. Recommendations for minimal laboratory testing panels in patients with COVID-19: Potential for prognostic monitoring. *Semin Thromb Hemost* 2020;46(03):379–382
- 18 Siddiqi HK, Mehra MR. COVID-19 illness in native and immunosuppressed states: a clinical-therapeutic staging proposal. *J Heart Lung Transpl* 2020. [Epub ahead of print] . Doi: 10.1016/j.healun.2020.03.012
- 19 Cui S, Chen S, Li X, Liu S, Wang F. Prevalence of venous thromboembolism in patients with severe novel coronavirus pneumonia. *J Thromb Haemost* 2020. [Epub ahead of print] . Doi: 10.1111/jth.14830
- 20 Klok FA, Kruip MJHA, van der Meer NJM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;S0049-3848(20)30120-1; [Epub ahead of print] . Doi: 10.1016/j.thromres.2020.04.013
- 21 Dolhnikoff M, Duarte-Neto AN, de Almeida Monteiro RA, et al. Pathological evidence of pulmonary thrombotic phenomena in severe COVID-19. *J Thromb Haemost* 2020. [Epub ahead of print] . Doi: 10.1111/jth.14844
- 22 Bikdeli B, Madhavan MV, Jimenez D, et al. COVID-19 and thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy, and follow-up. *J Am Coll Cardiol* 2020; S0735-1097(20)35008-7; [Epub ahead of print] . Doi: 10.1016/j.jacc.2020.04.031
- 23 Coppola A, Lombardi M, Tassoni MI, et al. COVID-19, thrombotic risk and thromboprophylaxis: learning lessons from the bedside, awaiting evidence. *Blood Transfus* 2020. [epub ahead of print] Doi: 10.2450/2020.0113-20
- 24 Coppola A, Tagliaferri A, Franchini M. The management of cardiovascular diseases in patients with hemophilia. *Semin Thromb Hemost* 2010;36(01):91–102
- 25 Franchini M, Coppola A. Atherothrombosis in von Willebrand disease: an analysis of the literature and implications for clinical management. *Semin Thromb Hemost* 2012;38(02):185–199
- 26 Schutgens RE, van der Heijden JF, Mauser-Bunschoten EP, Mannucci PM. New concepts for anticoagulant therapy in persons with hemophilia. *Blood* 2016;128(20):2471–2474
- 27 Arachchilage DRJ, Makris M. Choosing and using non-steroidal anti-inflammatory drugs in haemophilia. *Haemophilia* 2016;22(02):179–187
- 28 Sanders JM, Monogue ML, Jodlowski TZ, Cutrell JB. Pharmacologic Treatments for Coronavirus Disease 2019 (COVID-19): A Review. *JAMA* 2020. [Epub ahead of print] . Doi: 10.1001/jama.2020.6019
- 29 Narum S, Westergren T, Klemp M. Corticosteroids and risk of gastrointestinal bleeding: a systematic review and meta-analysis. *BMJ Open* 2014;4(05):e004587
- 30 Wang Y, Ye Z, Ge L, et al. Efficacy and safety of gastrointestinal bleeding prophylaxis in critically ill patients: systematic review and network meta-analysis. *BMJ* 2020;368:l6744
- 31 Yang X, Yang Q, Wang Y, et al. Thrombocytopenia and its association with mortality in patients with COVID-19. *J Thromb Haemost* 2020. [Epub ahead of print] . Doi: 10.1111/jth.14848
- 32 Terraube V, O'Donnell JS, Jenkins PV. Factor VIII and von Willebrand factor interaction: biological, clinical and therapeutic importance. *Haemophilia* 2010;16(01):3–13
- 33 Castaman G, Santoro C, Coppola A, et al; ad hoc Working Group. Emergency management in patients with haemophilia A and inhibitors on prophylaxis with emicizumab: AICE practical guidance in collaboration with SIBioC, SIMEU, SIMEUP, SIPMeL and SISET. *Blood Transfus* 2020;18(02):143–151