

A Hospital-Based Insight into the Antinuclear Antibody Patterns in Autoimmune Disorders

Prishni Gupta¹ Ritu Priya¹ Rachita Nanda^{1,✉} Suprava Patel¹ Eli Mohapatra¹

¹Department of Biochemistry, All India Institute of Medical Sciences, Raipur, Chhattisgarh, India

Address for correspondence Rachita Nanda, MBBS, MD, Department of Biochemistry, Medical College, AIIMS Raipur, Chhattisgarh, 492099, India (e-mail: dr.rachitananda@gmail.com).

J Lab Physicians:2020;12:115–120

Abstract

Background The incidence of autoimmune disorders has seen a rise in India in recent times. The symptoms and signs of these conditions are caused by a systemic autoimmune response, essentially characterized by the expression of the antinuclear antibodies (ANA). Presence of ANA in serum of patient could be the prime evidence of an autoimmune disorder.

Aim This study aimed to determine the antibody patterns and assess the clinical significance of ANA in patients of a teaching tertiary care hospital of central India.

Material and Methods This retrospective cross-sectional data analysis study retrieved 538 reports of individuals, who were prescribed the ANA test by indirect immunofluorescence assay over a period of 11 months, from the archives of the Department of Biochemistry. For continuous data, student *t*-test was used while Chi-square and Fisher exact was conducted for categorical data. A *p* value less than 0.05 were taken as significant.

Results Out of the 538 patients investigated for ANA testing by indirect immunofluorescence assay, 33% were positive, among which 74% were female, and majority belonged to the young adult age group. The most common pattern identified was nuclear, subpattern nuclear speckled.

Conclusion A high-ANA pattern positivity could be related to a high-autoimmunity prevalence in this region, and also promotes its use as a tool of evidence of suspected autoimmune disorders.

Keywords

- ▶ central India
- ▶ immunofluorescence assay
- ▶ nuclear

Introduction

In general, when a patient presents with clinical manifestations associated with autoimmune condition, the first test advised is antinuclear antibody (ANA)¹ that serve as the serological hallmarks of autoimmunity.² The ANA attack self-proteins within cell nucleus structures, now encompassing nuclear envelope components, mitotic spindle apparatus, cytosol, cytoplasmic organelles, and cell membranes.³ Their presence in serum may indicate either a systemic autoimmune disease like systemic lupus erythematosus (SLE), scleroderma and polymyositis/dermatomyositis, or an organ-specific condition like autoimmune thyroiditis and hepatitis.² Identification of ANA patterns and their target antigen on the basis of variable cellular

staining pattern can be correlated to specific autoimmune disease as part of diagnostic clinical immunology.^{2,4} Although a collection of tests are available for ANA detection, the indirect immunofluorescence (IIF)-ANA on HEp-2 cells remains the gold standard.⁵

The autoimmune disorders are responsible for considerable morbidity and mortality in Western countries, with prevalence of 3 to 5%. However, the condition in Asian countries like India is still blurred because only a few studies on individual diseases have been undertaken.⁶ Thus, the present study aims to determine the ANA positivity, study the ANA patterns, and assess their clinical significance in patients afflicted with autoimmune disorders from central India.

Materials and Methods

In this retrospective cross-sectional data analytic study, we retrieved and reviewed 538 ANA test reports, between January to November, 2018, from the archives of the Department of Biochemistry of a tertiary care center in Central India, irrespective of age and specialty departments.

IIF assay has been positioned as the gold standard for ANA testing by the American College of Rheumatology, with the advent of utilizing a cell-line substrate from human laryngeal carcinoma cells and Hep-2 cells, combined with a primate liver. In our center, ANA in serum was detected using the EUROIMMUN titer-plate technique. The kit comprised of Biochip slides embedded with HEp-20-10/liver (monkey) tissue sections. As per the manufacturer protocol, serum samples were diluted 1:100 with phosphate buffer saline (PBS)-Tween followed by tagging with fluorescein-labeled antihuman globulin. ANA binding to specific antigen leads to a fluorescent emission that was visualized under an inverted fluorescence microscope under 10× and 40× objectives. The fluorescent intensity of each sample was compared with both negative and positive control that were supplied in the kit. The grouping and classification of ANA patterns were as per the ICAP (International Congress on Antinuclear Antibody Patterns) nomenclature. The records that were analyzed were categorized based on demographic factors and clinical information available in the patient requisition forms.

Data Analysis

The data collected was analyzed in IBM SPSS 21. For continuous data, student *t*-test was used, while Chi-square and Fisher exact was done for categorical data. A *p* value less than 0.05 were taken as significant.

Results

Out of the 538 ANA reports that were reviewed, 33% (179) were positive for ANA. Although, the patients belonged from different disciplines of the hospital, general medicine that includes endocrinology contributed to maximum patient load (► **Table 1**).

ANA positivity was maximum in the 30 to 39 age group patient population, mean age being 37 ± 18 years ($p < 0.001$) (► **Table 2**). Females were more affected than males in ratio 1:2.8 ($p = 0.03$). All the age groups had females more positive than males, although the difference was not statistically significant (► **Table 3**).

The different patterns that were found in this study are depicted in ► **Table 4**. The nuclear pattern was the most common (68.7%), and amongst its subtypes, nuclear speckled pattern predominated the results (24%), followed by nuclear homogenous (21.2%); similarly, 18.9% patients had the cytoplasmic pattern, with 8.4% showing the cytoplasmic speckled pattern.

The patient population belonging to the 30 to 39 age group showed positivity for every pattern except nuclear envelope and mitotic intercellular bridge. The nuclear speckled pattern was seen in all age groups and so did the cytoplasmic speckled pattern but in fewer numbers, although with no patients in the 0 to 9 years age group. All the other subpatterns showed sporadic appearances in the different age groups (► **Supplementary Table S1**, online only).

Further, 162 patients were segregated based on available clinical data, symptoms and diagnosis. ► **Table 5** shows the disease spectra versus age group distribution numbers. Most patients were in the 30- to 39-years age group and maximum of them had an underlying skin disorder (acne vulgaris, pemphigus vulgaris, bulous pemphigoid), followed by patients with infections and inflammatory conditions. The next in majority were patients with various rheumatoid diseases with

Table 1 Distribution of total and positive cases across clinical disciplines

Departments	OPD	IPD	Total (n)	Positive (n)	Positivity %
General medicine/endo	175	60	235	74	41.34
Trauma/casualty	7	39	46	18	10.06
Pediatrics	35	17	52	15	8.38
Orthopaedics	68	3	71	24	13.41
Dermatology	43	9	52	21	11.73
Cardiology	8	0	8	3	1.68
Neurology	18	1	19	3	1.68
Pulmonary medicine	21	2	23	11	6.15
ENT	8	1	9	4	2.23
Ophthalmology	11	1	12	3	1.68
OBG	5	3	8	1	0.56
Psychiatry	1	1	2	1	0.56
Surgery	0	1	1	1	0.56
Total	400	138	538	179 (33%)	100

two age peaks, adolescent and young adult. A similar peaking was seen with the hematological conditions. While infections were more common in the adolescents, skin conditions were mostly seen in the early adult age group. Lupus had a homogenous distribution in all age groups till 70 years.

ANA positivity was observed in 54.5% patients afflicted with SLE or discoid lupus erythematosus (DLE), with a statistically significant bias toward nuclear homogenous pattern. A similar statistically significant distribution was seen in the rheumatoid disease group with a 38.9% positivity rate, and the skin conditions with a 47.8% positivity rate. The study did not find a positive ANA result in nephrotic syndrome, endocrine disorders, and Hansen's disease. The nuclear speckled pattern was seen in all conditions except in SLE and DLE, but the highest percentage was observed in dermatological disorders. While in all the conditions, ANA positivity was high, it was markedly high in chronic liver diseases (CLD) and interstitial lung diseases (ILD), which also showed positivity for certain uncommon patterns (→ **Supplementary Table S2**, online only).

Discussion

Autoimmune serology is an important diagnostic tool in a clinical setup in which the ANA is certainly one of the most frequently requested tests in India.⁷ The introduction of human cancer cell lines, Hep-2, as a substrate for the ANA test significantly increased the sensitivity, which however

often leads to high-false positive rates, making the interpretation for true positive difficult. This was solved by using a biochip that incorporated primate liver with the Hep-2 cells. The high-sensitivity of Hep-2 cells now needed to be confirmed by the response of the primate liver, which brought the rates of false positives down tremendously.⁸

In a country like India, a laboratory needs adequate, reliable and cheap screening tests. ANA-IIF is not only easy to perform but also cheap. Despite its subjectivity, some of the ANA patterns are specific and are extremely helpful in clinical diagnosis.⁹ With the rise in prevalence of autoimmune disorders in India and scarce documentation of prevalence, any information about their occurrence and clinical significance could be beneficial.

The present study was a patient-based study, and the first study of its kind in central India. ANA test reports of 538 patients were revisited, evaluating the ANA positivity status with the help of IIF on Hep-2 cell lines and primate liver. An ANA positivity rate of 33% was found. Only two other studies have been undertaken in India, in Bangalore by Sebastian et al⁸ and in Chandigarh by Minz et al.⁶ Although the frequency of positivity in Bangalore was similar to the present study, Chandigarh showed a lesser positivity (→ **Table 6**). Similar hospital-based studies done in Turkey by Zafer et al⁹ and Sener et al¹⁰ and in Belgium by Peene et al⁷ had lesser positivity rate compared with present study. Although the results of these studies cannot be considered as prevalence, yet an estimation of disease burden can be made. However, the population-based studies showed mixed results wherein Akmatov et al in Germany⁴ and Chanwit et al in Thailand¹¹ found ANA positivity rate to be 33% and 39.6%, respectively, while Guo et al¹² in China and Satoh et al² in the USA demonstrated much lower prevalence. The results contradict the fact that ANA and, in turn, autoimmunity is seen more in Western countries than in India, which could have been a result of previous poor screening facilities. Strikingly, a regional difference is found in India itself, which may be due to climatic variation, genetic variation, and controlling the different molecular mechanisms of autoimmunity, as explained by Atassi et al.¹³ This leaves scope and requirement for further analysis of autoimmunity in the country in different cultures and sects.

Table 2 Demographic variables distribution of ANA positivity

Demographic variables results	Age (years)	Gender (M/F)
Most positive	30–39	Female
Mean (SD)	37.47(17.78)	73.6%
Statistical test applied	t-test–Levene's test of equality	Chi-square–Fisher's exact
Value	49.852	3.777
p-Value	< 0.001	0.03

Abbreviation: ANA, antinuclear antibody; SD, standard deviation.

Table 3 Age and gender distribution and association in ANA positivity

Age group (years)	Male (P/T)	Female (P/T)	Exact sig. (2-sided)	Exact sig. (1-sided)
0–9	2/17	4/12	0.354	0.194
10–19	5/23	15/39	0.258	0.149
20–29	11/31	25/70	0.825	0.502
30–39	8/32	35/91	0.143	0.096
40–49	5/23	33/81	0.142	0.094
50–59	4/23	9/48	1.000	0.836
60–69	4/10	7/16	1.000	0.588
70–79	6/10	3/8	0.637	0.319
> 80	2/3	1/1	1.000	0.750
Total	47/172	132/366	0.062	0.033

Abbreviation: ANA, antinuclear antibody.

Table 4 Prevalence of different ANA patterns

Broad pattern	(n)	(%)	Subtypes of pattern	(n)	(%)
Nuclear	123	68.7	<i>Nuclear homogenous</i>	38	21.2
			<i>Nuclear speckled</i>	43	24.0
			<i>Nuclear dotted</i>	7	3.9
			<i>Nuclear DFS</i>	12	6.7
			<i>Nuclear nucleolar</i>	16	8.9
			<i>Nuclear coarse speckled</i>	4	2.2
			<i>Nucleus nuclear envelope</i>	2	1.1
			<i>Nuclear centromere</i>	1	0.6
Cytoplasmic	34	18.9	<i>Cytoplasmic AMA</i>	8	4.5
			<i>Cytoplasmic speckled</i>	15	8.4
			<i>Cytoplasmic fibrillar</i>	3	1.7
			<i>Cytoplasmic fibrillar filamentous</i>	5	2.8
			<i>Cytoplasmic golgi</i>	2	1.1
			<i>Cytoplasmic rods and rings</i>	1	0.6
Mitotic	15	8.4	<i>Mitotic spindle</i>	9	5.0
			<i>Mitotic intercellular bridge</i>	3	1.7
			<i>Mitotic centrosome</i>	1	0.6
			<i>Mitotic chromosomal</i>	1	0.6
			<i>Mitotic NuMa</i>	1	0.6
Mixed	7	3.9		7	3.9
Total	179			179	100

Abbreviations: AMA, antimitochondrial antibodies; ANA, antinuclear antibodies; DFS, dense fine speckled.

Table 5 Disease spectrum in different age groups

Diagnosis	Age groups in years									Total
	0–9	10–19	20–29	30–39	40–49	50–59	60–69	70–79	>80	
DLE/SLE	2	3	1	1	2	1	1	–	–	11
Rheumatic disorders	1	5	1	5	3	2	–	–	–	17
Joint pain/swelling	1	–	3	3	3	–	1	–	–	11
Skin disorders	–	2	6	7	4	4	–	–	–	23
Sclerosis/scleroderma	–	1	1	1	2	–	–	–	–	5
Hematological conditions	1	5	2	6	3	1	–	–	–	18
Infections/inflammations	4	6	3	8	1	–	–	–	–	22
CLD	–	1	1	1	4	2	1	–	–	10
ILD	–	1	1	–	2	1	–	1	1	7
Cardiac conditions	–	2	1	–	1	2	–	–	–	6
Nephrotic syndromes	2	2	–	0	1	–	–	–	–	5
Neuro and musculoskeletal conditions	–	1	5	3	3	1	2	–	–	15
Hansen's disease	–	–	–	–	–	1	–	–	–	1
Endocrine conditions	–	–	–	3	–	–	–	–	–	3
Miscellaneous	–	–	4	2	1	–	–	1	–	8
Total	11	29	29	40	30	15	5	2	1	162

Abbreviations: CLD, cystic lung disease; DLE, discoid lupus erythematosus; ILD, interstitial lung disease; SLE, systemic lupus erythematosus.

Table 6 Comparing studies done in India and worldwide

Author	Place of study	Study design	Sample size	ANA prevalence	Age	Male vs female	Most common ANA pattern
Present study	Raipur, India	Patient-based	536	33%	37 ± 18	Female 73.6%	Nuclear speckled
Akmatov et al ⁴	Germany	Population-based	1199	33%	49(38–60)	Female 56.2%	Nuclear Speckled
Minz et al ⁵	Chandigarh, India	Patient-based	650	18.9%	42 (20–50)	Female 75.9%	Nuclear speckled
Sener et al ¹⁰	Turkey	Patient-based	19996	21.9%	–	–	Nuclear homogenous
Sebastian et al ⁸	Bangalore, India	Patient-based	5066	38.2%	–	–	Nuclear homogenous
Zafer et al ⁹	Bolo, Turkey	Patient-based	3127	15.8%	43 ± 18	Female 74.7%	Nuclear speckled
Chanwit et al ¹¹	Thailand	Healthy controls	58	39.6%	Older age group	Female	Nuclear speckled
Guo et al ¹²	China	Population-based	20970	5.9%	32	Female 72.6%	–
Peene et al ⁷	Belgium	Patient-based	10550	23.5%	–	–	Nuclear speckled
Satoh et al ²	USA	Population-based	4754	13.8%	50–59	Female 9.6% vs. 17.8%	Nuclear

Abbreviations: ANA, antinuclear antibody.

Female predominance in the field of autoimmunity is already established, although the reasons remain obscure. ► **Table 6** clearly shows that all the studies including the present found ANA positivity rate higher in the female patient or population group. This dominance could be attributed to estrogen as a potential modifier of autoimmunity, as stated by Parks et al.¹⁴ Hayter et al included microchimerism-induced autoimmunity and action of X-chromosome encoded genes in the reason too.¹⁵ The female to male ratio, although biased toward the former, varies in proportion in the age groups. The difference peaks in the young adulthood and middle age, tailing at the other ends (► **Table 3**). This variation was also pointed out in studies by Guo et al¹² and Chanwit et al.¹¹ This result complemented the result of Parks et al, stating that childbearing may play an important role in initial antigen stimulation or breaking tolerance to self-antigens contributing to the development of ANA.¹⁴

According to Beeson, as stated in his compilation,¹⁶ the events of the autoimmune disorders are high in the 20 to 50 age group, while Hayter et al pin pointed the 20 to 29 age group with the highest prevalence of autoimmune disorders.¹⁵ This corresponds to the higher prevalence of autoimmune disorders (► **Table 5**) and of ANA positivity (► **Supplementary Table S1**, online only) in this group, as seen in present study, with the mean age being 37 ± 18 years. This result was similar to that of Guo et al in China (mean age 32 years).¹² A higher mean age was reported by Akmatov et al, Minz et al and Zafer et al (49 years, 42 years and 43 years, respectively),^{4,6,9} while Satoh et al and Chanwit et al found the highest prevalence in the older age group.^{2,11} This difference could be accounted to study population composition. The ANA in older people could suggest a senile condition rather than a pathological problem. Despite this age

distribution, certain conditions occur more in certain age groups like infections and rheumatic disorders in teenagers, neurological manifestations in old age while ILD in low rate appear equally in all age groups.

The nuclear pattern as a whole is unquestionably the most common in all the studies worldwide. Among this, the nuclear homogenous and nuclear speckled patterns come to the forefront. The nuclear speckled pattern is almost evenly distributed in all age groups except the pediatric group which also does not show nuclear homogenous pattern positivity just like Chanwit et al in Thailand.¹¹ The result of present study was similar to study done by Minz et al in Chandigarh and Zafer et al in Turkey, showing high-ANA positivity in SLE and DLE, scleroderma and rheumatoid patients.^{6,9} However, the most common pattern associated with SLE was nuclear homogenous in contrast to nuclear speckled in their studies. Minz et al in their article on review of diagnostics in autoimmunity stated that SLE showed positivity for both nuclear homogenous and nuclear speckled.⁵ This was in agreement to Hayter et al who pointed to the possibility of multiple tissue and molecular targets in SLE.¹⁵ It is often thought that the uncommon patterns have no role to play in diagnosis. However, the present study and studies done by Chanwit et al and Sener et al found the rare ANA patterns in significant number, and also disease conditions associated with them, especially chronic hepatic conditions and carcinoma.^{10,11} This supported the extensive study done by Pieter et al on rare ANA patterns, and his conclusion of nuclear envelope and nuclear dot patterns was associated with autoimmune hepatitis and hepatic and colon carcinoma.¹⁷ The cytoplasmic patterns are also considered nondisease-specific but their significance should not be ignored, as stated by Chanwit et al, since they might as well point to certain undetected conditions.¹¹

This study has certain limitations in terms of adequate clinical data available, lack of reference in neighborhood, and the study design being retrospective. However, its strength lies in the fact that it is the first of its kind in central India, providing an overview of ANA and autoimmunity status in this region..

Conclusion

Young age with dermatological manifestation in SLE was the commonest presentation. Although nuclear speckled was the most frequent pattern, few unusual patterns were also observed. A prospective study is needed to understand not only the epidemiology of autoimmune disorders but also the predictive value of ANA IIF in clinical setup, besides considering the possibility of taking up IIF as a screening tool for autoimmune disorders.

Ethical Approval

Approval taken from Institute Ethics Committee, AIIMS Raipur (745/IEC-AIIMSRPR/2019).

Guarantor

RN.

Contributorship

PG was involved in gaining ethical approval, performed data analysis, and wrote the first draft of the manuscript; RP conducted research work on literature, protocol development, and sample analysis; RN conceived the study, performed data analysis, and reviewed the manuscript; SP and EM reviewed and edited the manuscript. All authors approved the final version of the manuscript.

Conflict of Interest

None declared.

References

- Elena M, Hernández-becerril N, Perez-chiney AC, et al. Results in Immunology Predictive value of antinuclear antibodies in autoimmune diseases classified by clinical criteria: analytical study in a specialized health institute, one year follow-up. *Results Immunol* 2015;5:13–22
- Satoh M, Chan EKL, Ho LA, Rose M, Parks CG, Cohn RD, et al. Study design. *2013*;64(7):2319–27
- Gautam K. Anti-Nuclear antibodies: current concepts and future direction for diagnosing connective tissue disease. *Journal of Pathology of Nepal* 2015;5:766–773
- Akmatov MK, Röber N, Ahrens W, et al. Anti-nuclear autoantibodies in the general German population: prevalence and lack of association with selected cardiovascular and metabolic disorders-findings of a multicenter population-based study. *Arthritis Res Ther* 2017;19(1):127
- Kumar Y, Bhatia A, Minz RW. Antinuclear antibodies and their detection methods in diagnosis of connective tissue diseases: a journey revisited. *Diagn Pathol* 2009;4:1
- Minz RW, Kumar Y, Anand S, et al. Antinuclear antibody positive autoimmune disorders in North India: an appraisal. *Rheumatol Int* 2012;32(9):2883–2888
- Peene I, Meheus L, Veys EM, De Keyser F. Detection and identification of antinuclear antibodies (ANA) in a large and consecutive cohort of serum samples referred for ANA testing. *Ann Rheum Dis* 2001;60(12):1131–1136
- Sebastian W, Roy A, Kini U, Mullick S. Correlation of antinuclear antibody immunofluorescence patterns with immune profile using line immunoassay in the Indian scenario. *Indian J Pathol Microbiol* 2010;53(3):427–432
- Mengeloglu Z, Tas T, Kocoglu E, Aktas G, Karabörk S. Determination of anti-nuclear antibody pattern distribution and clinical relationship. *Pak J Med Sci* 2014;30(2):380–383
- Ag S, Afsar I. Evaluation of antinuclear antibodies by indirect immunofluorescence and line immunoassay methods 0: four years 0 data from Turkey. *APMIS* 2014;122(12):1167–1170
- Prapinjumrune C, Prucktrakul C, Sooktonglarng T, Thongprasom K. Serum antinuclear antibody in adult Thais. *Gerodontology* 2017;34(1):86–89
- Guo YP, Wang CG, Liu X, et al. The prevalence of antinuclear antibodies in the general population of china: a cross-sectional study. *Curr Ther Res Clin Exp* 2014;76:116–119
- Atassi MZ, Casali P. Molecular mechanisms of autoimmunity. *Autoimmunity* 2008;41(2):123–132
- Birnbaum LS, Jusko TA, Kissling GE, et al. HHS public access. *J Immigr Minor Health* 2015; (919):1–21
- Hayter SM, Cook MC. Autoimmunity reviews updated assessment of the prevalence, spectrum and case definition of autoimmune disease. *Autoimmun Rev* 2012;11(10):754–765
- Beeson PB. Age and sex associations of 40 autoimmune diseases. *Am J Med* 1994;96(5):457–462
- Vermeersch P, Bossuyt X. Autoimmunity reviews prevalence and clinical significance of rare antinuclear antibody patterns. *Autoimmun Rev* 2013;12(10):998–1003