


W. ZHANG, Z. ZHANG, J.-C. TANG, J.-T. CHE, H.-Y. ZHANG, J.-H. CHEN\*, Z. YANG\*  
 (PEKING UNIVERSITY AND PEKING UNIVERSITY SHENZHEN GRADUATE SCHOOL, P. R. OF CHINA)

Total Synthesis of (+)-Haperforin G

*J. Am. Chem. Soc.* **2020**, *142*, 19487–19492, DOI: 10.1021/jacs.0c10122.

## Synthesis of (+)-Haperforin G



**Significance:** Chen, Yang, and co-workers report the first total synthesis of (+)-haperforin G, a structurally complex member of the limonoid family of tetranortriterpenoid natural products. Their synthesis features an efficient radical coupling of two fragments.

**Comment:** Intermediate **G** is accessed from **A** in nine steps and transformed to **H** via a Pauson–Khand reaction. Further six steps give rise to **I**, which is then joined with **M** via a light-mediated radical-based coupling to obtain **O**.

**SYNFACTS Contributors:** Erick M. Carreira, Moritz J. Classen  
 Synfacts 2021, 17(02), 0121 Published online: 20.01.2021  
 DOI: 10.1055/s-0040-1719297; Reg-No.: C07920SF

© 2021, Thieme. All rights reserved.  
 Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of Natural Products and Potential Drugs

Key words

(+)-haperforin G

limonoids

tetranortriterpenoids

ring-closing metathesis

Pauson–Khand reaction

light-mediated radical coupling

aldol addition

Synfact  
 of the Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.