


Y. DENG, C.-P. H. YANG, A. B. SMITH, III* (UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, USA)

Enantioselective Total Synthesis of (+)-Peniciketals A and B: Two Architecturally Complex Spiroketal

J. Am. Chem. Soc. **2021**, *143*, 1740–1744, DOI: 10.1021/jacs.0c11424.

Total Synthesis of (+)-Peniciketal A



Significance: Smith and co-workers report the first total synthesis of (+)-peniciketal A, a complex spiroketal isolated from fungus *Penicillium raistrickii*. The benzannulated [6,6]-spiroketal has been shown to possess activity against human cancer cell lines and might serve as a promising drug lead candidate.

Comment: The authors employed a highly convergent approach that combines the two fragments **F** and **G** in a photoisomerization–cyclization cascade. Fragment **F** is generated from epoxide **A** by attack of dithiane **B**; Brook rearrangement then enables alkylation with **D** in a single pot. This is followed by selective silyl deprotection and in situ spiroketalization. Fragment **G** is obtained from atraric acid (**H**) via Negishi coupling and cross-metathesis.

SYNFACTS Contributors: Erick M. Carreira, Patrick Pfaff
Synfacts 2021, 17(04), 0363 Published online: 18.03.2021
DOI: 10.1055/s-0040-1719433; Reg-No.: C01121SF

© 2021, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of Natural Products and Potential Drugs

Key words

(+)-peniciketal A

spiroketals

[1,4]-Brook rearrangement

spiroketalization

photoisomerization

cyclization

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.