



An Interesting Case of Autoimmune Liver Disease

Paras Kathuria¹ Shilpa Arora¹ Rahul Karna² Naresh Kumar³ Suresh Kumar³ Premashis Kar^{3,✉}

¹Department of Medicine, Maulana Azad Medical College and Associated Lok Nayak Hospital, New Delhi, India

²Maulana Azad Medical College, New Delhi, India

³Maulana Azad Medical College and Associated Lok Nayak Hospital, New Delhi, India

Address for correspondence Premashis Kar, DM, PhD, E23 Nivedeta Kunj, Sector 10, R. K. Puram, New Delhi 110022, India (e-mail: premashishkar@gmail.com).

Ann Natl Acad Med Sci (India):2021;57:62–64

Abstract

Autoimmune liver diseases (AILD) are part of a broad spectrum of liver diseases with autoimmune etiology, usually present individually but at times have overlapping features. We present the case of a 60-year-old lady presenting with fatigue, itching and right upper quadrant abdominal pain. Further investigation showed cholestatic pattern of liver enzymes and evidence of portal hypertension without any evidence of extrahepatic obstruction. Autoimmune markers and liver biopsy showed overlapping features of both autoimmune hepatitis (AIH) and primary biliary cirrhosis (PBC), which is a separate diagnosis known as overlap syndrome, but by applying appropriate criterion, we were able to make a definite diagnosis of PBC. Differentiating PBC from overlap syndrome was important as therapy of both are different.

Keywords

- ▶ autoimmune hepatitis
- ▶ cholestasis
- ▶ primary biliary cirrhosis

Introduction

Autoimmune liver diseases (AILD) are part of a spectrum of autoimmune diseases primarily involving the liver and includes autoimmune hepatitis (AIH), primary biliary cirrhosis (PBC), and primary sclerosing cholangitis (PSC). These diseases usually occur in isolation but sometimes their features may overlap, which is a distinct entity with different therapy known as overlap syndrome. Here, we present an atypical case of PBC with features of AIH, thus presenting a diagnostic and therapeutic dilemma. It was important to differentiate PBC from overlap syndrome, as treatment of former focuses on ursodeoxycholic acid while latter needs steroids along with ursodeoxycholic acid.¹

Case Report

A 60-year-old female who was a known case of type 2 diabetes mellitus (DM) and hypothyroidism presented to the clinic with complaints of easy fatigability for 2 months, generalized itching for 1 month, and right hypochondriac pain for 15 days. She denied any history of fever, jaundice, or clay-colored stools. Physical examination

revealed a nontender, firm liver with sharp regular margins palpable 6 cm below costal margin and a palpable spleen 2 cm below costal margin. There was no evidence of scleral icterus, ascites, or stigmata of chronic liver diseases. Initial laboratories showed hemoglobin 8.2 g%, total leukocyte count 8,400 cells/mL, platelet count 2 lacs/mL, liver function tests (LFT) revealed total bilirubin 0.8 mg/dL, aspartate aminotransferase (AST) levels 186 U/L, alanine aminotransferase (ALT) levels 164 U/L, alkaline phosphatase (ALP) 760 μ/L, total protein 7.5 g/dL, albumin 2.5 g/dL, and prothrombin time (PT) 13 seconds. Ultrasonography showed liver span of 17 cm and heterogenous echotexture without any space-occupying lesion (SOL), portal vein diameter 12 mm with hepatopetal flow, normal common bile duct, gall bladder, and spleen 14 cm with normal echotexture without any SOL. Upper gastrointestinal (GI) endoscopy showed single column of grade 1 esophageal varices. Screening tests for hepatitis virus and human immunodeficiency virus were negative. Further laboratories showed thyroid-stimulating hormone (TSH) 60 μIU/mL, antithyroid peroxidase 112.9 IU/mL, positive antimitochondrial antibody (AMA) (4+) and antiliver kidney microsome (1+), IgM levels 7.58 g/L, IgG levels 26.8 g/L and negative

published online
December 30, 2020

DOI <https://doi.org/10.1055/s-0040-1722105>
ISSN 0379-038X.

© 2020. National Academy of Medical Sciences (India).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Thieme Medical and Scientific Publishers Pvt. Ltd. A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

antinuclear antibody, antismooth muscle actin, antisoluble liver antigen, antidouble stranded DNA, antitopoisomerase and anti-U₁ ribonucleoprotein. Magnetic resonance cholangiopancreatography (MRCP) did not show any evidence of intrahepatic biliary radical dilation. Liver biopsy showed lobular architecture disarray with portal-to-portal bridging fibrosis, focal ballooning of hepatocytes, edematous expansion of portal tracts with moderate chronic inflammation and interface hepatitis, bile duct injury and bile ductular proliferation (►Fig. 1). Copper-associated protein was present, indicative of chronic cholangiopathy. The overall picture presented a diagnostic dilemma for us. We initially made a diagnosis of AIH based on modified diagnostic criteria of the International Autoimmune Hepatitis Group, however, on using the extended criterion, that is, Revised Original Scoring System of the International Autoimmune Hepatitis Group, our patient falls in the probable diagnosis (10–15); for definite diagnosis > 15 score is needed.² Also the “Paris Criterion” used in various studies of overlap syndrome was not fulfilled in the case (►Table 1).¹ Hence, a final diagnosis of PBC was made and patient was started on 15 mg/kg UDCA and cholestyramine to which patient responded well both symptomatically and biochemically.




Fig. 1 High power (40x) view of hematoxylin and eosin-stained section of liver biopsy, showing interface hepatitis along with bile duct injury and proliferation.

Table 1 Paris criteria

PBC criteria	AIH criteria
Serum ALP levels at least two times the upper limit of normal values or serum GGT levels as least five times the upper limit of normal values	Serum ALT levels at least five times the upper limit of normal values
A positive test for AMA	Serum IgG levels at least two times the upper limit of normal values or a positive test for SMA
A liver biopsy specimen showing florid bile duct lesions	A liver biopsy showing moderate or severe periportal or periseptal lymphocytic piecemeal necrosis

Abbreviations: AIH, autoimmune hepatitis; ALP, alkaline phosphatase; ALT, alanine aminotransferase; AMA, antimicrobial antibody; GGT, gamma-glutamyl transferase; PBC, primary biliary cirrhosis; SMA, smooth muscle antibody.

Note: PBC-AIH overlap syndrome: 2 out of 3 criteria in each group are met either simultaneously or consecutively.

Discussion

PBC is an autoimmune disease of the liver involving intrahepatic ducts, leading to cholestatic manifestations, seen mainly in middle aged to elderly females. However, patients may also present late with features of cirrhosis and decompensation.³ Apart from historical pointers like pruritis, presence of easy fatigability is also a common feature (60–80%).³ Biochemically, ALP is raised, although AST/ALT may also be slightly increased. Bilirubin generally rises in advanced disease or after development of cirrhosis.⁴ The serological hallmark is the presence of AMA, which is seen in 95% of cases, although anti-PDCE2 or 2-oxo-glutaric acid dehydrogenase complex may also be seen. Liver biopsy is needed for assessing severity of disease or when there are certain atypical manifestations such as AMA-negative cases or raised AST/ALT. There is no definite treatment for PBC and patients are usually treated with ursodeoxycholic acid, which can slow down the course of the disease.

AIH is a chronic inflammatory disorder of liver usually affecting young females and presents with hepatocellular pattern of jaundice. Laboratory features reveal hypergammaglobulinemia, presence of antibodies such as characterized by periportal inflammation, hypergammaglobulinemia, and circulating autoantibodies including antinuclear factor, antisoluble liver antigen and liver kidney microsomal antibody. Interface hepatitis is the hallmark feature of AIH and usually has a better prognosis, being responsive to steroid treatment. Sometimes, the clinical picture does not fit any of these and have overlapping features. Such a case is usually termed as overlap syndrome and can be seen in 3–7% of patients with autoimmune liver disease (AILD).⁵ Recently, a case of overlap syndrome was reported by Bairy et al from south India, who was diagnosed on the basis of Paris criteria and treated with ursodeoxycholic acid.⁶ Rust et al reported another case of overlap syndrome treated with budesonide, ursodeoxycholic acid, and azathioprine.⁷ Differentiating overlap syndrome from AIH and PBC has therapeutic implications, as both ursodeoxycholic acid immunosuppression with steroids

would be required in patients with overlap syndrome. There are reports of beneficial effects of azathioprine and cyclosporine in corticosteroid-resistant patients with overlap syndrome.^{8,9}

In conclusion, overlap syndrome is an important differential to consider in patients with PBC showing features of AIH. However, it is important to differentiate patients with PBC showing AIH features from those of overlap syndrome to prevent unnecessary steroids and other immunosuppressants exposure and side effects.

Note

Written informed consent was taken from the patient who participated in this study.

Funding

None.

Conflict of Interest

None declared.

References

- 1 Chazouillères O, Wendum D, Serfaty L, Montebault S, Rosmorduc O, Poupon R. Primary biliary cirrhosis-autoimmune

- hepatitis overlap syndrome: clinical features and response to therapy. *Hepatology* 1998;28(2):296–301
- 2 Alvarez F, Berg PA, Bianchi FB, et al. International Autoimmune Hepatitis Group Report: review of criteria for diagnosis of autoimmune hepatitis. *J Hepatol* 1999;31(5):929–938
- 3 Sarin SK, Monga R, Sandhu BS, Sharma BC, Sakhuja P, Malhotra V. Primary biliary cirrhosis in India. *Hepatobiliary Pancreat Dis Int* 2006;5(1):105–109
- 4 Amarpurkar DN, Patel ND. Spectrum of autoimmune liver diseases in western India. *J Gastroenterol Hepatol* 2007;22(12):2112–2117
- 5 Czaja AJ. The overlap syndromes of autoimmune hepatitis. *Dig Dis Sci* 2013;58(2):326–343
- 6 Bairy I, Berwal A, Seshadri S. Autoimmune hepatitis - Primary biliary cirrhosis overlap syndrome. *J Clin Diagn Res* 2017;11(7):OD07–OD09
- 7 Rust C, Beuers U. Overlap syndromes among autoimmune liver diseases. *World J Gastroenterol* 2008;14(21):3368–3373
- 8 Chazouillères O, Wendum D, Serfaty L, Rosmorduc O, Poupon R. Long term outcome and response to therapy of primary biliary cirrhosis-autoimmune hepatitis overlap syndrome. *J Hepatol* 2006;44(2):400–406
- 9 Duclos-Vallée JC, Hadengue A, Ganne-Carrié N, Robin E, Degott C, Erlinger S. Primary biliary cirrhosis-autoimmune hepatitis overlap syndrome. Corticoreistance and effective treatment by cyclosporine A. *Dig Dis Sci* 1995;40(5):1069–1073