



Neurological Patients during the Covid-19 Pandemic

Pacientes neurológicos durante a pandemia de Covid-19

Débora Salles¹ Samara Ferreira Santino² Andréa Cristina de Moraes Malinverni^{1,3}
João Norberto Stávale³

¹ Molecular and Experimental Pathology Laboratory I, Universidade Federal de São Paulo, São Paulo, São Paulo, Brazil

² Escola Paulista de Enfermagem, Universidade Federal de São Paulo, São Paulo, São Paulo, Brazil

³ Department of Pathology, Escola Paulista de Medicina, Universidade Federal de São Paulo (EPM/Unifesp), São Paulo, São Paulo, Brazil

Address for correspondence Andréa Cristina de Moraes Malinverni, PhD, Universidade Federal de São Paulo, Rua Pedro de Toledo 781, 5° andar, ramal 1.118, Vila Clementino, São Paulo, SP, Brasil (e-mail: andreamoraesmalinverni@gmail.com).

Arq Bras Neurocir 2023;42(1):e66–e67.

Abstract

With the current pandemic caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), measures of social isolation were necessary, and this resulted in the interruption of several treatments. Regarding neuro-oncological patients, especially those with central nervous system (CNS) disorders, this interruption can cause serious damage or even compromise the success of the treatment in the future. It is essential that each case be evaluated separately to decide how to continue treatment during the pandemic, always considering the risk of SARS-CoV-2 infection and the benefits that the treatment will bring. The policy of not prescribing potentially toxic drugs, chemotherapy, and immunosuppressive therapies, as well as the use of techniques like stereotactic biopsy and telemedicine are important strategies at this time.

Keywords

- ▶ coronavirus
- ▶ neuro-oncology
- ▶ treatment

Resumo

Com a atual pandemia causada pelo coronavírus da síndrome respiratória aguda grave 2 (*severe acute respiratory syndrome coronavirus 2*, SARS-CoV-2, em inglês), foram necessárias medidas de isolamento social que resultaram na interrupção de alguns tratamentos médicos hospitalares. Em relação aos pacientes neuro-oncológicos, principalmente aqueles com distúrbios do sistema nervoso central (SNC), uma interrupção pode causar sérios danos ou até mesmo comprometer o sucesso do tratamento no futuro. Assim, de acordo com a literatura encontrada, é de fundamental importância que cada caso seja avaliado individualmente, para que se decida como prosseguir com o tratamento durante a pandemia, sempre considerando o risco de infecção por SARS-CoV-2 e os benefícios relacionados ao tratamento. São estratégias importantes neste momento a política de não prescrição de medicamentos potencialmente tóxicos, quimioterapia e terapias imunossupressoras, além do uso de técnicas como biópsia estereotáxica e telemedicina..

Palavras-chave

- ▶ coronavírus
- ▶ neuro-oncologia
- ▶ tratamento

received
September 18, 2020
accepted
March 9, 2021

DOI <https://doi.org/10.1055/s-0041-1730336>
ISSN 0103-5355.

© 2022. Sociedade Brasileira de Neurocirurgia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

The current coronavirus disease 2019 (Covid-19) pandemic has brought about measures of social isolation which have resulted in the interruption of several health treatments. Regarding neuro-oncological patients, especially those with central nervous system (CNS) disorders, this interruption can cause serious damage or even compromise the success of the treatment in the future, since surgical resections are being avoided as much as possible, and several rehabilitation services are not available due to the closure of units and the reorientation of teams to combat Covid-19. Therefore, certain information on how to organize these treatments in the current scenario is extremely relevant and should always be followed with due precautions, as these patients are more likely to develop the most severe forms of Covid-19, especially if they have undergone chemotherapy or surgery recently.^{1,2}

First, it is essential that each case be evaluated separately to decide how to continue treatment during the pandemic, always considering the risk of infection by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the benefits that the treatment will bring. In addition, emergency cases or those with a high chance of cure should not be extended.³ It is essential to consider the reuse of available drugs to treat and prevent brain diseases associated with Covid-19.⁴ However, more than ever, the policy of not prescribing potentially toxic drugs, systemic chemotherapy, and immunosuppressive therapies is mandatory, such as re-irradiation combined with steroids, for which the evidence of benefit is not clinically relevant in the current scenario. In situations in which there are no immediate urgent needs and long-term treatment is likely to bring benefits, such as in patients with low-grade astrocytomas or oligodendrogliomas, a more conservative dose of chemotherapy should be considered.⁵

The use of stereotactic biopsy has already been described, which at least offers a definitive diagnosis for severe conditions and can elucidate malignant tumor characteristics, with the advice that the best possible treatment should be followed in view of the current incapacity for maximum tumor resection. At the Mount Sinai Hospital, in New York, patients are always considered positive for Covid-19 until proven otherwise, and maximum precautions have been taken, including regarding the patients in need of intubation and in the postoperative setting.²

Another factor to be considered is telemedicine. In Brazil, the Federal Council of Medicine already recognizes the possibility of remote medical care during the fight against Covid-19, as well as in other countries in which the performance of these virtual consultations is made available when

feasible.⁶ However, for face-to-face procedures, patients should be tested for Covid-19. In institutions in which the test is not available, there should be a questionnaire regarding symptoms, and patients should be tested when available.⁷ In addition, some processes must remain accessible, such as ambulatory radiology, with the establishment of protocols that limit the number of patients in the waiting room and provide protective masks for everyone.⁸

It is also necessary to consider during this period the frequency of clinical follow-up at regular intervals for patients who have been stable and without intervention for years, as is the case of many individuals with brain tumors. Visits to the hospital environment can be divided into defined populations with more severe cases.⁴

Concluding, clinical follow-ups differ in relation to the conditions of the patients, according to the severity of each case. That is why the benefits and risks that the continuity of treatment will bring must be considered, with precaution and engagement on the part of the entire team involved.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Liang W, Guan W, Chen R, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *Lancet Oncol* 2020;21(03):335–337
- 2 Kessler RA, Zimering J, Gilligan J, et al. Neurosurgical management of brain and spine tumors in the COVID-19 era: an institutional experience from the epicenter of the pandemic. *J Neurooncol* 2020;148(02):211–219
- 3 Turri-Zanoni M, Battaglia P, Karligkiotis A, Locatelli D, Castelnovo P. Managing care for patients with sinonasal and anterior skull base cancers during the COVID-19 pandemic. *Head Neck* 2020;42(07):1503–1506
- 4 Saavedra JM. COVID-19, Angiotensin Receptor Blockers, and the Brain. *Cell Mol Neurobiol* 2020;40(05):667–674
- 5 Weller M, Preusser M. How we treat patients with brain tumour during the COVID-19 pandemic. *ESMO Open* 2020;4(Suppl 2):e000789
- 6 Hu YJ, Zhang JM, Chen ZP. Experiences of practicing surgical neuro-oncology during the COVID-19 pandemic. *J Neurooncol* 2020;148(01):199–200
- 7 Zacharia BE, Eichberg DG, Ivan ME, et al. Letter: Surgical Management of Brain Tumor Patients in the COVID-19 Era. *Neurosurgery* 2020;87(02):E197–E200
- 8 Ramakrishna R, Zadeh G, Sheehan JP, Aghi MK. Inpatient and outpatient case prioritization for patients with neuro-oncologic disease amid the COVID-19 pandemic: general guidance for neuro-oncology practitioners from the AANS/CNS Tumor Section and Society for Neuro-Oncology. *J Neurooncol* 2020;147(03):525–529