

The Effect of Neonatal Sepsis on Risk of Autism Diagnosis

Darios Getahun, MD, PhD, MPH^{1,2} Michael J. Fassett, MD^{3,4} Anny H. Xiang, PhD¹ Vicki Y. Chiu, MS¹
Harpreet S. Takhar, MPH¹ Sally F. Shaw, MPH, DrPH¹ Morgan R. Peltier, PhD⁵

¹Department of Research and Evaluation, Kaiser Permanente Southern California, Pasadena, California

²Department of Health Systems Science, Kaiser Permanente Bernard J. Tyson School of Medicine, Pasadena, California

³Department of Obstetrics and Gynecology, Kaiser Permanente West Los Angeles Medical Center, Los Angeles, California

⁴Department of Clinical Sciences, Kaiser Permanente Bernard J. Tyson School of Medicine, Pasadena, California

⁵Department of Foundations of Medicine, NYU-Long Island School of Medicine, Mineola, New York

Address for correspondence Darios Getahun, MD, PhD, MPH, Department of Research and Evaluation, Kaiser Permanente Southern California, 100 S. Los Robles Avenue, 2nd Floor, Pasadena, CA 91101 (e-mail: darios.t.getahun@kp.org).

Am J Perinatol 2023;40:858–866.

Abstract

Objective The study aimed to examine the association between neonatal sepsis and autism risk among children and whether the risk varied with the timing of exposure, child's sex, and race/ethnicity.

Study Design We conducted a retrospective cohort study using electronic health records (EHR) extracted from Kaiser Permanente Southern California Health Care System. Mother–child dyads were constructed by linking records of children born to member mothers and continuing to receive care through the system during the follow-up period with those of their biological mothers ($n = 469,789$). Clinical health records were used to define neonatal sepsis. Diagnosis of autism was made by medical specialists. Potential confounders included maternal sociodemographic factors, obstetrical history, child's age, sex, race/ethnicity, and maternal and child medical history. Incident rates and adjusted hazard ratios (aHR) were used to estimate the associations.

Results Compared with children without the diagnosis of autism, children with the condition were more likely to be from Asian/Pacific Islander descent and male sex. Exposed children showed higher rates of autism as compared with unexposed children (3.43 vs. 1.73 per 1,000 person-years, aHR: 1.67–95% confidence interval [CI]: 1.39–2.00). Both preterm (aHR: 1.47; 95% CI: 1.09–1.98) and term (aHR: 1.63; 95% CI: 1.29–2.06) births were associated with increased risk for autism. Although the magnitude of the HRs and incidence ratios for neonatal sepsis to increase autism risk varied between race ethnicities, neonatal sepsis was associated with significantly increased likelihood of autism diagnosis for all race-ethnic groups except for Asian/Pacific Islanders. Although neonatal sepsis was associated with significantly increased autism risk for both boys and girls, incident rates and HR point estimates suggested that the effect may be stronger in girls.

Conclusion Neonatal sepsis is associated with increased risk of autism diagnosis in preterm- and term-born children. The association was significant for both girls and boys and all race ethnicities except for Asian-Pacific Islanders.

Keywords

- ▶ neonatal sepsis
- ▶ autism
- ▶ autism spectrum disorder
- ▶ race
- ▶ ethnicity
- ▶ race/ethnicity

received

February 1, 2021

accepted after revision

May 12, 2021

article published online

July 5, 2021

© 2021. Thieme. All rights reserved.
Thieme Medical Publishers, Inc.,
333 Seventh Avenue, 18th Floor,
New York, NY 10001, USA

DOI <https://doi.org/10.1055/s-0041-1731648>.
ISSN 0735-1631.

Key Points

- Neonatal sepsis is associated with increased risk of autism diagnosis.
- The association was significant in preterm- and term-born children.
- The association was significant for all race/ethnicities except for Asian-Pacific Islanders.

Autism spectrum disorders (ASD) affect 1 in 59 children in the United States according to the latest data from the Centers for Disease Control and Prevention (CDC).¹ Its prevalence has increased dramatically over the last two decades.¹ The condition persists into adulthood and poses a significant burden to families and healthcare systems.^{2,3} In 2015, the annual direct healthcare expenditure associated with ASD and related costs were estimated at \$268 billion.¹ This accounts for 0.9 to 2.0% of the nation's gross domestic product and exceeds the costs of hypertension and stroke combined.⁴ Due to its fast-growing prevalence, the societal and healthcare costs will continue to rise in years to come.

The patho-etiology mechanism of ASD is largely unknown, but both genetic and environmental factors have been proposed.⁵⁻⁷ These include a history of mood disorders in family members, having a sibling with autism, advanced parental age, white race/ethnicity, and male sex.^{2,7-11} Medical and obstetrical factors including pregestational and gestational diabetes,^{12,13} thyroid disorders,¹⁴ hyperemesis gravidarum,¹⁵ multiple births, preterm birth, small for gestational age birth, preeclampsia, and child medical history have also been linked with an increased risk of ASD.¹⁶ Neonatal sepsis is a significant cause of infant morbidity and mortality, especially in neonates born preterm, at very low birth weight, or of black race. The incidence of the condition, especially for early onset (<72 hours) neonatal sepsis, has decreased since the implementation of guidelines¹⁷⁻¹⁹ in the mid-1990s for intrapartum antibiotic prophylaxis to prevent group B streptococcus (GBS) infections.²⁰ However, infections still affect as many as 2 in 1,000 live births.^{21,22} Neonatal sepsis is not only a major cause of hospitalization, but may also have profound effects on brain development that are not apparent until months and years later as the infant matures. Studies with animal models and human subjects have linked neonatal sepsis with adverse neurodevelopment.²³⁻²⁷ However, all were limited either by sample size or that they only considered short-term adverse neurological outcomes. Although the neonatal sepsis-related mortality rate has dropped in recent years, the impact of neonatal sepsis on the long-term neurodevelopmental sequelae of these surviving children is largely unknown. Furthermore, the risk-modifying effects of timing of diagnosis, gestational age at birth, the child sex, and race/ethnicity on associations between neonatal sepsis and autism require further investigation. Therefore, we examined medical records from a large, sociodemographically diverse population to evaluate the extent to which neonatal sepsis is associated with ASD, and how it is modified by timing of

sepsis diagnosis, the gestational age at birth, child's sex, and race/ethnicity.

Materials and Methods

Kaiser Permanente Southern California (KPSC) is a large integrated healthcare system that provides services to over 4.7 million members at 15 hospitals and 234 medical offices throughout Southern California. KPSC's member population is demographically diverse and broadly representative of the racial/ethnic groups living in Southern California. Services are provided through healthcare insurance with pharmaceutical benefits through group plans, individual plans, Medicare, Medicaid, and other low-income programs. On average, 40,000 pregnancies are delivered annually in the KPSC system, and the 5-year retention rate for these children is 75.4%.

We conducted a retrospective cohort study of singleton live born children delivered in all KPSC hospitals from January 1, 1991, to December 31, 2014 ($n = 469,789$; ►Appendix 1 [Fig. 1], available in the online version). Children were longitudinally linked to biological mothers by using unique maternal and infant medical record numbers. To be eligible for the study, children had to be born from a member mother and remain a member patient during the follow-up period (follow-up ended in December 31, 2018). The electronic health records (EHR) and pharmacy records contain detailed clinical information as well as drug-dispensing data on medications used to treat the mother and her baby for neonatal sepsis (antibiotics and immunoglobulin). Data on potential confounders and mediators that were extracted from EHRs included: maternal age (<20, 20-29, 30-34, and ≥ 35 years); educational attainment (<12, 12, and ≥ 13 years); median family household income based on census tract of residence; parity; smoking during pregnancy (yes/no); timing of prenatal care initiation (first trimester or late/no care); pre-pregnancy body mass index (BMI, kg/m²) and gestational weight gain; child's characteristics (child's age, sex, and race/ethnicity); and obstetrical history (gestational diabetes [GDM], hyperemesis gravidarum, birth weight, gestational age at delivery based on clinical estimates, Apgar's score, and mode of delivery [cesarean vs. vaginal] and maternal and child medical history [chronic hypertension, pregestational diabetes, hypo/hyperthyroidism, neonatal respiratory distress syndrome [RDS], bronchopulmonary dysplasia [BPD], mechanical ventilation). The accuracy of variables utilized in this study has been previously validated.²⁸⁻³⁰ Maternal and paternal race/ethnicity defined the child's race/ethnicity. A child was categorized non-Hispanic White (White) if born to non-Hispanic White mother and father. The same convention




Fig. 1 Kaplan–Meier curves for the accumulation of autism spectrum disorder diagnosis in children with and without sepsis in general (A) and stratified by sex (B).

was applied to non-Hispanic Black (Black), Hispanic, and Asian/Pacific Islander race/ethnicity groups. The other/mixed race/ethnicity category includes children born from interracial/interethnic relationships. Institutional review board approval with exemption of informed consent was obtained for this study.

Exposure and Outcome

The primary exposure measure was clinical diagnosis of sepsis during the neonatal period ascertained from KPSC's EHRs, regional claim records, and laboratory records. Neonatal sepsis was defined as meeting the following criteria: ICD-9-CM codes for septicemia (038.x) and septic shock (785.5x).

To ascertain a clinical diagnosis of ASD, we used the diagnostic criteria of Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV-TR)³¹ for ASD: autistic disorder, childhood disintegrative disorder, Rett's disorder, Asperger's disorder, or pervasive developmental disorder, not otherwise specified. Children, ages between 2 and 17 years, with at least one documented DSM-IV-TR code for ASD on any two separate visits during the follow-up period formed the ASD cases. Clinical experts validated the accuracy of ASD codes through medical record review.³² As per KPSC guidelines, last revised April 2013,³³ ASD services are covered by healthcare providers (i.e., child/adolescent psychiatrist, developmental/behavioral pediatrician, and child psychologist, or neurologist) who are experts in taking care of children with neurodevelopmental conditions. The experts are required to perform a series of behavioral and developmental surveillance at all well-child visits, as early as 4 months of age. If there is a sufficient degree of suspicion, a modified version of the Checklist for Autism in Toddlers [M-CHAT]³⁴ and developmental screening questionnaires for toddlers will be completed (as early as 18 months of life). For each diagnosed child, a treatment plan is developed and provided by a provider who is qualified in autism services. These criteria ensure that there is consistent testing and opportunity for diagnosis across all KPSC centers during the duration of the study.

Statistical Analysis

Perinatal and neonatal characteristics of children were compared by ASD status using Chi-square tests. Hazard ratios (HR) and their 95% confidence intervals (CI) derived from Cox's proportional hazard models were used to estimate the magnitude of associations before and after accounting for potential confounding factors. Follow-up of the children started from the delivery date until the first date of ASD diagnosis. Censoring occurred on the earliest of the following dates: health plan disenrollment, 17th birthday, non-ASD related death, or end of study (December 31, 2018).

Subgroup analyses for potential mediators: based on the timing of neonatal sepsis exposure, child's race/ethnicity, child's sex, and gestational age at birth were performed. Furthermore, to assess the relative influences of maternal and infant factors: periventricular leukomalacia, neonatal convulsion, other postnatal factors, maternal medical conditions (ischemic hypoxic conditions, endocrine abnormalities, hyperemesis gravidarum on the observed risk relationships, a sensitivity analysis was performed (→ Appendix 2 [Table 1], available in the online version) after (1) excluding children with only one ASD diagnosis; (2) excluding children with a history of congenital anomalies and developmental and emotional comorbidities (mental retardation, developmental dyslexia, deficits in language processing, conduct disorder, irritability, bipolar or anxiety disorders, and depression); (3) excluding pregnancies with a history of medical and/or perinatal adversities including gestational hypertension, diabetes, intrauterine growth restriction (IUGR) and low birth weight; (4) excluding children aged 2 to 3 years who may have had shorter follow-up period to capture events; (5) limiting the cohort to children without genetic predisposition; (6) accounting for maternal comorbidities (→ Appendix 2 [Table 2], available in the online version); (7) accounting for "year of birth"; (8) accounting for maternal pre-pregnancy BMI; and (9) examining the potential effects of residual confounders on the observed association using a range of E-values for the overall result and the lower limits of their 95% CI observed in Cox's regression

Table 1 Distribution of maternal and infant characteristics based on autism spectrum disorders status

| Characteristics | No ASD n = 461,342 | ASD n = 8,447 | p-Value |
|---|-----------------------|------------------|---------|
| Maternal age (%) | | | <0.001 |
| < 20 y | 27,632 (98.9) | 299 (1.1) | |
| 20–29 y | 200,884 (98.4) | 3,212 (1.6) | |
| 30–34 y | 140,057 (98.1) | 2,729 (1.9) | |
| ≥35 y | 92,769 (97.7) | 2,207 (2.3) | |
| Maternal education (%) | | | <0.001 |
| < 12 y | 47,188 (98.9) | 530 (1.1) | |
| 12 y | 130,869 (98.4) | 2,127 (1.6) | |
| ≥13 y | 268,056 (97.9) | 5,629 (2.1) | |
| Median family household income ^a (%) | | | <0.001 |
| < \$29,999 | 45,589 (98.4) | 744 (1.6) | |
| \$30,000–\$49,999 | 158,634 (98.3) | 2,768 (1.7) | |
| \$50,000–\$69,999 | 137,170 (98.1) | 2,658 (1.9) | |
| \$70,000–\$89,999 | 70,288 (98.1) | 1,344 (1.9) | |
| ≥\$90,000 | 46,807 (98.1) | 889 (1.9) | |
| Parity (%) | | | <0.001 |
| Parity 0 | 177,881 (97.9) | 3,857 (2.1) | |
| Parity 1 | 156,302 (98.2) | 2,808 (1.8) | |
| Parity ≥2 | 127,146 (98.6) | 1,782 (1.4) | |
| Smoking during pregnancy (%) | 31,433 (98.4) | 527 (1.6) | 0.038 |
| Late initiation of prenatal care (%) | 42,166 (98.5) | 649 (1.5) | <0.001 |
| Mode of delivery (%) | | | <0.001 |
| Cesarean | 109,624 (97.6) | 2,643 (2.4) | |
| Vaginal | 351,120 (98.4) | 5,794 (1.6) | |
| Child race/ethnicity (%) | | | <0.001 |
| Non-Hispanic White | 98,821 (98.2) | 1,843 (1.8) | |
| Non-Hispanic Black | 39,940 (98.2) | 741 (1.8) | |
| Hispanic | 172,994 (98.3) | 2,948 (1.7) | |
| Asian/Pacific Islander | 39,927 (97.9) | 844 (2.1) | |
| Other/mixed ^b | 103,461 (98.1) | 2,010 (1.9) | |
| Child's sex (%) | | | <0.001 |
| Female | 227,859 (99.3) | 1,589 (0.7) | |
| Male | 233,483 (97.1) | 6,858 (2.9) | |
| Gestational age at birth (%) | | | <0.001 |
| 28–36 wk | 33,536 (97.6) | 820 (2.4) | |
| 37–42 wk | 427,806 (98.2) | 7,627 (1.8) | |
| Birth weight (%) | | | <0.001 |
| < 1,500 g | 2,469 (97.0) | 77 (3.0) | |
| 1,500–2,499 g | 19,162 (97.6) | 469 (2.4) | |
| ≥2,500 g | 439,709 (98.2) | 7,901 (1.8) | |
| SGA birth at <10th percentile (%) | 41,529 (98.0) | 853 (2.0) | 0.002 |
| Apgar's score <7 at 1 min (%) | 23,181 (97.7) | 549 (2.3) | <0.001 |
| Apgar's score <7 at 5 min (%) | 2,689 (97.2) | 77 (2.8) | <0.001 |
| Neonatal RDS (%) | 9,336 (96.9) | 298 (3.1) | <0.001 |

(Continued)

Table 1 (Continued)

| Characteristics | No ASD <i>n</i> = 461,342 | ASD <i>n</i> = 8,447 | <i>p</i> -Value |
|---|------------------------------|-------------------------|-----------------|
| Bronchopulmonary dysplasia (%) | 199 (91.3) | 19 (8.7) | <0.001 |
| Bilirubin >15 mg/dL (%) | 26,845 (97.6) | 658 (2.4) | <0.001 |
| Mechanical ventilation (%) | 5,659 (96.7) | 195 (3.3) | <0.001 |
| NICU admission (1999–2014) ^c (%) | 36,242 (97.1) | 1,100 (2.9) | <0.001 |

Abbreviations: ASD, autism spectrum disorder; SGA, small for gestational age at <10th percentile; RDS, respiratory distress syndrome; NICU, neonatal intensive care unit.

^aMedian household income based on census tract information, values in U.S. dollars.

^bOther/mixed racial/ethnic category includes non-Hispanic children with multiple recorded races.

^cNICU admission (1999–2014), data on NICU admission was available starting from 1999.

Note: Row percentages in the table are with respect to the total number of observations in each row.

models.³⁵ The E-values were estimated by using the online calculator for HR with an outcome prevalence of <15%. All analyses were performed by using SAS version 9.4 (SAS institute, Cary, NC).

Results

Mothers of children with ASD were more likely to be older, have ≥13 years of formal education, have a high median family household income, have a history of smoking during pregnancy, and to be nulliparous (→Table 1) than mothers of children without the condition. Compared with children without ASD diagnosis, children with the condition were more likely to have been delivered by cesarean section, have

birth weight <2,500 g and shorter gestational ages, have an Apgar's score <7 at 1 and 5 minutes, be from Asian/Pacific Islander racial/ethnic group, be male sex (4:1 male to female ratio), and have the diagnosis of BPD and RDS and require mechanical ventilation during neonatal period (→Table 1).

Between 1991 and 2014, a total of 3,582 (0.8%) neonates were diagnosed with neonatal sepsis and were admitted to KPSC hospitals with NICU facilities (→Table 2). Of those children who met the diagnostic criteria for neonatal sepsis: 2,817 (79%) had the diagnosis made within 24 hours of birth, 162 (4.5%) had the diagnosis between 25 and 72 hours, and 603 (16.8%) had the diagnosis made after 72 hours. Distribution of mean child's age at ASD diagnosis in those with (mean ± standard deviation [SD]) = 5.13 ± 3.55 years) and

Table 2 Associations between neonatal sepsis and autism spectrum disorders

| Conditions | Total births <i>n</i> = 469,789 | ASD <i>n</i> (IR) ^a | Hazard Ratio (95% CI) | |
|--------------------------|------------------------------------|-----------------------------------|-----------------------|-----------------------|
| | | | Unadjusted | Adjusted ^b |
| No sepsis | 466,207 | 8,330 (1.73) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 3,582 | 117 (3.43) | 1.88 (1.57–2.26) | 1.67 (1.39–2.00) |
| Diagnosis timing | | | | |
| ≤3 d of age | 2,979 | 105 (3.77) | 2.05 (1.69–2.48) | 1.79 (1.47–2.17) |
| < 1 d of age | 2,817 | 100 (3.80) | 2.06 (1.69–2.51) | 1.80 (1.43–2.19) |
| 2–3 d of age | 162 | 5 (3.25) | 1.81 (0.75–4.35) | 1.59 (0.66–3.82) |
| > 3 d of age | 603 | 12 (1.92) | 1.10 (0.62–1.93) | 1.04 (0.59–1.84) |
| Gestational age at birth | | | | |
| 28–33 wk | 665 | 27 (4.29) | 1.60 (1.06–2.40) | 1.61 (1.07–2.42) |
| 34–36 wk | 586 | 18 (3.21) | 1.37 (0.86–2.19) | 1.23 (0.77–1.97) |
| 37–42 wk | 2,331 | 72 (3.24) | 1.83 (1.45–2.30) | 1.63 (1.29–2.06) |
| Birth weight | | | | |
| < 1,500 g | 346 | 17 (5.26) | 1.25 (0.80–1.95) | 1.79 (1.04–3.08) |
| 1,500–2,499 g | 678 | 20 (3.07) | 1.88 (1.10–3.23) | 1.13 (0.72–1.77) |
| ≥2,500 g | 2,558 | 80 (3.29) | 1.84 (1.47–2.29) | 1.63 (1.30–2.03) |

Abbreviations: ASD, autism spectrum disorder; CI, confidence interval.

^aIncidence rate per 1,000 person-years.

^bHazard ratios were adjusted for maternal age, education, smoking during pregnancy, parity, prenatal care, household income, child race/ethnicity, and sex.

without (mean \pm SD = 5.10 \pm 3.55 years) neonatal sepsis were comparable. The number of neonates with neonatal sepsis differed based on gestational age at delivery: 820 (2.4%) at early preterm (28–36 weeks) and 7,627 (1.8%) at term gestation.

Compared with other children, neonates with the diagnosis of sepsis were more likely to have the diagnosis of ASD (1.73/1,000 person-years vs. 3.43/1,000 person-years; adjusted odds ratio [aHR]: 1.67; 95% CI: 1.39–2.00; **Table 2**; **Fig. 1A**). Early neonatal sepsis was associated with a 2.06-fold increased risk of ASD (95% CI: 1.69–2.51). The observed associations of neonatal sepsis with ASD persisted after controlling for several potential confounding factors.

The incidence of ASD diagnosis varied based on child's sex and racial/ethnic background (**Table 3**). Neonatal sepsis significantly increased the risk of ASD diagnoses starting at the age of 4 years in boys and 5 years in girls (**Fig. 1B**). Although HR for neonatal sepsis was significant for boys and girls, the IR differences and HR were slightly higher for girls (**Table 3**), suggesting a stronger relationship. However, tests for heterogeneity of regression analysis of the

Kaplan–Meier plots did not detect an interaction between sepsis and child's sex ($p = 0.573$), suggesting similar impact on male and female children. We also did not detect significant interaction between race/ethnicity and neonatal sepsis on the risk of ASD ($p = 0.445$). Neonatal sepsis does seem to influence the incidence (**Appendix 3 [Fig. 1]**, available in the online version) and risk of ASD in both preterm (aHR: 1.47; 95% CI: 1.09–1.98) and term (aHR: 1.63; 95% CI: 1.29–2.06) gestation.

In a series of sensitivity analyses, we evaluated how different factors such as only one ASD diagnosis, a history of developmental and emotional comorbidities, congenital malformation, excluding children diagnosis prior to age 4 years not confirmed at later age, year of birth, familial predisposition to ASD, maternal BMI, and maternal medical and obstetrical comorbidities may have affected the observed associations. The finding of the overall analysis persisted after accounting for these factors (**Appendix 2 [Table 1]**, available in the online version). Furthermore, we computed an E-value to assess bias due to potential unmeasured confounders³⁵ and estimated it at 2.73 (95% CI: 2.66–2.80) for the overall adjusted

Table 3 Rates of neonatal sepsis and its associations with autism spectrum disorders by child's sex and race/ethnicity

| Child's characteristics | Total births <i>n</i> | ASD, <i>n</i> (IR) ^a | Hazard ratios (95% CI) ^b | |
|--------------------------|--------------------------|------------------------------------|-------------------------------------|------------------|
| | | | Crude | Adjusted |
| Female sex | | | | |
| No sepsis | 227,927 | 1,569 (0.66) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 1,521 | 20 (1.37) | 1.98 (1.27–3.08) | 1.90 (1.22–2.95) |
| Male sex | | | | |
| No sepsis | 238,280 | 6,761 (2.76) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 2,061 | 97 (4.98) | 1.70 (1.39–2.08) | 1.63 (1.33–1.98) |
| Non-Hispanic White | | | | |
| No sepsis | 100,117 | 1,821 (1.75) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 547 | 22 (4.22) | 2.31 (1.52–3.52) | 2.01 (1.32–3.06) |
| Non-Hispanic Black | | | | |
| No sepsis | 40,294 | 728 (1.56) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 387 | 13 (3.30) | 1.98 (1.15–3.43) | 1.96 (1.13–3.38) |
| Hispanic | | | | |
| No sepsis | 174,419 | 2,903 (1.63) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 1,523 | 45 (3.12) | 1.80 (1.34–2.42) | 1.56 (1.16–2.10) |
| Asian/Pacific Islander | | | | |
| No sepsis | 40,476 | 839 (2.10) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 295 | 5 (1.79) | 0.82 (0.34–1.98) | 0.75 (0.31–1.80) |
| Other/mixed ^c | | | | |
| No sepsis | 104,653 | 1,978 (1.89) | 1.00 (reference) | 1.00 (reference) |
| Sepsis | 818 | 32 (4.24) | 2.13 (1.50–3.02) | 1.91 (1.34–2.70) |

Abbreviations: ASD, autism spectrum disorder; CI, confidence interval.

^aIncidence rate per 1,000 person-years.

^bHazard ratios were adjusted for maternal age, education, smoking during pregnancy, perinatal care, parity, household income, child's sex, and year of diagnosis.

^cOthers, non-Hispanic children with multiple recorded races.

HR of 1.67 (95% CI: 1.39–2.00). Therefore, a minimum risk ratio of 2.66 would be required between unmeasured confounders and neonatal sepsis exposure, and between unmeasured confounders and ASD to account for the significant association we found between neonatal sepsis and child ASD.

Discussion

Using a large socioeconomically and racially/ethnically diverse sample of children, we demonstrated that neonatal sepsis is associated with increased risk of ASD. This association could not be explained by potential confounding factors, and any unknown, unmeasured, confounders would have to be unusually strong to impact the validity of our findings. The magnitude of association differed by the timing of exposure and was strongest when the exposure occurred during the first 24 hours of life. Neonatal sepsis is associated with increased ASD risk, regardless of the gestational age at birth. We detected an effect of neonatal sepsis for all racial/ethnic groups studied except for Asian/Pacific Islanders. This and the variability in the strength of the association as measured by IRs and HRs suggest that race/ethnicity may be a risk modifying factor. Although both girls and boys with sepsis were at increased risk for ASD, both the IRR (107 vs. 80%) and the HR (1.90 vs. 1.63) were higher for girls suggesting a stronger relationship; however, we did not detect a significant interaction. Therefore, further studies are needed to explore the possibility that different levels of ASD risk associated with neonatal sepsis.

Neonates, especially those born at a very preterm gestation, are at increased risk for developing severe inflammation. In the United States, the incidence of neonatal sepsis has decreased since the mid-1990s²⁰ due in part to the CDC,¹⁷ the American College of Obstetricians and Gynecologists,¹⁸ and the American Academy of Pediatrics¹⁹ recommendation for intrapartum antibiotic prophylaxis against GBS infection. Despite the implementation of this recommendation and other advancements in neonate care since then, neonatal sepsis remains a significant cause of infant morbidity and mortality. Immaturity or underdevelopment of the immune system has been previously proposed to increase the infants vulnerability to microorganism invasions and systemic infection during the neonatal period.³⁶ However, a different view on the patho-etiology of neonatal susceptibility to infections has emerged in recent years. Using animal models, Elahi et al³⁷ demonstrated that CD71+ cell-mediated immune suppression during neonatal period, as opposed to the immaturity of immune cells, make neonates vulnerable to developing infections.³⁷ Sepsis-induced inflammation and oxidative stress on neurons have often been reported for immuno-compromised patients. The same mechanism may be at work in neonates whose brain cellular components are especially susceptible to oxidative damage from infections resulting in adverse neurodevelopment.

Our findings are consistent with a previous study performed in Denmark that also reported an association between neonatal sepsis and ASD that appeared to be dependent on

gestational age at birth.³⁸ However, we also found significant race/ethnicity differences as well as timing of neonatal sepsis diagnosis that modify the observed associations.

Neonatal sepsis at a vulnerable period in brain development increases the risk for long-term adverse outcomes that impact the child's behavior and achievement. This is most clear when it occurs in the first 24 hours of life, as it has been shown in our study. This association has been supported by previous animal model and human epidemiological studies demonstrating an increase in the white matter injury and periventricular leukomalacia in early-onset neonatal sepsis.^{39–43} Furthermore, changes in synapse formation that may be affected by inflammation that could also be associated with ASD has been suggested.⁴⁴ Moreover, adjusting for potential pre- and postnatal factors did not affect the magnitude of association, suggesting an independent effect of neonatal sepsis on ASD risk.

Consistent with recent studies, we found substantial sex-related variation in the associations between perinatal factors and ASD diagnosis rates.^{7,14,15} After controlling for several confounding factors, ASD diagnosis rates were higher in males. However, the ASD risk in relation to neonatal sepsis was similar in both boys and girls (no interaction between sepsis and child's sex) suggesting that there are no neuroprotective effects of estrogens on sepsis-mediated neurodevelopmental changes in these very young age children as there is in older women with Parkinson's disease^{45,46} and Alzheimer's disease.^{47,48} Our study also shows important race/ethnic variation in ASD diagnosis and its relation to a history of neonatal sepsis. Using a stratified analysis, we found that the risk of ASD later in life was significantly higher in White, Black, Hispanic, and other racial/ethnic groups but not in Asian/Pacific Islander children with a history of neonatal sepsis compared with children from respective racial/ethnic background without a history of neonatal sepsis. These variations in the incidence rates of neonatal sepsis and the heterogeneity in ASD risk across racial/ethnic groups observed in our study may stem from differences in the rate of intrapartum antibiotic prophylaxis utilization and the known racial/ethnic differences in the rates of neonatal sepsis, preterm birth and low birth weight,^{21,49,50} but genetic polymorphism variation and socioeconomic deprivation may also contribute.⁵¹ Although we adjusted for gestational age, birth weight, maternal education and median family income in the race/ethnicity-specific analysis, we lack data on genetic polymorphisms that may account for these differences. The use of older siblings and mothers with ASD diagnosis as a proxy for genetic predisposition did not alter our findings (–Appendix 2 [Table 1], available in the online version), however.

Strengths of this study include health information extracted from a large integrated health care system's EHRs, use of a sociodemographically diverse population that is representative of the Southern California population, and controlling for several potential confounding factors. Other key strengths of this study are ASD diagnosis by medical specialists, maternal-child medical history extracted from virtually complete and already validated EHRs,^{28–30} and a follow-up period from birth until 17 years

of age. This ensures that all children in the study had the same baseline probability of being diagnosed with ASD by medical professionals.

Like all epidemiologic observational studies, this one also has some limitations. First, causality cannot be established because of the retrospective nature of this study design. Second, data on history of maternal smoking during pregnancy, a known risk factor for childhood behavioral and cognitive impairments,⁵² was self-reported by mothers at prenatal visits. The validity of self-reported tobacco use is often questioned because of pressure women feel to not smoke during pregnancy may result in underreporting of this common habit. However, Buka et al⁵³ found strong agreement between self-reported smoking and serum cotinine levels suggesting that self-report is a valid tool for quantifying tobacco use in pregnancy. Third, surveillance bias can occur when exposed and unexposed children differ in measurement (the intensity and diagnostic process) of health outcome during the follow-up period leading to unequal ascertainment of ASD events. We minimized this possibility by including only clinically diagnosed and validated ASD cases.^{32,33}

Conclusion

Our findings of this study suggest that neonatal sepsis is a predictor of ASD risk later in life. Therefore, preventive measures for early neonatal sepsis may be crucial to preventing adverse neurodevelopmental sequelae that do not become apparent until later in life. The strong association observed between neonatal sepsis within 24 hours of life and ASD risk suggest that early inflammatory processes may have profound effects. The association between neonatal sepsis and ASD risk is higher for White, Black, and Hispanic children, but not for Asian/Pacific Islanders. The drivers of these disparities are not clear and need to be investigated further.

Note

The opinions expressed are solely the responsibility of the authors and do not necessarily reflect the official views of the Kaiser Permanente Community Benefit Funds.

Authors' Contributions

D.G. obtained funding, conceptualized and designed the study, acquired data, analyzed and interpreted data, drafted the initial manuscript, and reviewed and revised the manuscript for important intellectual content. M.J.F., A.H.X., S.F.S., M.R.P., and H.S.T. conceptualized and designed the study and critically reviewed the manuscript for important intellectual content. V.Y.C. designed the study, acquired data, performed analyses, and reviewed and revised the manuscript.

Funding

This study was supported in part by Kaiser Permanente Direct Community Benefit Funds.

Conflict of Interest

None declared.

References

- Center for Disease Control and Prevention Autism Spectrum Disorder (ASD) Data and Statistics on Autism Spectrum Disorder. Accessed December 27, 2020 at: <https://www.cdc.gov/ncbddd/autism/data.html>
- Newschaffer CJ, Croen LA, Daniels J, et al. The epidemiology of autism spectrum disorders. *Annu Rev Public Health* 2007; 28:235–258
- Biao J Autism and Developmental Disabilities Monitoring Network Surveillance Year 2008 Principal Investigators Centers for Disease Control and Prevention. Prevalence of autism spectrum disorders—autism and developmental disabilities monitoring network, 14 sites, United States, 2008. *MMWR Surveill Summ* 2012;61(03): 1–19
- Leigh JP, Du J. Brief report: forecasting the economic burden of autism in 2015 and 2025 in the United States. *J Autism Dev Disord* 2015;45(12):4135–4139
- Hallmayer J, Cleveland S, Torres A, et al. Genetic heritability and shared environmental factors among twin pairs with autism. *Arch Gen Psychiatry* 2011;68(11):1095–1102
- Bailey A, Le Couteur A, Gottesman I, et al. Autism as a strongly genetic disorder: evidence from a British twin study. *Psychol Med* 1995;25(01):63–77
- Xie F, Peltier M, Getahun D. Is the risk of autism in younger siblings of affected children moderated by sex, race/ethnicity, or gestational age? *J Dev Behav Pediatr* 2016;37(08):603–609
- Fombonne E. Epidemiology of autistic disorder and other pervasive developmental disorders. *J Clin Psychiatry* 2005;66 (Suppl 10):3–8
- Burd L, Severud R, Kerbeshian J, Klug MG. Prenatal and perinatal risk factors for autism. *J Perinat Med* 1999;27(06):441–450
- Larsson HJ, Eaton WW, Madsen KM, et al. Risk factors for autism: perinatal factors, parental psychiatric history, and socioeconomic status. *Am J Epidemiol* 2005;161(10):916–925, discussion 926–928
- Croen LA, Grether JK, Selvin S. Descriptive epidemiology of autism in a California population: who is at risk? *J Autism Dev Disord* 2002;32(03):217–224
- Xiang AH, Wang X, Martinez MP, et al. Association of maternal diabetes with autism in offspring. *JAMA* 2015;313(14): 1425–1434
- Xiang AH, Chow T, Martinez MP, et al. Hemoglobin A1c levels during pregnancy and risk of autism spectrum disorders in offspring. *JAMA* 2019;322(05):460–461
- Getahun D, Jacobsen SJ, Fassett MJ, et al. Association between maternal hypothyroidism and autism spectrum disorders in children. *Pediatr Res* 2018;83(03):580–588
- Getahun D, Fassett MJ, Jacobsen SJ, et al. Autism spectrum disorders in children exposed in utero to hyperemesis gravidarum. *Am J Perinatol* 2019
- Getahun D, Fassett MJ, Peltier MR, et al. Association of perinatal risk factors with autism spectrum disorder. *Am J Perinatol* 2017; 34(03):295–304
- Centers for Disease Control and Prevention. Prevention of perinatal group B streptococcal disease: a public health perspective. *MMWR Recomm Rep* 1996;45(RR-7):1–24
- Committee on Obstetric Practice. American College of Obstetrics and Gynecologists. ACOG committee opinion. Prevention of early-onset group B streptococcal disease in newborns. Number 173–June 1996. *Int J Gynaecol Obstet* 1996;54(02):197–205
- American Academy of Pediatrics Committee on Infectious Diseases and Committee on Fetus and Newborn. Revised guidelines for prevention of early-onset group B streptococcal (GBS) infection. *Pediatrics* 1997;99(03):489–496
- Van Dyke MK, Phares CR, Lynfield R, et al. Evaluation of universal antenatal screening for group B streptococcus. *N Engl J Med* 2009; 360(25):2626–2636

- 21 Weston EJ, Pondo T, Lewis MM, et al. The burden of invasive early-onset neonatal sepsis in the United States, 2005–2008. *Pediatr Infect Dis J* 2011;30(11):937–941
- 22 Bizzarro MJ, Raskind C, Baltimore RS, Gallagher PG. Seventy-five years of neonatal sepsis at Yale: 1928–2003. *Pediatrics* 2005;116(03):595–602
- 23 Singh L, Das S, Bhat VB, Plakkal N. Early neurodevelopmental outcome of very low birthweight neonates with culture-positive blood stream infection: a prospective cohort study. *Cureus* 2018;10(10):e3492
- 24 Bolisetty S, Tiwari M, Sutton L, Schindler T, Bajuk B, Lui K. New South Wales and the Australian Capital Territory Neonatal Intensive Care Units' Data Registry. Neurodevelopmental outcomes of extremely preterm infants in New South Wales and the Australian Capital Territory. *J Paediatr Child Health* 2019;55(08):956–961
- 25 Cardoso FL, Herz J, Fernandes A, et al. Systemic inflammation in early neonatal mice induces transient and lasting neurodegenerative effects. *J Neuroinflammation* 2015;12:82
- 26 Smilga AS, Garfinkle J, Ng P, et al. Neonatal infection in children with cerebral palsy: a registry-based cohort study. *Pediatr Neurol* 2018;80:77–83
- 27 Deykin EY, MacMahon B. Pregnancy, delivery, and neonatal complications among autistic children. *Am J Dis Child* 1980;134(09):860–864
- 28 Getahun D, Rhoads GG, Fassett MJ, et al. Accuracy of reporting maternal and infant perinatal service system coding and clinical utilization coding. *J Med Stat Inform* 2013;1:1–3
- 29 Andrade SE, Scott PE, Davis RL, et al. Validity of health plan and birth certificate data for pregnancy research. *Pharmacoepidemiol Drug Saf* 2013;22(01):7–15
- 30 Koebnick C, Langer-Gould AM, Gould MK, et al. Sociodemographic characteristics of members of a large, integrated health care system: comparison with US Census Bureau data. *Perm J* 2012;16(03):37–41
- 31 American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th Edition Text Revision (DSM-IV-TR) Washington, DC: American Psychiatric Publishing; 2000
- 32 Coleman KJ, Lutsky MA, Yau V, et al. Validation of autism spectrum disorder diagnoses in large healthcare systems with electronic medical records. *J Autism Dev Disord* 2015;45(07):1989–1996
- 33 Clinical Practice Guideline. Kaiser Permanente Southern California Preventive Services for Children and Adolescents. Available at: First Issued: 6–1994, Last Reviewed/Revised: 4–2013. PMID. Accessed June 20, 2019 at: <http://cl.kp.org/pkc/scal/cpg/cpg/html/PrevSvcChild.html>
- 34 Robins DL, Fein D, Barton ML, Green JA. The Modified Checklist for Autism in Toddlers: an initial study investigating the early detection of autism and pervasive developmental disorders. *J Autism Dev Disord* 2001;31(02):131–144
- 35 VanderWeele TJ, Ding P. Sensitivity analysis in observational research: introducing the E-value. *Ann Intern Med* 2017;167(04):268–274
- 36 Raymond SL, Stortz JA, Mira JC, Larson SD, Wynn JL, Moldawer LL. Immunological defects in neonatal sepsis and potential therapeutic approaches. *Front Pediatr* 2017;5:14
- 37 Elahi S, Ertelt JM, Kinder JM, et al. Immunosuppressive CD71+ erythroid cells compromise neonatal host defence against infection. *Nature* 2013;504(7478):158–162
- 38 Atladóttir HÁ, Schendel DE, Parner ET, Henriksen TB. A descriptive study on the neonatal morbidity profile of autism spectrum disorders, including a comparison with other neurodevelopmental disorders. *J Autism Dev Disord* 2015;45(08):2429–2442
- 39 Polin RA. Systemic infection and brain injury in the preterm infant. *J Pediatr (Rio J)* 2008;84(03):188–191
- 40 Shah DK, Doyle LW, Anderson PJ, et al. Adverse neurodevelopment in preterm infants with postnatal sepsis or necrotizing enterocolitis is mediated by white matter abnormalities on magnetic resonance imaging at term. *J Pediatr* 2008;153(02):170–175, 175.e171
- 41 Verboon-Macolek MA, Groenendaal F, Cowan F, Govaert P, van Loon AM, de Vries LS. White matter damage in neonatal enterovirus meningoencephalitis. *Neurology* 2006;66(08):1267–1269
- 42 Han Q, Lin Q, Huang P, et al. Microglia-derived IL-1 β contributes to axon development disorders and synaptic deficit through p38-MAPK signal pathway in septic neonatal rats. *J Neuroinflammation* 2017;14(01):52
- 43 Simonsen KA, Anderson-Berry AL, Delair SF, Davies HD. Early-onset neonatal sepsis. *Clin Microbiol Rev* 2014;27(01):21–47
- 44 Petit-Pedrol M, Sell J, Planagumà J, et al. LGI1 antibodies alter Kv1.1 and AMPA receptors changing synaptic excitability, plasticity and memory. *Brain* 2018;141(11):3144–3159
- 45 Saunders-Pullman R, Gordon-Elliott J, Parides M, Fahn S, Saunders HR, Bressman S. The effect of estrogen replacement on early Parkinson's disease. *Neurology* 1999;52(07):1417–1421
- 46 Rocca WA, Bower JH, Maraganore DM, et al. Increased risk of parkinsonism in women who underwent oophorectomy before menopause. *Neurology* 2008;70(03):200–209
- 47 Tang MX, Jacobs D, Stern Y, et al. Effect of oestrogen during menopause on risk and age at onset of Alzheimer's disease. *Lancet* 1996;348(9025):429–432
- 48 Barron AM, Pike CJ. Sex hormones, aging, and Alzheimer's disease. *Front Biosci (Elite Ed)* 2012;4:976–997
- 49 Singh GK, Kogan MD. Persistent socioeconomic disparities in infant, neonatal, and postneonatal mortality rates in the United States, 1969–2001. *Pediatrics* 2007;119(04):e928–e939
- 50 Schrag SJ, Zywicki S, Farley MM, et al. Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med* 2000;342(01):15–20
- 51 Mustarim M, Yanwirasti Y, Jamsari J, Rukmono R, Nindrea RD. Association of Gene Polymorphism of Bactericidal Permeability Increasing Protein Rs4358188, Cluster of Differentiation 14 Rs2569190, Interleukin 1 β Rs1143643 and Matrix Metalloproteinase-16 Rs2664349 with Neonatal Sepsis. *Open Access Maced J Med Sci* 2019;7(17):2728–2733
- 52 Hultman CM, Sparén P, Cnattingius S. Perinatal risk factors for infantile autism. *Epidemiology* 2002;13(04):417–423
- 53 Buka SL, Shenassa ED, Niaura R. Elevated risk of tobacco dependence among offspring of mothers who smoked during pregnancy: a 30-year prospective study. *Am J Psychiatry* 2003;160(11):1978–1984