

Fig. 1 Several skeleton structures of heterocyclic compounds that are widely used in clinical setting.

With the character of high efficiency, specificity, and mildness, biocatalysis has been a hotspot in the field of green synthetic chemistry. Biocatalysts lipases are the most commonly used enzymes in synthetic organic chemistry, and favor the reaction with mild reaction conditions, as well as require low amounts of coenzymes.^{6,7} This article reviews the latest research in lipase-catalyzed synthesis of heterocyclic compounds, including five- and six-membered nitrogen-containing and oxygen-containing heterocyclic compounds or others synthesized in recent years, which may provide a valuable reference for green synthesis of heterocyclic compounds.

Lipase-Catalyzed Synthesis of Heterocyclic Compounds

Synthesis of Five-Membered Heterocyclic Compounds

Synthesis of Five-Membered Heterocyclic Compounds Containing Nitrogen

Pyrrole and its derivatives occur widely in nature and possess bioactivity in medicine field, such as captopril (an angiotensin-converting enzyme inhibitor for hypertension therapy),⁸ atorvastatin (a hydroxymethylglutaryl coenzyme A reductase inhibitor for blood lipid regulation),⁹ and the anticancer drug pyrotinib (an irreversible pan-ErbB inhibitor).¹⁰ In addition to the medicine field, the application of pyrrole compounds in materials science is also emerging, expanding and deepening gradually.¹¹

As we all know, the Paal–Knorr reaction represents an effective and simple method to synthesize pyrrole. In 2013, Zheng et al used enzymes to catalyze the Paal–Knorr reaction for the preparation of pyrrole derivatives (**►Scheme 1**).¹² They screened samples of commercially available enzymes for their ability to catalyze the model reaction of the Paal–Knorr reaction using 2,5-hexanedione and aniline as reaction materials. They found that lipase led to moderate yields






Fig. 2 Structures of indole natural products and pharmaceuticals.

(47 and 53% yields for Amano lipase M from *Mucor javanicus* and Lipase AT30, respectively) when compared with those of amylase and protease (60–99% yields).

Indole (benzopyrrole) is one of the important pyrrole derivatives with various bioactive activities, including antibacterial, anti-inflammatory, antitumor, antioxidation, and hypoglycemic activity, and has been extensively studied in the fields of medicine, materials science, food, fragrances, dyes, etc.^{13,14} **►Fig. 2** lists some indole derivatives that are clinically used as pharmaceuticals. In 2021, Li et al first reported *Candida rugosa* lipase (CRL)-catalyzed reaction of 1,3-diketones with fumaronitrile to synthesize polycyano-substituted indole with moderate to excellent yields (**►Scheme 2A**).¹⁵ This report also suggested that steric hindrance of R_1 and R_2 decreased the total yield of the target product. Encouragingly, when the substrate is an asymmetric 1,3-diketone, a higher regioselectivity and total yield of the reaction were achieved when compared with a chemical catalysis method. Unfortunately, in this reaction, R_1 and R_2 are limited to electron-donating alkyl groups, and the 1,3-dione compound should not be a β -ketoester (for example, diethyl malonate and methyl acetoacetate). This result suggested that choosing CRL as a catalysis favored a higher regioselectivity of the reaction, which made this enzymatic method more attractive than the chemical method.¹⁶ A plausible reaction mechanism for this process is depicted in **►Scheme 2B**: briefly, an enol anion is generated from acetyl acetone **1** under the action of the Asp-His dyad and stabilized by the oxyanion; the interaction between the enol



Scheme 1 Enzyme-catalyzed Paal–Knorr reaction for the synthesis of pyrrole compounds.




Scheme 2 (A) Lipase-catalyzed synthesis of cyano-containing multisubstituted indoles, and (B) a plausible reaction mechanism.

anion and the fumaronitrile **2** leads to the formation of carbanion, followed by attacking the carbonyl group to generate intermediate **4**, which is converted into 1,5-diketone enolate **5**; then **5** undergoes protonation/deprotonation process to produce intermediate **6**; subsequently, **6** reacts with another molecule of fumaronitrile via the Michael addition, and then undergoes cyclization, protonation/deprotonation, and aromatization to generate aniline intermediates **7**. The amino group in the structure of **7** attacks the carbonyl group in the molecule under the activation of lipase to form a five-membered heterocyclic intermediate **8**; finally, **8** undergoes dehydration and isomerization to produce the target product, polycyano-substituted indole **3**.

Xiang et al first suggested lipase-catalyzed the reaction between aldehydes and indole to obtain bisindole.¹⁷ This

process was conducted in a solvent mixture [1,4-dioxane containing 20% (v/v) water] at 50°C using *porcine pancreas* Type II (PPL) as a catalyst. The yield was highly associated with the electronic nature of aldehyde substituents: aromatic aldehydes with electron-withdrawing substituents showed higher reactivity than those with electron-donating groups, and aliphatic aldehydes. Our research group has improved this reaction by using the immobilized lipase TLIM (derived from *Thermomonas*) to catalyze this reaction in pure water. Excitingly, this method has a broad substrate tolerance, including various aldehydes such as aromatic aldehydes, heterocyclic aldehydes, and aliphatic aldehydes substituted by various electron-withdrawing groups, electron-donating groups, and large steric hindrance groups. The amount of enzyme (2 mg/mL) is much lower than that of



$R_1 = \text{H, Cl, CH}_3, \text{OCH}_3$

$R_2 = \text{H, CH}_3$

$R_3 = \text{H, CH}_3$

$R_4 = \text{H, C}_3\text{H}_7, \text{C}_6\text{H}_5$; 75%-89% yield

4-Cl-C₆H₄, 3-NO₂-C₆H₄, 4-NO₂-C₆H₄, 2-F-C₆H₄, 4-CN-C₆H₄;
85%-99% yield

4-CH₃-C₆H₄, 4-OCH₃-C₆H₄, 4-OH-3-OCH₃-C₆H₃, 2-OH-3-OCH₃-C₆H₃;
88%-95% yield


2-thienyl, 2-pyridyl, 4-*t*-C₄H₉-C₆H₄; 72%-95% yield

Scheme 3 Lipase-catalyzed synthesis of bis(indolyl)methanes.

ordinary enzymatic reactions, and after repeated use, the catalytic activity of enzymes is still good (►Scheme 3).¹⁸

Synthesis of Five-Membered Heterocyclic Compounds Containing Oxygen


Furan compounds have been widely used in the field of medicine. Currently, there are more than 10 furan drugs in clinical application.¹⁹ In 2021, Liu et al presented a novel synthesis method of polyhydroxyalkyl furans, which is based on the lipase-catalyzed Knoevenagel/oxo-Michael/Thorpe-Ziegler cascade reaction in pure water.²⁰ This method used reducing sugars and malononitrile as substrates, selected various lipases, such as lipase B from *Candida antarctica* (CALB), *Pseudomonas cepacia* lipase (PSL), Novozym435 (immobilized CALB), PPL as well as CRL, as catalysts, and acquired moderated to good yield (55–88%) (►Scheme 4A). It is worth to notify that Novozym435 showed the highest catalytic efficiency in comparison to other lipases. The authors proposed the catalytic mechanism as follows: first,




Scheme 5 Lipase-catalyzed synthesis of five-membered lactones.

the active methylene group in the malononitrile structure is deprotonated by the lipase-catalyzed triad and the oxygen anion hole; second, another substrate glucose and deprotonated malononitrile undergo the Knoevenagel condensation reaction to produce intermediate **1**; then, intermediate **1** rapidly undergoes the lipase-catalyzed oxo-Michael/Thorpe-Ziegler reaction to obtain intermediate **2**; finally, polyhydroxyalkyl furan is generated by spontaneous aromatization and ring opening reaction of intermediate **2** (►Scheme 4B).

The five-membered lactone ring is a special furan derivative structure. In 2019, Xu and Hu used cascade lipase CALB-catalyzed lactonization to achieve dynamic hemithioacetal formation, leading to efficient synthesis of five-membered lactone with chiral discrimination (►Scheme 5).²¹ Solvent-dependent regioselectivity was observed for the selective formation of 1,3-oxathiolan-5-one in weakly polar solvents, and γ -butyrolactone in the polar tetrahydrofuran (THF) solvent. Gutman et al realized that PPL could catalyze intramolecular transesterification of γ -hydroxy esters in the organic phase to obtain optically active γ -butyrolactones (►Scheme 6).²² Organic solvents with strong hydrophobicity are beneficial to the reaction. The initial reaction rate in *n*-hexane is two to four times those of solvents, such as ether, THF, and chloroform, yet, the reaction cannot proceed in an aqueous system. The study also suggested that the initial reaction rate of the γ -position of the substrate



Scheme 4 (A) Lipase-catalyzed synthesis of polyhydroxyalkyl furans, and (B) a plausible reaction mechanism.²⁰



- a: R₁ = H, R₂ = H
 b: R₁ = CH₃, R₂ = H
 c: R₁ = C₂H₅, R₂ = H
 d: R₁ = C₆H₁₃, R₂ = H
 e: R₁ = C₈H₁₇, R₂ = H
 f: R₁ = C₆H₅, R₂ = H
 g: R₁ = *p*-Me-C₆H₄, R₂ = H
 h: R₁ = *p*-MeO-C₆H₄, R₂ = H
 i: R₁ = *p*-Br-C₆H₄, R₂ = H
 j: R₁ = H, R₂ = CH₃
 k: R₁ = H, R₂ = Br
 l: R₁ = H, R₂ = CH₃CO

Scheme 6 Lipase-catalyzed synthesis of optically active γ -butyrolactones.

substituted by an alkyl group is about one order of magnitude faster than that substituted by an aryl group, and the configuration is just the opposite (the configuration of the alkyl-substituted lactone product is *S*, and the aryl-substituted the lactone product's configuration is *R*). Moreover, both the initial reaction rate and stereospecificity are significantly reduced when the substrate has a substituent at the α -position.

Synthesis of Six-Membered Heterocyclic Compounds

Synthesis of Six-Membered Heterocyclic Compounds Containing Nitrogen

Pyridine and its derivatives have a wide range of applications. They can be used as denaturants, dye assistants, photosensitizers, color developers, etc. Some of them are important components of vitamins, enzymes, and coenzymes.²³ Imidazo[1,2-*a*]pyridine is a valuable framework found in several nitrogen-containing heterocycles, and several drugs containing a pyridine core have been commercialized, such as zolpidem (a hypnotic drug), alpidem (anxiolytic drug), saripidem (anxiolytic drug), zolimidine (the antiulcer agent), as well as olprinone (a PDE-3 inhibitor for heart and circulatory failure therapy) (**Fig. 3**). Budhiraja et al dis-






Fig. 3 Examples of drugs containing the imidazo[1,2-*a*]pyridine framework.




Scheme 7 Lipase-catalyzed synthesis of imidazo[1,2-*a*]pyridine derivatives.

closed the CALB-catalyzed Groebke–Blackburn–Bienaymé (GBB) multicomponent reaction, which was used for the synthesis of imidazo[1,2-*a*]pyridine derivatives from 2-aminopyridine, aromatic benzaldehyde, and isocyanides (**Scheme 7**).²⁴ They suggested that the reaction was greatly influenced by the different substituents on the aryl benzaldehyde and 2-aminopyridine ring. When there was a strong electron-withdrawing group 5-NO₂ in the 2-aminopyridine ring structure, the reaction could not proceed. Molecular docking and molecular dynamics simulations showed that Thr40 and Ser105 residues in the CALB peptide chain played a crucial role in catalyzing the multicomponent reaction of GBB.

Quinoline (benzopyridine) is a research hotspot in nitrogen-containing six-membered heterocyclic benzo systems, and is widely used in medicine, agriculture, printing, dyeing and other fields.^{25,26} In 2012, Zheng et al reported a lipase-catalyzed reaction between 2-amino-3,5-dibromobenzaldehyde and cyclopentanone to synthesize quinoline derivatives (**Scheme 8A**).²⁷ The reaction could be catalyzed by PPL, pepsin, albumin from bovine, and other enzymes with the best catalytic effect being found in PPL (15 mg/mL). The yield can be achieved at 95%. The authors proposed the reaction mechanism as the following: lipase promotes the formation of enolate intermediates, which undergo aldol reaction with



Scheme 8 (A) Lipase-catalyzed synthesis of quinoline derivatives, and (B) a plausible reaction mechanism.



Scheme 9 Lipase-catalyzed synthesis of 4*H*-pyran derivatives with spiro structure.

the substrate 2-amino-3,5-dibromobenzaldehyde, and subsequently, intramolecular cyclization was performed through enzyme-assisted dehydration to generate the quinoline products (►Scheme 8B).

Synthesis of Six-Membered Heterocyclic Compounds Containing Oxygen

Pyran compounds are common six-membered heterocyclic compounds containing one oxygen heteroatom. There are two isomers of pyran: α -pyran (2*H*-pyran) and γ -pyran (4*H*-pyran) with neither of them being found naturally, but their derivatives are widely found in nature. Among the pyran derivatives, benzopyrans, namely chromones, have exhibited a broad range of physiological activities, such as antioxidant, anticancer, antibacterial, antiallergic and anti-human immunodeficiency virus (HIV) activities.^{28,29} Pyran compounds also exhibit inhibitory activity of cholinesterase,³⁰ and therefore can be used to treat neurodegenerative diseases such as Parkinson's disease, Alzheimer's disease, Huntington's disease, etc. Chai et al reported that PPL can efficiently catalyze the Knoevenagel condensation–Michael addition cascade reaction of isatin, malononitrile/ethyl cyanoacetate, and various carbonyl compounds in EtOH/H₂O solvent to produce 4*H*-pyran compounds with a spiro ring structure (►Scheme 9), and the yield can be achieved as high as 82 to 95%.³¹


In 2011, Xu et al first reported that PPL could efficiently catalyze the one-pot condensation of aldehyde, 1,3-cyclohexanedione, and malononitrile in a 10% (v/v) ethanol to produce 4*H*-pyridine in a yield of more than 90%.³² However, when the malononitrile was replaced by ethyl cyanoacetate, the yield would be significantly reduced. Based on the previous study, our research group further optimized the reaction by screening the catalytic activity of other lipases. We found that immobilized lipase TLIM could efficiently catalyze the condensation reaction of aromatic aldehyde/aliphatic aldehyde/heterocyclic aldehyde, 1,3-cyclohexanedione, and malononitrile/ethyl cyanoacetate, with a good to excellent yield (82–99%). When 1,3-cyclohexanedione was replaced by 4-hydroxycoumarin, the reaction also proceeded successfully (yield: 75–99%) (►Scheme 10A). Our data also showed that lipase TLIM had a broad spectrum of substrates, and the yield was still good after five repeated uses of the enzyme.³³ Compared with the traditional chemical methods, our reaction system offers great advantages in terms of temperature, catalyst reusability, and target product yield.^{34–36} The reaction mechanism is proposed as following: first, the Knoevenagel condensation

reaction of 4-chlorobenzaldehyde **1** and malononitrile **2** takes place to obtain intermediate product **4**; then, **4** combined with 4-hydroxycoumarin **3** via the Michael addition reaction to obtain intermediate **5**, which undergoes enolization, cyclization, and isomerization to produce the target product **6** (4*H*-pyran) (►Scheme 10).


In 2015, Yang et al suggested the PPL-catalyzed reaction of salicylaldehyde, 2,4-pentanedione, and aliphatic alcohol for the synthesis of 2*H*-chromene derivatives in aqueous DMF/H₂O solvent (►Scheme 11).³⁷ With the increase in carbon chain, the reaction rate and yield will be decreased, and when the carbon chain length is greater than that of pentanol, the reaction cannot proceed.

Xanthenes are also important six-membered heterocyclic compounds containing an oxygen heteroatom, which are sensitive to nucleophiles and light energy, so they are widely used in many fields such as medicine, materials science, etc.^{38,39} In 2018, our research group discovered a novel method for the synthesis of xanthenes: lipase TLIM-catalyzed Knoevenagel–Michael cascade reactions of 1,3-diketones with aromatic aldehydes/heterocyclic aldehydes/aliphatic aldehydes in *n*-hexane (►Scheme 12).⁴⁰ The yield can be up to 80 to 97%. Target products of this reaction were usually ring-opened xanthone compounds, yet, ring-closed xanthenes were formed when pyridine-2-carboxaldehyde/2,6-dichlorobenzaldehyde reacted with 1,3-cyclohexanedione under a high temperature or strong acid. It is worth mentioning that the structure of the target product may be different from that obtained by ordinary chemical methods when benzaldehyde (containing hydroxyl in the ortho position) reacted with 1,3-cyclohexanedione using this method. After three repeated uses of the lipase, the yield was significantly reduced.

Dicoumarol derivatives have a wide range of pharmacological and physiological activities. They can be used as a vitamin K antagonist to prevent blood clotting.⁴¹ They also have antibacterial activity and can be used as antifever medicine, etc.⁴² In 2019, Fu et al for the first time reported immobilized lipase RMIM (from *Rhizomucor miehei*)-catalyzed Knoevenagel–Michael cascade reaction, which was used to generate dicoumarol derivatives (►Scheme 13).⁴³ The method tolerates the reaction of 4-hydroxycoumarin with various aldehydes (aromatic aldehydes/heterocyclic aldehydes/aliphatic aldehydes) in a pure water with a yield of 81 to 98%. Circular dichroism spectroscopy revealed that the secondary structure of RMIM was changed after five times of repeated use of the enzyme. When the ratio of α -helix and β -fold was decreased, the enzyme activity was reduced, yet, the yield of the reaction was still 81%.



Scheme 10 (A) Lipase-catalyzed synthesis of 4*H*-pyran derivatives, and (B) a plausible reaction mechanism.



Scheme 11 Lipase-catalyzed synthesis of 2*H*-pyran derivatives.


Synthesis of Other Heterocyclic Compounds

Oxazolidinone derivatives are widely used in the treatment of acute bacterial skin infections and other severe infections induced by gram-positive bacteria and MRSA (methicillin-resistant *Staphylococcus aureus*).⁴⁴ In 2015, the synthesis of chiral oxazolidinone through lipase-catalyzed kinetic resolution was first described.⁴⁵ Lipase PS-IM (from *Burkholderia cepacia*) catalyzed the reaction of α -amino alcohol and bisacyl donor carbonate in *tert*-butyl methyl ether at room temperature (r.t.), and the chiral oxazolidinone can be obtained with a certain *ee*% value of up to 95% (→ **Scheme 14**). However, to achieve the complete conversion of selected enantiomer, a longer reaction time was necessarily needed. Fortunately, a catalytic amount of the Lewis acid SiO₂ can shorten the reaction time from 4 days to 3 hours with no

erosion on the enantioselectivity of the product. The reaction mechanism is proposed as follows: the first step was kinetics-controlled carbamylation, followed by lipase-catalyzed cyclization to obtain enantiomerically enriched oxazolidinone derivatives (→ **Scheme 14B**).

Lipase can be used to catalyze intramolecular transesterification of hemithioacetals to synthesize 1,3-oxathiolan-5-one derivatives (→ **Scheme 15**) with the best reactivity being found in lipase CALB or novozyme 435.⁴⁶ The reaction mechanism may be attributed to the following: first, under the catalysis of triethylamine, **1** (2-methylpropanal) and **2** (octanal) [or, **3** (pyridine-2-carbaldehyde)] react with mercaptan **A** (methyl 2-mercaptoacetate) or mercaptan **B** (methyl 3-mercaptoacetate) to produce γ - and δ -hydroxyl ester intermediates **1A-3A** and **1B-3B**, respectively. Subsequently, lipase-catalyzed intramolecular transesterification to selectively generate γ -lactone. δ -Hydroxy ester intermediates (**1B-3B**) did not react during the enzymatic catalysis. The study also found that the rigidity of the enzyme structure will be increased at low temperatures, which can improve substrate selectivity and product enantioselectivity.

1,3-Oxazines are one class of the most important heterocyclic compounds, which possess a variety of pharmacological



Scheme 12 Lipase-catalyzed synthesis of xanthone derivatives.



Scheme 13 Lipase-catalyzed synthesis of dicoumarol derivatives.


activities, such as bactericidal, antitumor, anti-HIV, anti-inflammatory, etc., and have been widely used in medicine.⁴⁷ Zhang et al reported a combined catalytic system using lipase CALB and Ru(bpy)₃Cl₂ for the reaction of naphthol and ethyl *N*-arylglycinate (instead of ethyl *N*-alkylglycinate) to produce 1,3-oxazine derivatives (►Scheme 16).⁴⁸ The reaction was conducted at r.t. in acetonitrile for 48 hours under a 12W fluorescent bulb (Philips), and the yield achieved was 19 to 69%. Ethyl *N*-arylglycinate with its electron-rich and electron-neutral nature showed higher reactive activity in comparison to that with electron-withdrawing nature. Mechanistically, the protocol consists of sequential enzymatic hydrolysis and visible-light-excited decarboxylation of ethyl *N*-aryl glycinate to generate α -amino radicals, which are oxidized to give iminium ions. The Mannich reaction with β -naphthols, transimination, and intramolecular cyclization furnished the 1,3-oxazine derivatives.

Heterospirocyclic compounds widely exist in various pharmacological agents and agricultural products, and can be used as anticancer, anticonvulsant, and antimicrobial agents.⁴⁹ Wang et al reported the synthesis of spiroox-


azine derivatives through immobilized CALB-initiated multicomponent reaction from nitrostyrene, cyclohexanone, *p*-nitrobenzaldehyde, and acetamide substrates, enabling six C-C/C-N bonds and two heterocycles in one-pot operation (►Scheme 17A).⁵⁰ When *p*-nitrobenzaldehyde was replaced by other aromatic aldehydes, the yield was very low. Substituted nitrostyrene with an oxygen group afforded better yields in comparison to other nitrostyrenes. Besides, during the reaction, acetamide could play a synergistic role with enzyme. Through isotope labeling and control experiments, a reasonable mechanism was proposed as the following: first, aldol reaction of *p*-nitrobenzaldehyde and cyclohexanone was catalyzed by CALB and acetamide to produce β -hydroxyketone (compound **A**) followed by condensation and hydrolysis in the presence of CALB to form enamine intermediate **C**, then **C** underwent Michael addition with nitrostyrene and the intramolecular Nef reaction to produce intermediate **D**, which was condensed with another molecule of cyclohexanone to form the target spiro product **E** (►Scheme 17B).

Conclusion and Perspectives


With the development of lipase versatility, lipases catalyze a wide range of reactions, especially for the synthesis of many heterocyclic compounds that are difficult to be prepared by classical chemical methods. Lipase may represent a promising strategy for the green synthesis of drugs and intermediates. However, lipases from natural sources have shortcomings, such as intolerance to high temperature, extreme pH, as well as poor stability in organic solvents, which limit its use in industry practice.



Scheme 14 (A) Lipase-catalyzed synthesis of chiral oxazolidinone derivatives, and (B) a plausible reaction mechanism.




Scheme 15 Lipase-catalyzed synthesis of 1,3-oxathiolan-5-one derivatives.



Scheme 16 Lipase-catalyzed synthesis of 1,3-oxazine derivatives.

X-ray crystallography makes the spatial structure of lipases readily available to scientists. With the development

of molecular biology, lipase can be improved through molecular biological methods, such as directed evolution, site-directed mutagenesis, chemical modification, immobilization, etc. to acquire stronger environmental tolerance, greater stability, as well as higher catalytic efficiency abilities. Furthermore, combined with the new technology of bioinformatics, the underlying mechanism of lipase-catalyzed reactions can be stated more clearly and in-depth. Last but not least, lipase-catalyzed strategies combined with traditional and novel organic synthesis methods may provide more options for the synthesis of target products, including chiral heterocyclic compounds.



Scheme 17 (A) CALB and acetamide-catalyzed synthesis of spirooxazine derivatives, and (B) a plausible reaction mechanism.

Funding

This work was financially supported by the National Natural Science Foundation of China (Grant No. 21676143) and National Students' Platform for Innovation and Entrepreneurship Training Program (Grant No. 202010291006Z).

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Mohammadi Ziarani G, Kheilkordi Z, Gholamzadeh P. Ultrasound-assisted synthesis of heterocyclic compounds. *Mol Divers* 2020; 24(03):771–820
- Dhameliya TM, Donga HA, Vaghela PV, et al. A decenary update on applications of metal nanoparticles (MNPs) in the synthesis of nitrogen- and oxygen-containing heterocyclic scaffolds. *RSC Advances* 2020;10(54):32740–32820
- Zhao Q, Meng G, Nolan SP, Szostak M. N-Heterocyclic carbene complexes in C-H activation reactions. *Chem Rev* 2020;120(04):1981–2048
- Mathur R, Negi KS, Shrivastava R, Nair R. Recent developments in the nanomaterial-catalyzed green synthesis of structurally diverse 1,4-dihydropyridines. *Rsc Adv* 2021;11(03):1376–1393
- Sadjadi S. Magnetic (poly) ionic liquids: a promising platform for green chemistry. *J Mol Liq* 2021;323:114994
- Barriuso J, Vaquero ME, Prieto A, Martínez MJ. Structural traits and catalytic versatility of the lipases from the *Candida rugosa*-like family: a review. *Biotechnol Adv* 2016;34(05):874–885
- Bilal M, Fernandes CD, Mehmood T, Nadeem F, Tabassam Q, Ferreira LFR. Immobilized lipases-based nano-biocatalytic systems - a versatile platform with incredible biotechnological potential. *Int J Biol Macromol* 2021;175:108–122
- Oz M, Lorke DE, Kabbani N. A comprehensive guide to the pharmacologic regulation of angiotensin converting enzyme 2 (ACE2), the SARS-CoV-2 entry receptor. *Pharmacol Ther* 2021; 221:107750
- Köhler-Forsberg O, Otte C, Gold SM, Østergaard SD. Statins in the treatment of depression: hype or hope? *Pharmacol Ther* 2020; 215:107625
- Li X, Yang C, Wan H, et al. Discovery and development of pyrotinib: a novel irreversible EGFR/HER2 dual tyrosine kinase inhibitor with favorable safety profiles for the treatment of breast cancer. *Eur J Pharm Sci* 2017;110:51–61
- Sorokin AB. Recent progress on exploring μ -oxo bridged binuclear porphyrinoid complexes in catalysis and material science. *Coord Chem Rev* 2019;389:141–160
- Zheng H, Shi Q, Du K, Mei Y, Zhang P. A novel enzyme-catalyzed synthesis of N-substituted pyrrole derivatives. *Mol Divers* 2013; 17(02):245–250
- Qin HL, Liu J, Fang WY, Ravindar L, Rakesh KP. Indole-based derivatives as potential antibacterial activity against methicillin-resistance *Staphylococcus aureus* (MRSA). *Eur J Med Chem* 2020;194:112245
- Iacopetta D, Catalano A, Ceramella J, et al. Synthesis, anticancer and antioxidant properties of new indole and pyranindole derivatives. *Bioorg Chem* 2020;105:104440
- Li F, Xu Y, Wang C, Wang C, Zhao R, Wang L. Efficient synthesis of cyano-containing multi-substituted indoles catalyzed by lipase. *Bioorg Chem* 2021;107:104583
- Pike RAS, Sapkota RR, Shrestha B, et al. K_2CO_3 -catalyzed synthesis of 2,5-dialkyl-4,6,7-tricyano-decorated indoles via carbon-carbon bond cleavage. *Org Lett* 2020;22(08):3268–3272
- Xiang Z, Liu Z, Chen X, Wu Q, Lin X. Biocatalysts for cascade reaction: porcine pancreas lipase (PPL)-catalyzed synthesis of bis(indolyl)alkanes. *Amino Acids* 2013;45(04):937–945
- Fu YJ, Lu ZP, Fang K, He XY, Xu HJ, Hu Y. Enzymatic approach to cascade synthesis of bis(indolyl)methanes in pure water. *RSC Adv* 2020;10(18):10848–10853
- Don-Lawson C, Nweneka DO, Okah R, Orié KJ. Synthesis, characterization and bioactivity of 1,1-bis(2-carbamoylguanidino)furran-2-ylmethane. *Am J Anal Chem* 2020;11(06):261
- Liu JX, Li FX, Zheng X, et al. Lipase-catalyzed synthesis of polyhydroxyalkyl furans from unprotected sugars and malononitrile. *Process Biochem* 2021;101:99–103
- Xu J, Hu L. Asymmetric one-pot synthesis of five- and six-membered lactones via dynamic covalent kinetic resolution: exploring the regio- and stereoselectivities of lipase. *Tetrahedron Lett* 2019;60(12):868–871
- Gutman AL, Zuobi K, Bravdo T. Lipase-catalyzed preparation of optically active gamma-butyrolactones in organic solvents. *J Org Chem* 1990;55(11):3546–3552

- 23 Shi YJ, Zhou Q, Wang Y, et al. Synthesis and biological activities of novel pyrazole amide derivatives containing substituted pyridyl group. *Chin J Org Chem* 2018;38(09):2450
- 24 Budhiraja M, Kondabala R, Ali A, Tyagi V. First biocatalytic Groebke-Blackburn-Bienaymé reaction to synthesize imidazo [1,2-*a*]pyridine derivatives using lipase enzyme. *Tetrahedron* 2020;76(47):131643
- 25 Stringer T, De Kock C, Guzgay H, et al. Mono- and multimeric ferrocene congeners of quinoline-based polyamines as potential antiparasitics. *Dalton Trans* 2016;45(34):13415–13426
- 26 Zablotskaya A, Segal I, Geronikaki A, Shestakova I, Nikolajeva V, Makarenkova G. N-Heterocyclic choline analogues based on 1,2,3,4-tetrahydro(iso)quinoline scaffold with anticancer and anti-infective dual action. *Pharmacol Rep* 2017;69(03):575–581
- 27 Zheng H, Liu J, Mei YJ, Shi QY, Zhang PF. A novel enzymatic synthesis of quinoline derivatives. *Catal Lett* 2012;142(05):573–577
- 28 Keri RS, Budagumpi S, Pai RK, Balakrishna RG. Chromones as a privileged scaffold in drug discovery: a review. *Eur J Med Chem* 2014;78:340–374
- 29 Semwal RB, Semwal DK, Combrinck S, Viljoen A. Health benefits of chromones: common ingredients of our daily diet. *Phytochem Rev* 2020;19(04):761–785
- 30 Sameem B, Saeedi M, Mahdavi M, et al. Synthesis, docking study and neuroprotective effects of some novel pyrano[3,2-*c*]chromene derivatives bearing morpholine/phenylpiperazine moiety. *Bioorg Med Chem* 2017;25(15):3980–3988
- 31 Chai SJ, Lai YF, Xu JC, Zheng H, Zhu Q, Zhang PF. One-pot synthesis of spirooxindole derivatives catalyzed by lipase in the presence of water. *Adv Synth Catal* 2011;353(2–3):371–375
- 32 Xu JC, Li WM, Zheng H, Lai YF, Zhang PF. One-pot synthesis of tetrahydrochromene derivatives catalyzed by lipase. *Tetrahedron* 2011;67(49):9582–9587
- 33 Fu Y, Lu Z, Ma X, et al. One-pot cascade synthesis of benzopyrans and dihydropyrano[*c*]chromenes catalyzed by lipase TLIM. *Bioorg Chem* 2020;99:103888
- 34 Gupta M, Gupta M, Rajnikant R, Gupta VK. Salicyldimine-based Schiff's complex of copper(II) as an efficient catalyst for the synthesis of nitrogen and oxygen heterocycles. *New J Chem* 2015;39(05):3578–3587
- 35 Bhosale RS, Magar CV, Solanke KS, Mane SB, Choudhary SS, Pawar RP. ChemInform abstract: molecular iodine: an efficient catalyst for the synthesis of tetrahydrobenzo[*b*]pyrans. *ChemInform* 2008;39(18):
- 36 Mohammadi AA, Asghariganjeh MR, Hadadzahmatkesh A. Synthesis of tetrahydrobenzo[*b*]pyran under catalysis of $\text{NH}_4\text{Al}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ (Alum). *Arab J Chem* 2017;10:S2213–S2216
- 37 Yang FJ, Wang Z, Wang HR, Zhang H, Yue H, Wang L. ChemInform abstract: enzyme catalytic promiscuity: lipase catalyzed synthesis of substituted 2*H*-chromenes by a three-component reaction. *ChemInform* 2015;46(05):25633–25636
- 38 Liu J, Zhang J, Wang H, et al. Synthesis of xanthone derivatives and studies on the inhibition against cancer cells growth and synergistic combinations of them. *Eur J Med Chem* 2017;133:50–61
- 39 Grzelakowska A, Kolinska J, Zaklos-Szyda M, Sokolowska J. Novel fluorescent fluorescent probes for L-cysteine based on the xanthone skeleton. *J Photochem Photobiol Chem* 2020;387:112153
- 40 Fu Y, Fan B, Chen H, Huang H, Hu Y. Promiscuous enzyme-catalyzed cascade reaction: Synthesis of xanthone derivatives. *Bioorg Chem* 2018;80:555–559
- 41 Sun C, Zhao W, Wang X, Sun Y, Chen X. A pharmacological review of dicoumarol: an old natural anticoagulant agent. *Pharmacol Res* 2020;160:105193
- 42 Li J, Sui YP, Xin JJ, et al. Synthesis of biscoumarin and dihydropyran derivatives with promising antitumor and antibacterial activities. *Bioorg Med Chem Lett* 2015;25(23):5520–5523
- 43 Fu Y, Lu Z, Fang K, He X, Huang H, Hu Y. Promiscuous enzyme-catalyzed cascade reaction in water: Synthesis of dicoumarol derivatives. *Bioorg Med Chem Lett* 2019;29(10):1236–1240
- 44 Kalam MA, Iqbal M, Alshememry A, Alkholief M, Alshamsan A. UPLC-MS/MS assay of Tedizolid in rabbit aqueous humor: application to ocular pharmacokinetic study. *J Chromatogr B Analyt Technol Biomed Life Sci* 2021;1171:122621
- 45 Zhang Y, Zhang Y, Ren Y, Ramström O. Synthesis of chiral oxazolidinone derivatives through lipase-catalyzed kinetic resolution. *J Mol Catal, B Enzym* 2015;122:29–34
- 46 Sakulsombat M, Zhang Y, Ramström O. Dynamic asymmetric hemithioacetal transformation by lipase-catalyzed γ -lactonization: in situ tandem formation of 1,3-oxathiolan-5-one derivatives. *Chemistry* 2012;18(20):6129–6132
- 47 Lathwal A, Mathew BP, Nath M. Syntheses, biological and material significance of dihydro[1,3]oxazine derivatives: an overview. *Curr Org Chem* 2021;25(01):133–174
- 48 Zhang GY, Xiang Y, Guan Z, He YH. Enzyme and photoredox sequential catalysis for the synthesis of 1,3-oxazine derivatives in one pot. *Catal Sci Technol* 2017;7(09):1937–1942
- 49 Yan ZY, Li JL, Ye GT, et al. Fused multicyclic polyketides with a two-spiro-carbon skeleton from mangrove-derived endophytic fungus *Epicoccum nigrum* SCNU-F0002. *RSC Advances* 2020;10(48):28560–28566
- 50 Wang JL, Chen XY, Wu Q, Lin XF. One-pot synthesis of spirooxazino derivatives via enzyme-initiated multicomponent reactions. *Adv Synth Catal* 2014;356(05):999–1005