
T. AMATOV, N. TSUJI, R. MAJI, L. SCHREYER, H. ZHOU, M. LEUTZSCH, B. LIST* (MAX-PLANCK-INSTITUT FÜR KOHLENFORSCHUNG, MÜLHEIM AN DER RUHR, GERMANY AND HOKKAIDO UNIVERSITY, SAPPORO, JAPAN)

Confinement-Controlled, either syn- or anti-Selective Catalytic Asymmetric Mukaiyama Aldolizations of Propionaldehyde

J. Am. Chem. Soc. 2021, 143, 14475–14481, DOI: 10.1021/jacs.1c07447.

Catalytic Asymmetric Aldolization of Propanal Enol Silanes



Significance: List and co-workers report an asymmetric Mukaiyama aldolization controlled by a confined chiral imidodiphosphorimidate catalyst. By using catalysts 1 and 2, syn- or anti-aldol products can be obtained from (E)- or (Z)-enolsilanes, respectively, in high yields and with high diastereo- and enantioselectivities.

Comment: Compared with the reported pioneering work (S. E. Denmark, S. K. Ghosh Angew. Chem. Int. Ed. 2001, 40, 4759), this method presents an improved, fully atom- and step-economic strategy. Ultimately, it could streamline the syntheses of complex oligopropionates.

SYNFACTS Contributors: Benjamin List, Yihang Li Synfacts 2021, 17(11), 1257 Published online: 19.10.2021 DOI: 10.1055/s-0041-1737044; Reg-No.: B08721SF

Category

Organo- and Biocatalysis

Key words

confinement asymmetric catalysis stereoselectivity Mukaiyama reaction

