


C. N. UNGAREAN, P. GALER, Y. ZHANG, K. S. LEE, J. M. NGAI, S. LEE, P. LIU\*, D. SARLAH\* (UNIVERSITY OF PITTSBURGH AND UNIVERSITY OF ILLINOIS, URBANA, USA)

Synthesis of (+)-Ribostamycin by Catalytic, Enantioselective Hydroamination of Benzene  
*Nat. Synth.* 2022, DOI: 10.1038/s44160-022-00080-x.

## Total Synthesis of (+)-Ribostamycin



**Significance:** Sarlah, Liu, and co-workers developed a rapid route to access (+)-ribostamycin, an aminoglycoside (AG) featuring a central 1,3-diaminocyclohexanetriol moiety. AGs are a well-known class of carbohydrate-derived antibiotics, which suppress protein synthesis by binding to prokaryotic rRNA.

**Comment:** The key step in the construction of (+)-ribostamycin is an initial enantioselective dearomative hydroamination of benzene. Deprotection of urazole **F** initiates an intramolecular epoxide opening forming an urazole-bridged heterobicyclic system **G**. After two glycosylation reactions, the urazole is fragmented and a global reduction releases the natural product without the need for further purification.

**SYNFACTS Contributors:** Erick M. Carreira, Henrik R. Wilke  
*Synfacts* 2022, 18(08), 0827 Published online: 19.07.2022  
DOI: 10.1055/s-0041-1738236; Reg-No.: C04122SF

© 2022, Thieme. All rights reserved.  
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of Natural Products and Potential Drugs

Key words

(+)-ribostamycin  
aminoglycosides  
dearomative hydroamination  
glycosylation  
urazole

Synfact  
of the  
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.