



Risk Factors Associated with Colorectal Cancer in Octogenarians Can Help Stratify the Need for Colonoscopy

Amy Y. Xiao¹  Sulakchanan Anandabaskaran¹  Maggie M. Ow^{1,2} 

¹Department of Medicine, University of Auckland, Auckland, New Zealand

²Department of Gastroenterology and Hepatology, Auckland City Hospital, Auckland, New Zealand

Address for correspondence Maggie M. Ow, MBChB, FRACP, MD, Department of Medicine, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand (e-mail: m.ow@auckland.ac.nz).

J Coloproctol 2022;42(2):146–151.

Abstract

Objective Colonoscopy is increasingly performed in octogenarians for the detection of colorectal cancer (CRC), but its benefits may be outweighed by its risks. The aim of the present study was to identify the risk factors for CRC in octogenarians presenting for colonoscopy to help stratify the need for this procedure.

Methods A retrospective analysis of 434 patients aged ≥ 80 years referred for a colonoscopy between January 2018 and December 2019. Comparisons were made between those with and without CRC and advanced adenoma (AA). The primary endpoint was to identify the clinical variables predictive of CRC and AA, and the secondary endpoints were complications and death 30 days after the procedure.

Results Colonoscopy was performed in 434 octogenarians, predominantly for symptoms, with CRC in 65 (15.0%) patients. Iron deficiency was associated with a higher risk of having CRC identified on colonoscopy (odds ratio [OR]: 2.33; 95% confidence interval [95%CI] = 1.36–4.00), but not symptoms such as bleeding, weight loss, or diarrhea. A colonoscopy in the last 10 years was protective, with a lower risk of CRC (OR: 0.45; 95% CI = 0.22–0.93). Patients with both normal iron stores and a colonoscopy within 10 years had a 92.5% chance of not having CRC. No variables were predictive of AA. Patients with complications, including death, were older and more likely to have underlying cardiorespiratory disease.

Conclusion Iron status and colonoscopy within 10 years can be used to predict the risk of CRC in octogenarians. Those with low predicted risk, especially if older and with cardiorespiratory disease, should be considered for non-invasive tests, such as computed tomography (CT) colonography, over colonoscopy.

Keywords

- ▶ colonoscopy
- ▶ colorectal cancer
- ▶ elderly
- ▶ risk factors

received
September 7, 2021
accepted after revision
October 25, 2021
published online
January 31, 2022

DOI <https://doi.org/10.1055/s-0041-1742256>.
ISSN 2237-9363.

© 2022. Sociedade Brasileira de Coloproctologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Introduction

Colorectal cancer (CRC) is the second leading cause of cancer-related deaths globally, with 1.9 million new cases a year worldwide.¹ The incidence of CRC increases with age, beginning around 40 years,² and, with longer life expectancy, the number of colonoscopies in the elderly is rising rapidly in the Western world. Colonoscopy is the investigation of choice for the evaluation of lower gastrointestinal (GI) symptoms and detection of colorectal pathology. The symptoms and cases of serious pathology increase with age; therefore, the yield of colonoscopy, especially for colorectal neoplasia, also rises with increasing age.³ A study conducted in the United Kingdom with almost 1,000 colonoscopies in predominantly symptomatic patients reported a diagnostic yield for CRC of 20% in octogenarians, compared with 7.4% in those younger than 80 years of age.⁴ The benefit of a higher diagnostic yield of colonoscopy in the elderly, however, is offset by a shorter life expectancy in this group,⁵ higher frequency of comorbidities,⁶ and increased risk of adverse events and hospitalisations postprocedure.^{7–10} It can also be technically more difficult to perform colonoscopy in the elderly, with lower completion rates and issues around inadequate bowel preparation.^{2,11}

Therefore, the decision to perform colonoscopy in patients of advanced age, especially the very elderly, requires careful consideration. Most studies^{12,13} on colonoscopy in the elderly have, to date, focused on diagnostic yield, technical factors, adverse events, and mortality. However, there is no data on who amongst the elderly is more likely to have an advanced colorectal neoplasm at the time of colonoscopy. This is an important consideration, especially in those who are frail or comorbid, as the benefit of colonoscopy may be outweighed by the potential risks if there is a low chance of an underlying CRC. The aim of the present study is to identify the risk factors associated with the presence of CRC in octogenarians undergoing colonoscopy.

Methods

We performed a retrospective study of all patients aged ≥ 80 years who underwent colonoscopy for any indication at our tertiary institution between January 2018 and December 2019. The institutional Ethics in Research Committee approved the study prior to commencement, and informed consent was obtained from the participants.

All endoscopy data are entered into an electronic database using software developed to report endoscopy (Provation MD, Provation, Minneapolis, MN, US). The data included indication, comorbidities, sedation, tolerance, quality of the bowel preparation, cecal intubation, endoscopic findings, therapeutics, withdrawal time, and complications. History of CRC or endoscopic procedures was also recorded.

The standard bowel preparation prior to colonoscopy at our unit was 3 L of polyethylene glycol, as a split-dose regimen. The option of having conscious sedation and the associated pros and cons were discussed with each patient, as part of the consent process. Conscious sedation was administered by the endoscopist using intravenous midazolam and fentanyl.

The level of tolerance was assessed by the endoscopist using the Modified Gloucester Comfort Score (MGCS): 1–none, talking comfortably throughout; 2–one or two episodes of mild discomfort, well tolerated; 3–more than 2 episodes of discomfort, adequately tolerated; 4–significant discomfort experienced several times during the procedure; and 5–extreme discomfort frequently during the test. The quality of the bowel preparation was graded by the endoscopist as excellent; good; adequate; fair; or poor. The excellent, good, and adequate grades were considered acceptable preparation. The proximal extent reached was documented by photography. A complete examination was defined as visualisation and/or intubation of the ileocecal valve (or ileocolonic anastomosis in patients with previous resection).

Study Endpoints

All patients were followed up until 30 days postcolonoscopy or death, whichever came first. The primary endpoint was to identify the clinical variables associated with the occurrence of CRC and advanced adenomas (AAs) on colonoscopy. Advanced adenomas were defined as those with a diameter ≥ 10 mm, with high-grade dysplasia or villous features on histology. The secondary endpoints were complications and death 30 days after the colonoscopy.

Statistical Analysis

The statistical analysis was performed using the Statistical Package for the Social Sciences (IBM SPSS Statistics for Windows, IBM Corp., Armonk, NY, US) software, version 26.0. The results were expressed as medians with interquartile ranges (IQRs), unless stated otherwise. The categorical data were compared using the Fisher exact test. Continuous data between patients with CRC/AA and patients without CRC/AA were compared using Mann-Whitney-U test. Univariate logistic regression analysis was performed to determine the variables associated with CRC. Multivariate logistic regression analysis was performed regarding the variables shown to be significant in the univariate analysis. For each variable, the odds ratio (OR) and 95% confidence interval (95%CI) were calculated to determine the risk associated with CRC. The levels of statistical significance were determined by two-tailed tests ($p < 0.05$).

Results

A total of 4,733 colonoscopies were performed between January 1st, 2018, and December 31st, 2019, 434 (9.2%) of which were in patients aged ≥ 80 years. The clinical indications for colonoscopy are shown in **Table 1**. The vast majority (97.9%; $n = 425/434$) of the patients were referred for symptomatic reasons. The most common indication was rectal bleeding. Prior to inclusion in the study, 32 patients had a history of colonic resection: 29 for previous CRC, 1 for Crohn disease, 1 for a large adenoma, and 1 for diverticular disease.

Of the 434 procedures, 399 (91.9%) were performed under conscious sedation, with both intravenous midazolam and fentanyl administered in 92.5% ($n = 369/399$) of sedated procedures, only midazolam in 1.8% ($n = 7/399$) of sedated

Table 1 Clinical indication for colonoscopy in 434 patients aged ≥ 80 years

Variable	Number of patients = 434
Median age, years (interquartile range)	84 (81–86)
Male gender – n (%)	221 (50.9)
Clinical indication – n (%) ^a	
Iron deficiency with/without anemia	145 (33.4)
Other anemia ^b	87 (20.0)
Rectal bleeding	156 (35.9)
Diarrhea	92 (21.2)
Other changes in bowel habits	63 (14.5)
Abdominal pain	71 (16.4)
Weight loss	81 (18.7)
Polyp surveillance	6 (1.4)
Postcolorectal cancer resection surveillance	3 (0.7)
Outpatient procedure – n (%)	252 (58.1)

Notes: ^aSome patients had more than one indication. ^bAnemia without evidence of iron deficiency.

procedures, and only fentanyl in 5.8% ($n = 23/399$) of sedated procedures. The median dose of midazolam was of 1.5 mg (IQR: 1.0–2.0 mg), and the median dose of fentanyl was of 75 mcg (IQR: 50–100 mcg). Among those who were sedated, 97.0% ($n = 387/399$) tolerated their colonoscopy very well (MGCS: ≤ 2). The unsedated group also tolerated the procedure very well, with all but 1 patient with MGCS ≤ 2 (97.1%; $n = 34/35$). The quality of the bowel preparation was acceptable in 59.2% ($n = 257/434$) of the procedures. The overall completion rate was of 90.3% ($n = 391/434$), and the adjusted completion rate (excluding obstructing lesions and poor preparation) was of 97.0% ($n = 421/434$). The median withdrawal time (in procedures without maneuvers) was of 8.2 minutes (IQR: 6.4–11.1 minutes).

Colorectal cancer and Advanced Adenoma

Of the 434 patients, 15.0% ($n = 65/434$) were diagnosed with CRC on colonoscopy. Three types of cancer other than CRC were detected: invasive prostate carcinoma, metastatic melanoma, and neuroendocrine carcinoma. A total of 15.7% ($n = 68/434$) of the patients had at least one AA: 12.2% ($n = 53/434$) had an adenoma with a diameter ≥ 10 mm; 9.9% ($n = 43/434$) had an adenoma with villous features; and 6.0% ($n = 26/434$) had an adenoma with high-grade dysplasia.

► **Table 2** shows a comparison of the demographics and clinical characteristics of the patients with and without CRC. The univariate analysis (► **Table 3**) identified age, iron deficiency, and previous colonoscopy within 10 years as signifi-

Table 2 Comparison of 434 patients aged ≥ 80 years stratified by the presence and absence of colorectal cancer on colonoscopy

Variable	With CRC; n = 65	Without CRC; n = 369	p-value
Median age, years (IQR)	85 (82–87)	83 (81–86)	0.048
Male gender	33 (50.8)	188 (50.9)	1.000
Previous CRC	4 (6.2)	25 (6.8)	1.000
Iron deficiency with/without anemia	33 (50.8)	112 (30.4)	0.002
Other anemia ^a	9 (13.8)	78 (21.1)	0.239
Rectal bleeding	25 (38.5)	131 (35.6)	0.675
Any changes in bowel habits	18 (27.7)	137 (37.1)	0.162
Diarrhea	11 (16.9)	81 (22.0)	0.414
Abdominal pain	12 (18.5)	59 (16.0)	0.589
Weight loss	10 (15.4)	71 (19.2)	0.604
Previous colonoscopy within 10 years	10 (15.4)	109 (29.5)	0.023

Abbreviations: CRC, colorectal cancer; IQR, interquartile range.

Notes: Values are expressed as numbers and percentages, unless stated otherwise. ^aAnemia without evidence of iron deficiency.

cant variables predicting CRC on colonoscopy. On the multivariate analysis (► **Table 3**), only iron deficiency and colonoscopy within 10 years were significant predictors of CRC. Iron deficiency was independently associated with an almost 2.5-fold increase in risk (OR: 2.33; 95%CI = 1.36–4.00; $p = 0.002$). In contrast, a previous colonoscopy within 10 years was associated with halving of the risk of CRC (OR: 0.45; 95%CI = 0.22–0.93; $p = 0.031$). Patients with both normal iron stores and a colonoscopy within 10 years had a 92.5% chance of not developing CRC.

However, none of the clinical variables in ► **Table 2** were associated with the presence of AAs.

Complications and Mortality

Patients who developed complications, including death, were significantly older and more likely to have underlying cardiorespiratory disease (► **Table 4**). Intraprocedural factors, such as patient tolerance, the sedation used, or the quality of the bowel preparation were not significant.

At 30 days, 19 (4.4%) patients developed a complication – there were 6 cardiac events, 4 cases of bleeding, 4 falls (with 2 fractures of the neck of the femur), 3 cases of pneumonia, 1 case of dehydration, 1 stroke, 1 urinary retention, and 1 perforation. In total, 1 patient had 3 complications, 7 (1.6%) died within 30 days of the colonoscopy – 2 following their operations for fracture of the neck of the femur, 1, from stroke, 1, from ischemic colitis (the presenting complaint), and 3, from their underlying cancers (1 colorectal, 1 gastric, 1 pancreatic).

Table 3 Univariate and multivariate analysis of colorectal cancer on colonoscopy in patients aged ≥ 80 years

Variable	Univariate analysis		Multivariate analysis	
	OR (95%CI)	p-value	OR (95%CI)	p-value
Age (years)	1.07 (1.00–1.15)	0.050	1.06 (0.98–1.14)	0.157
Male gender	0.99 (0.59–1.68)	0.979		
Previous colorectal cancer	0.90 (0.30–2.68)	0.853		
Iron deficiency with/without anemia	2.37 (1.39–4.04)	0.002	2.33 (1.36–4.00)	0.002
Other anemia ^a	1.61 (0.94–2.76)	0.084		
Rectal bleeding	1.14 (0.66–1.96)	0.647		
Any changes in bowel habit	0.60 (0.33–1.08)	0.091		
Diarrhea	0.72 (0.36–1.45)	0.362		
Abdominal pain	0.88 (0.41–1.88)	0.740		
Weight loss	0.68 (0.32–1.43)	0.303		
Previous colonoscopy within 10 years	0.43 (0.21–0.88)	0.021	0.45 (0.22–0.93)	0.031

Abbreviations: 95%CI, 95% confidence interval; OR, odds ratio.

Note: ^aAnemia without evidence of iron deficiency.

Table 4 Comparison of patients aged ≥ 80 years with and without complications, including death up to 30 days after the colonoscopy

Variable	No complications; n = 412	All complications, including death; n = 22	p-value
Median age, years (IQR)	83 (81–86)	86 (84–88)	0.007
Male gender	210 (51.0)	11 (50.0)	1.000
Outpatient procedure	243 (59.0)	9 (40.9)	0.120
Median dose of midazolam, mg (IQR)	1.5 (1.0–2.0)	1.3 (0.5–2.1)	0.996
Median dose of fentanyl, mcg (IQR)	75 (50–100)	75 (50–100)	0.342
Median patient tolerance, MGCS ^a (IQR)	1.0 (1.0–1.0)	1.0 (1.0–2.0)	0.406
Acceptable bowel preparation	244 ((59.2)	13 (59.1)	1.000
Comorbidities			
Cardiorespiratory disease	170 (41.3)	14 (63.6)	0.047
Diabetes	83 (20.1)	5 (22.7)	0.786
Stroke	53 (12.9)	3 (13.6)	1.000
Cognitive impairment/dementia	31 (7.5)	3 (13.6)	0.402
Renal replacement therapy	5 (1.2)	0 (0.0)	1.000
Other active (non-CRC) cancers	32 (7.8)	3 (13.6)	0.407

Abbreviations: CRC, colorectal cancer; IQR, interquartile range; MGCS, Modified Gloucester Comfort Score.

Notes: Values are expressed as numbers and percentages, unless stated otherwise. ^aSee the Methods section for the MGCS scoring criteria.

Discussion

Colonoscopy in the very elderly is a clinical decision that requires careful evaluation, with factors such as patient tolerance, risk of complications, and coexisting comorbidities contributing to this decision.¹⁴ The readily-available alternative, computed tomography (CT) colonography (CTC), is increasingly used in this population due to its easy preparation, better patient tolerance, and good safety profile, without any of the risks associated with sedation and invasive procedures.¹⁵ Nevertheless, serious pathology is prevalent in this age group,

and it is important to be able to identify the patients most likely to develop CRC, as this subgroup requires tissue diagnosis in a timely fashion, and it is best that they proceed straight to colonoscopy. To our knowledge, the present is the first study to report on the clinical predictors of CRC on colonoscopy in the elderly. Our cohort consists of patients who are extremely elderly, as this is an age group that is not often included in studies on colonoscopy but in whom management can be particularly challenging.

In our population of octogenarians, the vast majority underwent colonoscopy for symptomatic indications, and

the diagnostic yield for CRC was high, as expected: 15.0%.^{3,4,11,12,16-18} We found that patients with iron deficiency and those who did not undergo a colonoscopy in the last 10 years were significantly more likely to have a diagnosis of CRC on colonoscopy. Iron-deficiency anemia is recognised as a marker of occult GI malignancy. The prevalence of CRC in individuals with iron deficiency varies, with higher rates in men and postmenopausal women and in those with concurrent GI symptoms.¹⁹ The present study confirmed that iron deficiency is an important risk factor for CRC in the very elderly, and this is independent of any GI symptoms. Interestingly, none of the symptoms, such as bleeding or diarrhea, was independently associated with CRC. Rectal bleeding was the most common indication for colonoscopy in the present study, and is often described as a red flag for CRC. In our population, however, rectal bleeding was not predictive of CRC. This may be because hematochezia is often due to diverticular disease in this age group, which is a highly-prevalent condition, affecting over 60% of octogenarians,²⁰ and may not be specific enough as a risk factor for CRC.

In the present study, a colonoscopy in the last 10 years was protective and found to be a negative predictor of CRC: patients with a previous colonoscopy had half the risk of developing CRC compared to those who had not had a colonoscopy within 10 years. This is consistent with previous data from case-control and population studies documenting a reduced risk of CRC over a 10-year period in individuals who had a negative colonoscopy.²¹ In our population, around one third of patients had had a colonoscopy in the last 10 years, and this subgroup, especially if iron stores are normal as well, have a relatively low risk of having a diagnosis of CRC on colonoscopy, with 92.5% having only benign pathologies found. Therefore, it is better to submit this subgroup to a non-invasive test, such as a CTC, as their initial investigation, which is safer and better tolerated than colonoscopy, and with similar detection rates for large polyps and CRC.¹⁵

An important difference between the present study and many of the previous studies on colonoscopy in the very elderly is that our cohort comprised a large proportion of inpatients. Elderly inpatients are acutely unwell, may be more comorbid or deconditioned, and this may influence factors such as bowel preparation, procedure tolerance, and safety outcomes. In the present study, almost 40% of octogenarians had unacceptable bowel preparation, somewhat higher than the 15% to 25% reported in other studies.¹² Having said this, our cohort had a very respectable completion rate (unadjusted: 90%; adjusted: 97%), which is comparable to published data on this population,¹² despite issues with bowel preparation. Patient comfort during the procedure also rated well, with 97% reporting no more than 1 or 2 episodes of mild discomfort. With respect to complications, the risks of perforation, bleeding, and cardiorespiratory events in the present study were comparable to those of previous reports.^{22,23} Older patients and those with underlying cardiorespiratory disease were more likely to develop a complication, including death. We also observed a higher mortality rate (1.6%) in our cohort compared to those of other studies, in which the incidence of

mortality at 30 days is reported to be of approximately 1 to 5 per 1,000 colonoscopies.⁷ This increased mortality may be due to a higher proportion of inpatients in the present study, and further emphasizes the importance of careful evaluation when deciding in whom to perform a colonoscopy.

There are limitations to the present study. First, it was a retrospective study, and, as so, at risk of selection bias, as the patients studied may have been those who were more physically fit to undergo colonoscopy. Nevertheless, our data can help inform clinicians when making decisions on requesting colonoscopy in the elderly. Second, the cohort represents a sample from a tertiary centre, which introduces referral bias and limits the generalizability of the data to some extent. There is likely to be a greater representation of inpatients with complex medical problems requiring tertiary level care in our cohort. Thirdly, the quality of the bowel preparation was not measured using a validated quantitative scale such as the Boston Bowel Preparation score. Lastly, as the study was retrospective, only complications that required presentation to hospital or that were reported by the patient to the endoscopy unit would be recorded, and minor adverse events may not have been captured. Notwithstanding this, patients are given postprocedure instructions, which include a list of symptoms to look out for and recommendations to contact the endoscopy unit if these occur; hence, serious complications would have been accurately captured.

Conclusion

In summary, colonoscopy in octogenarians has a high diagnostic yield for CRC, and several clinical factors are useful to help stratify which patient has a greater risk of having a CRC at presentation, and therefore should proceed straight to colonoscopy. Octogenarians referred with iron deficiency, especially if they have not had a colonoscopy in the last 10 years, are at the highest risk of having a CRC, and should be strongly considered for colonoscopy. Conversely, those who have normal iron stores and have had a colonoscopy in the last 10 years should be considered for a non-invasive test, such as CTC, first, especially if they are older octogenarians and have underlying cardiorespiratory disease, as they have a low chance of having CRC and are more likely to suffer a complication after the colonoscopy. Colonoscopy is generally safe, with few adverse events, but does carry a risk of death, which is not inconsiderable. Therefore, prior to any decision to attempt colonoscopy in a very elderly patient, a careful assessment should include an evaluation of the aforementioned clinical predictors to determine the likelihood of detecting a CRC. Further prospective multi-centre studies are needed to validate the findings of the present study.

Disclosure Statement

This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Ferlay J, Ervik M, Lam F, et al. Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer Accessed February 2021 from: <https://gco.iarc.fr/today>
- 2 Lukens FJ, Loeb DS, Machicao VI, Achem SR, Picco MF. Colonoscopy in octogenarians: a prospective outpatient study. *Am J Gastroenterol* 2002;97(07):1722–1725
- 3 Stevens T, Burke CA. Colonoscopy screening in the elderly: when to stop? *Am J Gastroenterol* 2003;98(08):1881–1885
- 4 Arora A, Singh P. Colonoscopy in patients 80 years of age and older is safe, with high success rate and diagnostic yield. *Gastrointest Endosc* 2004;60(03):408–413
- 5 Wilson JA. Colon cancer screening in the elderly: when do we stop? *Trans Am Clin Climatol Assoc* 2010;121:94–103
- 6 Kahi CJ, Azzouz F, Juliar BE, Imperiale TF. Survival of elderly persons undergoing colonoscopy: implications for colorectal cancer screening and surveillance. *Gastrointest Endosc* 2007;66(03):544–550
- 7 Day LW, Kwon A, Inadomi JM, Walter LC, Somsouk M. Adverse events in older patients undergoing colonoscopy: a systematic review and meta-analysis. *Gastrointest Endosc* 2011;74(04):885–896
- 8 Costa-Moreira P, Silva M. Screening Colonoscopy in Older Patients: Elder Care or Elder Abuse? *Dig Dis Sci* 2020;65(07):1874–1876
- 9 Grossberg LB, Papamichael K, Leffler DA, Sawhney MS, Feuerstein JD. Patients over Age 75 Are at Increased Risk of Emergency Department Visit and Hospitalization Following Colonoscopy. *Dig Dis Sci* 2020;65(07):1964–1970
- 10 Tran AH, Man Ngor EW, Wu BU. Surveillance colonoscopy in elderly patients: a retrospective cohort study. *JAMA Intern Med* 2014;174(10):1675–1682
- 11 Cha JM, Kozarek RA, La Selva D, et al. Risks and benefits of colonoscopy in patients 90 years or older, compared with younger patients. *Clin Gastroenterol Hepatol* 2016;14(01):80–6.e1
- 12 Lin OS. Performing colonoscopy in elderly and very elderly patients: Risks, costs and benefits. *World J Gastrointest Endosc* 2014;6(06):220–226
- 13 Travis AC, Pievsky D, Saltzman JR. Endoscopy in the elderly. *Am J Gastroenterol* 2012;107(10):1495–1501, quiz 1494, 1502
- 14 Maratt JK, Calderwood AH. Colorectal cancer screening and surveillance colonoscopy in older adults. *Curr Treat Options Gastroenterol* 2019;17(02):292–302
- 15 Kim DH, Pickhardt PJ, Taylor AJ, et al. CT colonography versus colonoscopy for the detection of advanced neoplasia. *N Engl J Med* 2007;357(14):1403–1412
- 16 Virk GS, Jafri M, Ashley C. Colonoscopy and colorectal cancer rates among octogenarians and nonagenarians: nationwide study of US veterans. *Clin Interv Aging* 2019;14:609–614
- 17 Karajeh MA, Sanders DS, Hurlstone DP. Colonoscopy in elderly people is a safe procedure with a high diagnostic yield: a prospective comparative study of 2000 patients. *Endoscopy* 2006;38(03):226–230
- 18 Loffeld RJ, Liberov B, Dekkers PE. Yearly diagnostic yield of colonoscopy in patients age 80 years or older, with a special interest in colorectal cancer. *Geriatr Gerontol Int* 2012;12(02):298–303
- 19 Ioannou GN, Rockey DC, Bryson CL, Weiss NS. Iron deficiency and gastrointestinal malignancy: a population-based cohort study. *Am J Med* 2002;113(04):276–280
- 20 Lorenzo D, Gallois C, Lahmek P, et al; Groupe des Hémorragies Digestives Basses de l'ANGH (Association Nationale des Hépatogastroentérologues des Hôpitaux Généraux) Middle-term mortality and re-bleeding after initial diverticular bleeding: A nationwide study of 365 mostly elderly French patients. *United European Gastroenterol J* 2017;5(01):119–127
- 21 Singh H, Turner D, Xue L, Targownik LE, Bernstein CN. Risk of developing colorectal cancer following a negative colonoscopy examination: evidence for a 10-year interval between colonoscopies. *JAMA* 2006;295(20):2366–2373
- 22 Causada-Calo N, Bishay K, Albashir S, Al Mazroui A, Armstrong D. Association Between Age and Complications After Outpatient Colonoscopy. *JAMA Netw Open* 2020;3(06):e208958
- 23 Clarke GA, Jacobson BC, Hammett RJ, Carr-Locke DL. The indications, utilization and safety of gastrointestinal endoscopy in an extremely elderly patient cohort. *Endoscopy* 2001;33(07):580–584