

## Endoscopic resection of a giant esophageal fibrovascular polyp


A 66-year-old woman consulted because of dysphagia and occasional regurgitation of a lump of tissue during episodes of vomiting over the previous 9 months. A computed tomography (CT) scan showed a giant esophageal mass emerging in the upper esophagus. Evaluation by endoscopic ultrasound (EUS) showed that the mass was very hypoechoic, and it was suspected to be lipoma.

An endoscopic procedure was performed with the patient intubated, using a large-channel gastroscope (Pentax, Japan) and CO<sub>2</sub> insufflation. The endoscopy showed a polyp with a diameter of 5 cm at its base and a length of 15 cm (Fig. 1a). The mucosal appearance was normal and its lower part was ulcerated.

Resection was commenced with the submucosal injection of saline mixed with adrenaline (1:10000) at the base of the

polyp. The polyp was gradually dissected step by step at its base, using first a triangle-tip electro-surgical knife (Fig. 1b,c) then a hook knife (Olympus, Japan) (Fig. 1d; Video 1) using the Endocut current. Hemostasis was achieved using Coagrasper forceps (Olympus). The polyp was then caught with a snare (Fig. 1e) and extracted using an over-tube (Fig. 2). No complications such as bleeding or perforation occurred. Histological analysis confirmed the fibrovascular nature of the polyp, which had been completely resected.

Even though fibrovascular polyps are rare, they may cause morbidity and mortality, which makes their resection mandatory [1,2]. In such situations, two types of



**Fig. 1** Endoscopic views showing: **a** a fibrovascular polyp with its base implanted just under the Killian area; **b** the initial stages of dissection using a triangle-tip electro-surgical knife; **c** the ongoing dissection; **d** completion of the dissection using a hook knife; **e** the polyp being caught with a snare.



**Fig. 2** Macroscopic appearance of the resected polyp.

### Video 1



Endoscopic resection of a giant fibrovascular polyp using first a triangle-tip knife and then a hook knife.

therapeutic approach are available: surgical or endoscopic [1]. Because of the risk of uncontrolled bleeding, it is conventionally recommended that polyps larger than 80 mm are not treated endoscopically [1, 3]. Esophageal surgery is however known to be complex, associated with morbidity, and not always technically feasible.

By using a dissection knife, we were able to safely remove this large polyp endoscopically. This is a new endoscopic approach, as in all the previously published cases, fibrovascular polyps have been removed with a polypectomy snare [3–5]. In conclusion, endoscopic treatment of a giant fibrovascular polyp can be performed safely and effectively using appropriate equipment.

Endoscopy\_UCTN\_Code\_TTT\_1AO\_2AG

**Competing interests:** None

**Diane Lorenzo, Jean Michel Gonzalez, Marc Barthet**

Service de Gastroentérologie, Hôpital Nord, Marseille, France

#### References

- 1 Lee SY. Recurrent giant fibrovascular polyp of the esophagus. *World J Gastroenterol* 2009; 15: 3697
- 2 Mishra PK, Goel N, Saluja SS et al. Management of giant fibrovascular polyp of esophagus. *Am Surg* 2012; 78: E538–E540
- 3 Park J-S, Bang BW, Shin J et al. A case of esophageal fibrovascular polyp that induced asphyxia during sleep. *Clin Endosc* 2014; 47: 101
- 4 Murino A, Eisendrath P, Blero D et al. A giant fibrovascular esophageal polyp endoscopically resected using 2 gastroscopes simultaneously (with videos). *Gastrointest Endosc* 2014; 79: 834–835
- 5 Zhang J. Successful endoscopic removal of a giant upper esophageal inflammatory fibrous polyp. *World J Gastroenterol* 2009; 15: 5236

#### Bibliography

**DOI** <http://dx.doi.org/10.1055/s-0042-101857>  
Endoscopy 2016; 48: E69–E70  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

#### Corresponding author

**Diane Lorenzo, MD**  
Hôpital Nord  
chemin des Bourrelly  
Marseille  
France  
Fax: +33-4-91961311  
[diane.lorenzo@gmail.com](mailto:diane.lorenzo@gmail.com)