


Cushing's syndrome managed by endoscopic ultrasound-guided radiofrequency ablation of adrenal gland adenoma

A 38-year-old woman presented with “moon face,” “buffalo hump,” and weight gain of 9 kg in 12 months. Overnight, 1 mg dexamethasone failed to suppress the morning level of cortisol, and the 24-hour urine cortisol level was elevated to 101 µg/day (normal range 0–50). Initial contrast-enhanced abdominal computed tomography (CT) showed a 2.8-cm left adrenal mass enhanced in arterial phase, and the patient was diagnosed with Cushing's syndrome due to left adrenal adenoma (► **Fig. 1 a**). She refused surgical treatment but agreed to undergo endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA; STARmed, Koyang, Korea) (► **Video 1**).

Prior to RFA, contrast-enhanced EUS with SonoVue (Bracco, Inc., Milan, Italy) was performed. Findings of early enhancement and delayed washout were compatible with adrenal adenoma (► **Fig. 2 a**). A 19-gauge needle electrode was positioned inside the adenoma. Using real-time EUS imaging, RFA (50 W) was performed at five different sites (► **Fig. 2 b**). Four days later contrast-enhanced EUS revealed viable tissue remaining at the marginal edge of the previously ablated portion of the adenoma (► **Fig. 3 a**). EUS-RFA was repeated at five more sites in the remaining viable tissue (► **Fig. 3 b**). Follow-up CT at 1 week showed the adrenal mass almost completely replaced with necrotic tissue, without complications (► **Fig. 1 b**). Serum and urine cortisol levels returned to normal the following day and remained normal for the next 2 months, with no adverse events related to RFA. However after the third month, the cortisol levels were re-elevated and this time the patient agreed to surgery.

Until recently, there were only a few case reports of RFA for the treatment of Cushing's syndrome; all of them were treated via the CT-guided percutaneous method [1,2]. The present case is the first in



► **Fig. 1** Computed tomography scan showing a 2.8-cm left adrenal adenoma with arterial enhancement in the coronal view (arrows). **a** Before endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA). **b** After EUS-RFA.

which EUS-RFA was used to manage Cushing's syndrome caused by adrenal adenoma. This case report supports EUS-RFA as a safe and feasible alternative method that should be considered in patients who refuse surgical treatment. Further evidence and experiences are required.

Endoscopy_UCTN_Code_TTT_1AS_2AD

Competing interests

None

The Authors


Sun-Ho Lee, Dong Wan Seo, Dongwook Oh, Tae Jun Song, Do Hyun Park, Sang Soo Lee, Myung-Hwan Kim

Division of Gastroenterology, Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea


VIDEO 1



► **Video 1:** Endoscopic ultrasound-guided radiofrequency ablation performed on a left adrenal adenoma for the management of Cushing's syndrome.



► **Fig. 2** Endoscopic ultrasound-guided radiofrequency ablation (EUS-RFA) of the left adrenal adenoma. **a** Contrast-enhanced EUS with early enhancement. **b** The EUS-RFA needle positioned inside the adenoma.



► **Fig. 3** Contrast-enhanced endoscopic ultrasound (EUS) 4 days after the first EUS-guided radiofrequency ablation (RFA) treatment. **a** Central hypo-echogenicity with enhancement remained at the marginal edge of the adenoma. **b** Repeat EUS-RFA was performed.

Corresponding author

Dong Wan Seo, MD, PhD

Division of Gastroenterology, Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, 88, Olympic-Ro 43-Gil, Songpa-gu, Seoul 138-736, Korea,
Fax: +82-2-4855782,
dwseoamc@amc.seoul.kr

References

- [1] Nishi N, Tanaka J, Minagawa A. Cushing syndrome treated by radiofrequency ablation of adrenal gland adenoma. *Jpn J Radiol* 2012; 30: 274–276
- [2] Nunes TF, Szejnfeld D, Xavier AC et al. Percutaneous ablation of functioning adrenal adenoma: a report on 11 cases and a review of the literature. *Abdom Imaging* 2013; 38: 1130–1135

Bibliography

DOI <http://dx.doi.org/10.1055/s-0042-118165>
Endoscopy 2017; 49: E1–E2
© Georg Thieme Verlag KG Stuttgart · New York
ISSN 0013-726X