Barriers and Benefits of Information Communication Technologies Used by Health Care Aides

Hector Perez¹ Noelannah Neubauer¹ Samantha Marshall¹ Serrina Philip¹ Antonio Miguel-Cruz^{1,2,3} Lili Liu¹

¹School of Publich Health Sciences, Faculty of Health, University of Waterloo, Waterloo (ON), Canada

²Glenrose Rehabilitation Hospital, Edmonton (AB), Canada

³Department of Occupational Therapy, Faculty of Rehabilitation

Medicine, University of Alberta, Edmonton (AB), Canada

Address for correspondence Lili Liu, PhD, University of Waterloo, 200 University Avenue West, Waterloo, ON N2L 3G5, Canada (e-mail: lili.liu@uwaterloo.ca).

Appl Clin Inform 2022;13:270-286.

Abstract	Packaround Although information and communication technologies (ICT) are be
Abstract	background Although information and communication technologies (icr) are be-
	coming more common among nealth care providers, there is little evidence on now ICT
	can support health care aides. Health care aides, also known as personal care workers,
	are unlicensed service providers who encompass the second largest workforce, next to
	nurses, that provide care to older adults in Canada.
	Objective The purpose of this literature review is to examine the range and extent of
	barriers and benefits of ICT used by health care workers to manage and coordinate the
	care-delivery workflow for their clients.
	Methods We conducted a literature review to examine the range and extent of ICT
	used by health care aides to manage and coordinate their care delivery, workflow, and
	activities. We identified 8.958 studies of which 40 were included for descriptive
	analyses.
	Results We distinguished the following five different purposes for the use and
	implementation of ICT by health care aides: (1) improve everyday work. (2) access
	electronic health records for home care (3) facilitate client assessment and care
	planning (4) enhance communication and (5) provide care remotely. We identified
	128 barriers and 130 benefits related to adopting ICT. Most of the barriers referred to
	incomplete bardware and software features time consuming ICT adeption heavy or
	incomplete hardware and software features, time-consuming fer adoption, neavy of
Kannanda	increased workloads, perceived lack of userulitiess of ict, cost of budget restrictions,
Keywords	security and privacy concerns, and lack of integration with technologies. The benefits
Information	for nealth care aldes' adoption of ICT were improvements in communication, support
technology	to workflows and processes, improvements in resource planning and health care aides'
delivery of health care	services, and improvements in access to information and documentation.
health personnel	Conclusion Health care aides are an essential part of the health care system. They
 workflow 	provide one-on-one care to their clients in everyday tasks. Despite the scarce informa-

aged

hey mation related to health care aides, we identified many benefits of ICT adoption.

received September 7, 2021 accepted after revision January 3, 2022

© 2022. Thieme. All rights reserved. Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

DOI https://doi.org/ 10.1055/s-0042-1743238. ISSN 1869-0327.

Background and Significance

The world's aging population is growing at a significant rate.¹ Globally, a higher increase is noted among individuals aged 65 years and older compared with other age groups.² In addition, with advancing age, the risk of developing chronic diseases, such as depression and dementia, becomes higher.³ All of these combined represent significant challenges for health care systems and providers.

In Canada, health care providers, such as health care aides and personal support workers, provide direct care for the elderly, ill, or persons who require assistance with their activities of daily living and multiple tasks in different settings.^{4–7} These tasks may include self-care, medication management, social interactions,^{8,9} emotional support, meal preparation, companionship, socialization, and housekeeping,¹⁰ all while under the supervision of regulated nursing staff or other health professionals.^{11,12}

Although health care aides are unlicensed support personnel, they comprise the second largest workforce within the health care sector,¹³ next to nurses, providing 80% of the direct care to Canadian seniors.^{8,9} The situation regarding professional regulation for this group of health care workers is similar between the United States,¹⁴ Canada,¹⁵ and Europe, such that the regulation may vary from one country to another.¹⁶ Depending on the location, this workforce is recognized by different names,⁸ including personal support workers, personal care attendants, long-term care aides, resident care workers, nursing aides, nursing assistants, health care assistants, and care aides.^{8,9,11,16–18} For this study, we will refer to them as health care aides.

The population of health care aides is difficult to estimate.¹⁹ The exact size of the health care aide workforce in Canada is unknown since many are on a contractually, parttime or on-call basis. Most of these workers are women.^{18,20,21} The shortage of health care aides in Canada and several high-income countries around the world will continue to grow as the world's senior population is expected to increase in the upcoming years.^{8,16,22–26} The necessity for more health care providers is a concern, not unique to Canada. Health care aides are an essential component for the health care sector.^{8,16,27,28}

These workers are vital for high quality care in continuing care facilities,^{12,29} including home care, supportive living, and long-term care.^{11,12,29} However, they do not work under a standardized scope of practice.^{7,18} Instead, their duties are assigned depending on the employer, work setting, and client needs.^{5,9–11}

Information and communication technologies (ICT) has created numerous avenues for advocating and promoting health care delivery among various age groups. ICT represent an enormous opportunity to improve health care by providing quality, accessibility, and affordability for multiple users.^{30–32} Current technologies within the scope of practice of nurses³³ and unlicensed health care aides, include telemonitoring and care delivering mobile application.^{15,34,35}

ICT implementation and adoption are key and beneficial to assist health care aides' workflows and activities.^{36,37}

Existing supporting health care tools can potentially provide a range of applications and platforms to facilitate health care aides' assignments,³⁸ transmit observations, and enhance communication between families, clients, and health care providers within the ecosystem of care.³⁹ ICT can help to deliver real-time support and keep clients' information confidential and available,^{40,41} and ultimately improve caring activities and health care delivery.³⁷

Despite the various technology-based interventions that exist, none have been designed with health care aides in mind or intended to support their workflows.³⁷ For example, Steele Gray et al⁴² found that ICT for health care aides are inefficient and often do not fit their workflows. Also, Sterling et al³⁷ described how health care aides consistently identified existing ICT as outdated and ineffective; and not convenient for their needs, they suggested that current technologies exacerbate health care aides' existing communication challenges by not allowing real-time communication with different health care provides.

The existing evidence suggests that health care aides face multiple challenges to access and share patient information^{15,37} and communicate with other health care providers.⁴³ This has a negative impact on their ability to care for their clients.¹⁵ Saari et al¹⁵ identified technology as having the ability to facilitate effective communication and address information gaps for health care aides. Consequently, opportunities for ICT intended for health care aides include features of portability,⁴⁴ ubiquity,⁴⁵ interoperability,⁴⁶ communication tools to assist care plans,⁴³ and integration with remote care, telemonitoring, and assistive technologies.³⁶

To our knowledge, no comprehensive review has studied the range and extent of ICT used by health care aides to support their workflows and practice. Likewise, it is important to understand the barriers and benefits of implementation and adoption of ICT by health care aides. This understanding can inform the design, implementation, and adoption of future technologies and tools, since the role of health care aides is becoming more important, as they provide direct care to clients.

Objectives

To date, there is little evidence on how ICT can support health care aides' workflow and practice as the barriers and benefits of ICT use and adoption remain unknown. The purpose of this literature review is to examine the range and extent of barriers and benefits of ICT used by health care aides to manage and coordinate the care-delivery workflow for their clients.

Methods

We conducted a literature review based on Daudt et al,⁴⁷ modifications of the methodology by Arksey and O'Malley.^{47,48} This methodology followed six steps: (1) determine the research questions following the PICOS (population, intervention, comparison, outcome, and study design)

framework, (2) identify the relevant studies, (3) study selection, (4) chart data, (5) collect and summarize, and (6) report the results. Modification made by Daudt et al⁴⁷ of this methodology includes an interprofessional team in step (2) and use a three-tiered approach to cross-check and select the articles in step (3).^{47,49}

Data Sources and Search Strategy

We examined peer-reviewed literature published between January 2010 and March 2020, considering the past 10 years of technology development. We searched three academic databases: Medline, EMBASE, and CINAHL. We defined the search strategy in collaboration with an expert subject expert librarian. Then, we formulated and followed two main concepts to extract relevant studies from the electronic databases (N.N. and S.M.):

- 1. Health care professionals providing care to people who require health-related personal care.
- 2. Use of technologies within care.

We combined keywords for the above concepts, thesaurus terms including subject headings and the Medical Subject Headings (MeSH) terms, and free-text words using the Boolean operators of "AND" and "OR" (N.N., S.M., and A.M. C.). In addition, we customized search terms, strategies, and limits for each database (N.N., S.M., and A.M.C). For more details about the search strategy, see **Supplementary Table S1** (available in the online version).

Study Selection Process

We exported all studies to EndNote, a reference management software, and removed duplicates (N.N., S.M., and H.P.). Then, we transferred the remaining studies to Covidence, a primary screening and data extraction tool (N.N., S.M., and H.P.). Following, we removed additional duplicates using Covidence or manually as we progressed in the review process (N.N., S.M., H.P., and S.P.). Then, two researchers engaged in the title and abstract screening independently (N.N. and S.M.) where we compared studies with the inclusion and exclusion criteria. We resolved conflicts between the researchers through a discussion where consensus was reached (N.N., S.M., A.M.C., S.P., and H.P.). If a disagreement persisted after the discussion, one senior researcher or domain expert researcher resolved the conflict (N.N., A.M.C., and L.L.).

Next, two researchers assessed the full texts (N.N. and S. M.). A third rater decided if those studies should be included in the data extraction phase if disagreement occurred (A.M. C.). All researchers were thoroughly calibrated and trained on applying the inclusion and exclusion criteria before evaluating studies.

Inclusion Criteria

We included papers that:

- Examined the use of ICT primarily by healthcare workers, including health care aides and home care nurses.
- Reported ICT intended to assist in providing caregiving services at home.

- Were published and available in full text in peer-reviewed journals, doctoral or master's theses, and conference proceedings.
- Were published in English.

Exclusion Criteria

We excluded papers that:

- Did not report the use of ICT.
- Were related to care providers with a professional designation or regulated scope of practice.
- Reported ICT that were not intended for caregiving services at home.
- Described primary ICT use by someone other than health care aides or home care nurses.
- Did not provide enough information to categorize or extract information from.
- Were beyond the scope of this literature review.
- Were not available in full text or published in peerreviewed journals, doctoral or master's theses, or conference proceedings.

Data Extraction Process

Three researchers (N.N., S.M., and A.M.C.) extracted data from the selected studies in an Excel spreadsheet file where we operationalized the variables. We reviewed each study in full and extracted data for the areas of interest for this review (N.N., S.M., S.P., H.P., and A.M.C.). Finally, two researchers met regularly during the study extraction phase to discuss differences and reach a consensus (S.M. and N.N.).

Data Analysis

Four researchers conducted the data analysis (S.M., N.N., A.M. C., and H.P.). Due to the diversity of the included articles, we decided to use a qualitative approach, and conducted content analysis on the extracted data. We categorized the studies into the following five main groups according to the purpose of each study. In addition, we calculated descriptive statistics for population, study characteristics, ICT features, barriers, and benefits of ICT. Finally, after several rounds of discussions between the research team, we classified the barriers and benefits that were identified in this review. In **– Table 1**, we present a detail description of the selected studies including characteristics and ICT features.

Results

Study Selection

Fig. 1 shows the scholarly literature article search results. In the initial search, we identified 8,958 studies from academic databases. After the removal of 1,065 duplicates, 7,893 publications entered in the selection process. The agreement level between raters⁵⁰ during the title and abstract phase and the full-text screening phase was high, with a 95.57% level of agreement for the abstracts and 84.71% for the full papers.

Most selected studies were conducted in the United States (30%, 12/40), Sweden (12.5%, 5/40), and Norway (12.5%, 5/40). The remaining studies were conducted across nine

study purpose
by
classified
studies
selected
of
1 Characteristics
Table '

Study purpose	Authors	Age of participants (range)	Gender of participants	Study design	Data collection method	Hardware used	Software used
Improve everyday work	Stroulia et al ⁵⁴	Not reported	Not reported	Multiple methods	Interviews, focus groups, survey Questionnaire	Tablet	Application
Improve everyday work	Sassen et al ⁵⁵	Intervention group mean: 38.6, con- trol group mean: 39.7	Both genders (mostly female)	Quantitative	Survey Questionnaire	Laptop	Web based
Improve everyday work	Nilsson and Fagerström ⁵⁶	Not reported	Not reported	Qualitative	Focus groups	Not reported	Not reported
Improve everyday work	Lexis et al ⁶¹	Not reported	Female	Quantitative	Survey Questionnaire	Desk computer	Web based
Improve everyday work	Kaunda-Khangamwa et al ⁵²	Not reported	Not reported	Qualitative	Interviews	Phone	Text messaging
Improve everyday work	Gund et al ⁵⁷	38-56; mean=48	Not reported	Multiple methods	Interviews	Desk computer	Web based
Improve everyday work	Gars and Skov ⁵³	Not reported	Not reported	Qualitative	Focus groups, observations	Tablet	Application
Improve everyday work	Danilovich et al ⁶⁰	Not reported	Female	Qualitative	Interviews, Focus groups	Tablet	Application
Improve everyday work	Brown et al ⁵¹	Mean of case managers=36.4	Both genders	Multiple methods	Focus groups, survey Questionnaire	Tablet and computer	Web based and application
Improve everyday work	Andersen et al ⁵⁸	High strain mean: 47.9, Iow and moderate strain mean: 37.9	Both genders (90% female)	Quantitative	Survey Questionnaire	Phone	Web based
Improve everyday work	Alhuwail and Koru ⁵⁹	30->51	Not reported	Qualitative	Interviews, Focus groups	Tablet	Application
Improve everyday work	Alhuwail and Koru ⁴⁶	30- > 52	Not reported	Qualitative	Interviews, Focus groups, observations	Not reported	Not reported
Access electronic health records for home care	Yang et al ⁶³	25-47	Both genders (mostly female)	Qualitative	Observations	Laptop	Web based
Access electronic health records for home care	Westra et al ⁶²	Not reported	Not reported	Quantitative	Not reported	Not reported	Web based
Access electronic health records for home care	Tapper et al ⁶⁴	Not reported	Not reported	Quantitative	Survey Questionnaire	Tablet	Web based
Access electronic health records for home care	Sockolow et al ⁶⁵	Median age = 49 years	Both genders (88% female)	Mixed methods	Retrospective	Laptop	Application
							(Continued)

(Continued)
1
Ð
P
P

Software used	Not reported	Web based	Application and web based	Not reported	Web based	Application	Web based	Web based	Web based	Web based	Web based				
Hardware used	Not reported	Desk computer	Desk computer	Desk computer	Desk computer	Tablet	Phone and tablet	Not reported	Phone and tablet	Tablet	Tablet	Phone	Desk computer	Desk computer	Desk computer
Data collection method	Interviews	Secondary analysis	Interviews, focus groups	Not applicable	Interviews, Focus groups, observations	Survey Questionnaire	Focus groups, obser- vations, survey Questionnaire	Survey Questionnaire	Interviews, focus groups	Interviews	Focus groups	Secondary analysis	Not applicable	Interviews, observations	Focus groups, observations
Study design	Qualitative	Quantitative	Qualitative	Not applicable	Quantitative	Quantitative	Multiple methods	Multiple methods	Qualitative	Qualitative	Qualitative	Quantitative	Not applicable	Qualitative	Qualitative
Gender of participants	Not reported	Female	Not reported	Not reported	Female	Both genders (mostly female)	Both genders (mostly female)	Not reported	Not reported	Not reported	Both genders (mostly female)				
Age of participants (range)	Not reported	Not reported	Not reported	Not reported	30–51	28–63	Not reported	Not reported	28–59; mean=47	23-65; mean=43	29–66; mean=48	Not reported	Not reported	Not reported	Not reported
Authors	Han et al ⁶⁶	Gjevjon and Hellesø ⁶⁷	De Vliegher et al ⁶⁹	Bercovitz et al ⁶⁸	Alhuwail et al ⁷⁰	Vasalampi ⁷¹	Stutzel et al ⁷⁵	Santi et al ⁷²	Göransson et al ⁷³	Farsjø et al ⁷⁶	Dowding et al ⁷⁹	Dean et al., 2016 ⁷⁴	Bastide et al ⁷⁷	Arbaje et al ⁷⁸	Petersen et al ⁴⁰
Study purpose	Access electronic health records for home care	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Facilitate client assess- ment and care planning	Enhance communication				

G
Ð
Ē
- 5
Ħ
5
,0,
\cup
\sim
\sim
_
-
<mark>e</mark> 1
ole 1 (
ble 1 (
able 1 (
Table 1 (

 \sim

Study purpose	Authors	Age of participants (range)	Gender of participants	Study design	Data collection method	Hardware used	Software used
Enhance communication	Lyngstad et al ⁸¹	Intervention group: 38.6–40.5 (mean = 39.6) Control group: 38.4–40.4 (mean = 39.4)	Both genders (mostly female)	Quantitative	Survey Questionnaire	Desk computer	Web based
Enhance communication	Lyngstad and Hellesø ⁸⁰	Intervention group: 38.6–40.5 (mean = 39.6) Control group: 38.4–40.4 (mean = 39.4)	Both genders (mostly female)	Quantitative	Survey Questionnaire	Desk computer	Web based
Enhance communication	Lindberg et al ¹⁰¹	Not reported	Not reported	Systematic review	Not applicable	Tablet and laptop	Web based and application
Enhance communication	Gentles et al ⁸²	Not reported	Not reported	Scoping review	Not applicable	Computer and phone	Web based and phone based
Enhance communication	Chiang and Wang ²	27–54	Female	Qualitative	Interviews	Phone	Application
Enhance communication	Bossen et al ⁸³	30–75	Both genders	Qualitative	Interviews, observa- tions, focus groups	Tablet and computer	Web based
Provide care remotely	Mangwi Ayiasi et al ⁸⁴	Not reported	Not reported	Quantitative	Survey Questionnaire	Phone	Application
Provide care remotely	Little et al ⁸⁵	Not reported	Not reported	Qualitative	Observations	Phone	Application
Provide care remotely	Katalinic et al ³⁵	Not reported	Not reported	Quantitative	Survey Questionnaire	Tablet	Application



Fig. 1 Scholarly literature article search results. ICT, information and communication technologies.

other countries. Half of the studies were published from 2010 to 2015 and a half after 2015 (**Supplementary Figs. S1** and **S2**, available in the online version).

Characteristics of the Research Conducted

Population

Of the total included studies, 38 (97.5%) did not use the term "health care aide" or "personal support worker" to refer to the health care workers. However, we considered these studies as they were conducted within samples with a similar scope of practice. Also, most studies did not report the age of their participants. For that reason, it was not possible to extract this information. Also, 24 (60%) studies

did not report participant gender or sex. Only 11 (27.5%) studies reported male and female participants, and 5 (12.5%) reported only female participants.

Study Design

Most studies were qualitative (n = 17, 42.5%) and quantitative (n = 13, 32.5%), with a smaller proportion of studies that included multiple (n = 5, 12.5%) or mixed-method designs (n = 1, 2.5%). One systematic review and one literature review were also included. Two studies did not report their study designs. We extracted 53 data collection methods from the selected studies, as some authors used multiple methods. For that reason, this denominator is not adding up to the total number of selected studies.



Fig. 2 Distribution of barriers reported in the selected studies.

We found five different types of data collection techniques: interviews (n = 14, 26.4%), focus groups (n = 13, 24.5%), survey/questionnaires (n = 12, 24.5%), observation (n = 9, 17%) and one for workshops; and two different types of data analyses: secondary analysis (n = 3, 5.6%) and retrospective analysis (n = 1, 1.8%). We present a summary of study designs and data collection methods in **–** Supplementary Table S2 (available in the online version).

Study Purpose

We classified the selected studies in five groups, according to the main purpose of ICT use by health care aides: (1) improve everyday work (n = 12, 30%),^{46,51-61} (2) access electronic health records for home care (n = 9, 22.5%),⁶²⁻⁷⁰ (3) facilitate client assessment and care planning (n = 9, 22.5%),⁷¹⁻⁷⁹ (4) enhance communication (n = 7, 17.5%),^{2,40,80-83} and (5) provide care remotely (n = 3, 7.5%).^{35,84,85}

Features of the Reported Information and Communication Technologies

In **-Table 1**, we report our findings regarding the hardware and software used in the ICT referred in the selected studies. Desk computers were the most used form of hardware (n = 11, 27.5%), followed by electronic tablets (n = 9, 22.5%), mobile phones (n = 6, 15%), and laptops (n = 3, 7.5%). Some studies reported a combination of hardware between tablets and desk computers (n = 2, 5%), tablets and mobile phones (n = 2, 5%), tablets and laptops (n = 1, 2.5%), or desk computers and mobile phones (n = 1, 2.5%). Five studies (12.5%) did not report the type of hardware used.

Regarding software, most studies used a web-based platform (n = 21, 52.5%), followed by mobile applications (n = 10, 25%). A few studies used mobile applications and a webbased platform (n = 4, 10%). One study (2.5%) reported a text messaging technology, and four studies (10%) did not report the type of software.

Barriers and Benefits Identified in Adopting Information and Communication Technologies by Health Care Aides

In this section, we present the information regarding barriers and benefits of ICT used by health care aides.

Barriers

Several studies have explored the barriers of ICT implementation and adoption for various health care professionals.^{45,86–95} Still, the evidence of ICT related barriers for health care aides is limited. The concept of barriers is associated with attitudes, education, training, and limitations for the adoption and use of ICT.⁹² Barriers vary, and there is a need to address them to implement ICT.⁸⁶ In the literature, the evidence reports that barriers could be related but not limited to inadequate access to useful, relevant, and appropriate hardware and software when implementing ICT.⁹⁴ ICT barriers have been observed from the technology, organizational, socioeconomic, and ethical perspectives.⁹⁶

In **Fig. 2**, we present a list of the barriers we identified for the adoption and implementation of ICT for health care aides. In **Supplementary Table S3** (available in the online version), we include an entire list of barriers category, frequency, and the corresponding studies.

In this review, we found 128 barriers related to the adoption and use of ICT by health care aides. The overarching categories were as follows: (1) incomplete hardware and software features, (2) time-consuming ICT adoption, (3) heavy or increased workloads, (4) perceived lack of usefulness of the ICT, (5) cost or budget restrictions, (6) security and privacy concerns, (7) lack of collaboration and integration with current technologies, (8) data quality or validity and communication issues, (9) technical support, and (10) related to the provider–client relationship.

In the group of barriers related to hardware and software, we found that users reported system malfunctions, poor design, and issues regarding software configuration.^{35,40,46,63,66,67,70,71,85} In these studies, the software did not reflect users' needs and expectations or did not

support their information needs.⁷⁰ Some studies described the lack of documentation features.^{35,46,65,70,72} In addition to usability issues,⁵⁹ we found connectivity issues, as a significant hardware barrier in this group.^{56,64,71,84} Some studies reported incomplete hardware features as a significant barrier to adopting and implementing ICT.^{56,65,77,86}

Time-consuming adoption and implementation was barrier another significant for health care aides.^{2,35,40,51,53,55,56,59,63-65,85} This further amplifies the need for communication with ICT providers in order for an effective ICT solution.^{71,72,76} In some studies, users reported the ICT implementation process as time consuming.^{40,52,53,63–65,85} A few studies described the lack of time for documentation between clients and providers as a significant barrier.53 These groups of studies reported that ICT required more time from health care aides.^{2,51,56} Likewise, we found that users referred to ICT as a waste of time, particularly during the documentation process.

Studies presenting barriers related to heavy or increased workloads^{40,52,55,59,69,71–73,83} argued that users perceived a growth in daily workloads due to the use and implementation of ICT. For example, Petersen et al⁴⁰ reported problematic ICT use due to report writing taking precedence over the care of their patients. Heavy workload of health care aides was referred as a major barrier of adoption and implementation of new ICT.^{71,72}

Other studies on the usefulness of ICT,^{55,57,58,64,70,71,74} suggesting that ICT did not meet health care aides' needs⁶⁴ or represent an immediate advantage.⁵⁵ Another significant barrier we found was the associated cost or budget restriction.^{35,52,57,64,76} A few studies referred to concerns such as return on investment,⁶⁴ licensing and purchasing costs,³⁵ and financial restrictions to implement or accept the ICT solution.⁷⁶

Security and privacy concerns were also highlighted as significant barriers to ICT adoption by health care aides.^{2,40,60,72,77} In some studies, users perceived risk regarding their information being leaked, stolen,^{2,60} or tracked by malicious users.⁷⁷ Other studies highlighted the privacy of information on the devices as an issue.⁴⁰

We found barriers associated with the lack of collaboration and integration with current technologies^{57,59,68,71} and existing systems.^{35,72,76} Such barriers included lack of technical knowledge or skills for implementing and effectively using ICT,^{53,83} concerns about interoperability issues,^{64,66,69} validity of information (i.e., incompleteness, errors, and inaccurate information),^{73,78} lack of continuity in the use of the ICT solution,^{40,76} and limitations in software features and system designs.^{57,59}

Data quality and validity^{40,62,72} and communication issues were also reported^{40,81} as barriers. A few studies reported technical support as a major barrier.^{56,70} Other ICT were reported as not aligned with actual workflows⁵⁶ or posed difficulty for scaled or customized implementation.³⁵ The lack of technical support and learning or training requirements, as well as limited flexibility, were also described as barriers.⁴⁰ Finally, some studies indicated that the provider-client relationship was negatively affected by the implementation of ICT.^{63,64,71} In summary, the most predominant barriers related to the ICT for health care aides were related to incomplete or limited software features, ^{35,40,46,52,63,66,67,70,71,85} time-consuming ICT (i.e., in use, adoption, or implementation), ^{2,35,40,51,53,55,56,59,63} ^{64,65,85} heavy or increased workloads, ^{40,52,55,59,69,71-73,83} and health care aides perceived ICT as not convenient or a useful solution. ^{55,57,58,64,70,71,74,83}

Benefits

Several authors have documented the benefits of ICT for health care providers.^{66,78,85,87,95,97} However, there is little evidence on the benefits of technology for health care aides. Benefits of ICT are described as tangible or perceived outcomes derived from the implementation or adaptation of ICT.^{98,99} The benefits of using new ICT are usually described with great positivity. A clear description of the benefits of ICT for health care aides can inform the decision-making process regarding the implementation and adoption of new technologies for health care aides.³⁶ The evidence suggested that benefits related to ICT for health care professionals. These benefits are related to improvements in documentation, monitoring, and quality of care, and support

to existing workflows and information exchange, and enhancement in communications.¹⁰⁰

In **Fig. 3**, we report the distribution of benefits identified for ICT used by health care aides. In **Supplementary Table S4** (available in the online version), we present an entire list of the benefits we found.

The main benefits we found in this review were related to (1) improvements in communication, (2) support to work-flows and processes, (3) improvements in resource planning and health care aides services associated cost and time, (4) improve access to information and documentation, (5) improve documentation quality and efficiency, and (6) improvements in the coordination of care and patient's care follow-up at home, extending care for patients in the community.

Improvements in communication compared with conventional methods, such as fax, traditional mailing, and telephone,^{2,80,81,101} was a consistently reported benefit of ICT for health care aides. In these studies, users reported that ICT improved communication between caregivers, patients,^{51,54,72} and staff^{64,65} by introducing tools such as video calls and telehealth.^{35,57} Several studies reported the



benefits of telehealth and remote care provided by ICT.^{32,102} We observe health care aides' opportunity to provide services in distant places.^{102,103} For example, ICT can support health care aides to deliver high-quality health care in remote territories and regions in rural communities and isolated populations.^{63,64,102} The need for telehealth services for older adults, especially those living with dementia, has increased over the last few years.¹⁰⁴ A few authors have described how telehealth can improve health care service delivery.^{35,54,102,104,105}

Also, studies reported that ICT enabled a better cultural understanding by delivering information in different languages.⁷⁰ In addition, we found that ICT supported communication across locations, personnel, and families.⁷⁶ Moreover, a few studies reported that ICT improve older adults communication capacity about their health concerns.⁷³ Finally, ICT were found to allow health care aides to respond to issues rapidly.⁸³

Another group of studies reported that ICT supported workflows and processes^{2,35,53,54,66,69,71,72,74,77,81,83} by improving time management,^{74,77,81} simplifying and standardizing procedures,⁷² reducing repetitive actions,² and supporting schedule planning.^{2,40,52,61,77,82,84,101} As an example, Stroulia et al⁵⁴ reported that ICT for health care aides supported automatization and adaptative scheduling which could potentially reduce travel times and missed appointments. Katalinic et al³⁵ reported how ICT can support health care aides' remote practice to provide caregiving services to clients in distant locations.

We found a group of studies reporting benefits related to better work and staff organization associated with ICT.⁷¹ Lastly, ICT were observed as a factor for improving staff planning,^{66,71} enable caregiving services in a timely manner,⁸⁴ and promote time savings in comparison with existing methods of communication.⁴⁰ Regarding the benefit of costs and service utilization,^{2,101} Chiang et al² reported the reduction of medical costs and service consumptions through the use of smartphones applications to provide home care services. Also, Lindberg et al¹⁰¹ identified several publications describing how ICT for health care aides can reduce cost of caregiving services compared with traditional modes of delivery.

Furthermore, a few studies described accessibility to information and documentation as an enormous benefit^{40,53,54,57,59,64,80,101} by providing health care aides quick access or real-time updates to their clients information.^{54,59} A significant group of studies reported that ICT improved documentation quality and efficiency,^{64,65,68} by improving consistency in documentation,⁷² with more precise and updated information.^{54,57,59,76}

Lastly, we found another group of benefits that reported improvements in the coordination of care and patient followup at home. These benefits have proven to extend the care for patients in the community,^{2,35,82,84,101} and improve interactions between clients and health care aides.^{52,56,57,61,75} The effectiveness of ICT was described as a benefit by providing easy access to health care plans,^{35,53,57,72} promoting flexibility and interoperability,^{40,63,69} and assisting patient safety.^{54,57,71,73}

Discussion

This review examined the range and extent of ICT used by health care aides to manage and coordinate care delivery and their workflow with clients. We included 40 studies selected from the academic literature. We organized our findings into five categories of purposes for implementing ICT among health care aides. Additionally, we found a diversity of settings and arrangements for hardware and software to deploy these technologies within health care services and organizations.

In the literature, we identified evidence suggesting that current technologies for health care aides are inefficient and often do not fit their workflows,⁴² or outdated and ineffective.³⁷ From a benefits perspective, ICT is presented as a facilitator of effective communication for health care aides.¹⁵ In this review, we shared additional knowledge about the range and extent of ICT used by health care aides, as well as enhanced our understanding of the barriers and benefits associated with the use, adoption, and implementation of ICT by health care aides.

The literature suggests that across the health care sector, the barriers and benefits of ICT are common for health care professionals.^{87,95} Overall, we identified 128 barriers and 130 benefits associated with ICT used by health care aides. Even though ICT can face significant barriers in implementation, we found positive information regarding their benefits. Our findings are consistent with the academic literature that exposed the barriers and benefits of use and implemention of ICT in the health care sector for other providers.^{15,35,42,45,66,89,97,106,107}

Several authors acknowledge the difficulties related to implementing technologies in health care settings.⁴⁵ The common barriers for the adoption and implementation of ICT within this sector are related to resource requirements and costs,^{108,109} technological limitations, and lack of global standards and privacy concerns.^{110–112} These barriers are aside from the previously established structural organizational barriers.⁸⁶

In the academic literature, there are several examples describing the barriers related to the adoption and implementation of ICT in health care settings that can be analyzed under the lens of technology acceptance models.^{45,113–115} Furthermore, the expected benefits of using ICT could inform future implementations.¹¹⁶ Benefits of good-quality software, compatibility and interoperability with other information systems¹¹⁷ would promote the adoption of these technologies among health care aides.¹¹⁸

In this review, our findings suggested that health care aides could experience multiple barriers to use and adopt new ICT in their workflows and practice. The main barriers we found were related to incomplete hardware and software features, time-consuming ICT adoption, and heavy or increased workloads. In past studies, authors have described the financial, organizational, structural, cultural, and technical difficulties to adopting ICT⁸⁹ and additional barriers to ICT use and adoption by health care aides, such as professional skills¹⁰² and digital health education and literacy.¹¹⁹ To overcome these barriers, authors have recommended a

more comprehensive¹²⁰ and participatory design process involving health care aides' needs and experiences.^{60,121,122}

Likewise, we observed that health care aides could adopt and use ICT to support new models of care, such as telehealth for client follow-up, and consultation. However, traditional work practices have shaped health care aides' current workflows and procedures. As a result, health care aides face multiple barriers to adopt and use ICT in their practice. One significant barrier is the organizational and provider infrastructure,⁴² as providers generally use ICT-specific electronic patient information systems. By increasing their own awareness about the use, capabilities, and benefits of integrating ICT in critical activities of integrated care, health care aides could take advantage of the potential of these technologies to improve activities, such as prevention,⁴² collaboration, and delivery of care in innovative ways using ICT.

Barriers to access to data, even at regional levels, present an obstruction to deliver care by health care aides.⁴² Accessing and updating information about client's observations and medical conditions are essential for health care aides to provide better caregiving services. Evidence suggests that information exchange also can improve health care aides' communication with other health professionals.¹⁵ Thus, we support the argument that access to information can enhance the ability of health care aides to provide higher quality caregiving services^{37,97,102} and make better decisions to improve the quality of care.³⁶

Since health care aides face structural challenges in their profession, such as lack of regulation and standardized scope of practice, it is crucial to consider these barriers beforehand when implementing new ICT.^{8,9,12,29,123} New ICT developments should consider both, the structural and the technological barriers within the health care sector.¹²⁴

ICT can support health care aides' work by facilitating caregiving¹²⁵ and support the long-term relationship between patients and health care providers.⁹⁷ By implementing and adopting ICT in health care aides' professional practice, they could support caregiving services for numerous people. In summary, ICT could serve as a strategy to enable socially isolated people to utilize health care aides' services,^{36,102,126,127} and address the lack of direct support from professionals available to provide care services¹²⁸ and leverage the shortage of skilled and well-trained health care professionals.¹²⁹

Involving health care aides into the process of designing, implementing, and customizing new ICT is crucial to increase the rate of adoption.^{36,37} As the older adult population demands more caregiver professional services, we could expect an increasing number of new ICT to support health care aides.^{33,36,125} For that reason, we should consider the findings reported in this review related to barriers and benefits of ICT for health care aides.

Future Recommendations

After reviewing an extensive compilation of studies for this review, we recognize the need for more straightforward ICT aligned with health care aides' workflows. In addition, we must consider the involvement of end-users in design, training, and education, and clarity about the use of ICT. The voice of the end-user is essential for developing ICT tools. Future developments should include feedback from users in every step of development and the mindset of improving care. This could be done in participatory design methods¹³⁰ to engage users in the ICT.

The novel coronavirus disease 2019 (COVID-19) pandemic has accelerated the adoption of ICT tools,^{29,105} such as telehealth applications.¹⁰⁵ These applications include remote consultations for seniors,¹³¹ specifically persons living with dementia, to minimize the risk of contracting COVID-19.¹³² The pandemic has impacted the role of health care aides as caregivers by limiting their interactions and communications with their clients and families,¹³³ as well as it has revealed limitations in their ability to provide remote services. This is considering the gaps between different health care facilities and their preparedness to provide telehealth services with patients who have no internet connection or phone service.¹³⁴

At the same time, the pandemic revealed opportunities for the adoption and implementation of ICT tools to monitoring COVID-19 symptoms¹⁰⁹ and coordinating and delivering care.^{135,136} In the upcoming years, we can expect a greater demand, adoption, and use of ICT to improve and enable the quality of care. Identifying the health care aides' needs and requirements for design is crucial,¹²¹ as this will pave the road to acceptance and adoption of the ICT. We also believe that critical sociocultural factors need to be understood before developing ICT.¹²¹ In the future, health care aides with technical training may be more receptive to ICT.³⁸ This could help health care aides to overcome multiple barriers identified in this study.

Finally, although the literature reporting information specific to health care aides is scarce, and there is limited research on evidence-based practices for this population, we cannot understimate the importance of health care aides to sustain the health care system. Indeed, many countries around the world face a shortage of these groups of service providers. In this review, we have seen how ICT can help health care aides in their practice. However, more research is needed to support evidence-based ICT.

Limitations of the Study

This literature review has some limitations. First, despite our efforts to conduct a comprehensive and exhaustive search, the academic literature lacks consistency concerning our search terms. We were looking for healthcare workers, such as health care aides and personal support workers. Due to the limited literature involving this population, we decided to include nurses and personal support providers in studies that referred home care services. We assumed that their experiences are similar, considering different ways of pointing health care aides across Canada and other countries. Second, it was hard to classify barriers and benefits, as some articles reported these indistinctly. For that reason, we defined meanings for each group, and after a process of

open discussion, such definitions were organized accordingly. Third, although we identified several barriers and benefits in selected studies, not all studies comprehensively reported all perceived benefits and barriers associated with ICT for health care aides.

Conclusion

Despite challenges, the literature identifies that ICT can improve health care aides' workflow and quality of care for their clients. The main issue we experienced was the scarce literature related specifically to health care aides. We achieved the aim of this literature review to understand the barriers and potential benefits of ICT adoption for health care aides. We classified this information by purpose as reported in the selected studies. Future directions should examine the impact of the ICT on workflows of health care aides.

Clinical Relevance Statement

This literature review showed the information and communication technologies (ICT) used by health care aides to manage and coordinate the care-delivery workflow for their clients. In addition, we have explored the factors that would assist in the development and adoption of electronic platforms, which could improve the overall efficiency of care provided by health care aides.

Multiple Choice Questions

- 1. What is the most common barrier to implement and adopt information and communication technologies (ICT) faced by health care aides?
 - a. Incomplete hardware and software features
 - b. Time-consuming ICT adoption
 - c. Heavy or increased workloads
 - d. Perceived lack of usefulness of ICT

Correct Answer: The correct answer is option a, as health care aides reported system malfunctions, poor design, and issues regarding software configuration. In these studies, the software did not reflect users' needs and expectations or did not support their information needs. Some studies described the lack of documentation features. In addition to usability issues we found connectivity issues, as a hardware significant barrier in this group. Some studies reported incomplete hardware features as a significant barrier to adopting and implementing ICT.

- 2. What is the most common benefit for health care aides to implement and adopt information and communication technologies (ICT)?
 - a. Improvements in communication
 - b. Improvements in planning
 - c. Improvement in organization staffing
 - d. Increase overall productivity

Correct Answer: The correct answer is option a. We found this benefit as the most common in this review. Improvements in communication include communication between caregivers and patients and staff, better cultural understanding. In addition, ICT can support communication across locations, personnel, and families and allow clients to inform their concerns to health care aides.

Author Contributions

All authors contributed to the study's conception and design. Data extraction, analysis, and synthesis were performed by H.P., N.N., and S.M. The first draft of the manuscript was written by H.P., N.N., S.M., and S.P., and all authors commented on previous versions of the manuscript. L.L., A.M.C., and N.N. read and approved the final manuscript.

Funding

This project is supported by the Centre for Aging and Brain Health Innovation (CABHI) and Alberta Innovates grant number G2019000525 AICE-ABH, received by L.L. as principal investigator.

Conflict of Interest

None declared.

Acknowledgments

We acknowledge Lauren McLennan, Peyman A. Khaneghah, Wang C. Tang, and Elyse Letts who provided support as research assistants.

References

- 1 World Health Organization. Ageing and health. Accessed May 3, 2021 at: https://www.who.int/news-room/fact-sheets/detail/ ageing-and-health
- 2 Chiang KF, Wang HH. Nurses' experiences of using a smart mobile device application to assist home care for patients with chronic disease: a qualitative study. J Clin Nurs 2016;25 (13,14):2008-2017
- ³ Alzheimer Society of Canada. Report summary prevalence and monetary costs of dementia in Canada (2016): a report by the Alzheimer Society of Canada. Health Promot Chronic Dis Prev Can 2016;36(10):231–232
- 4 Personal support worker. Accessed March 20, 2021 at: https:// settlement.org/alternative-jobs/nurse/personal-support-worker/
- 5 Personal Support Worker HQ. Personal support worker? Accessed January 21, 2022 at: https://www.centennialcollege.ca/programs-courses/full-time/personal-support-worker/
- 6 Online ONIP. Personal support worker (PSW) in Canada. Accessed March 2, 2021 at: http://www.onip.ca/personal-support-worker/
- 7 Saari M, Xiao S, Rowe A, et al. The role of unregulated care providers in home care: a scoping review. J Nurs Manag 2018;26 (07):782–794
- 8 Hewko SJ, Cooper SL, Huynh H, et al. Invisible no more: a scoping review of the health care aide workforce literature. BMC Nurs 2015;14(01):38
- 9 Berta W, Laporte A, Deber R, Baumann A, Gamble B. The evolving role of health care aides in the long-term care and home and community care sectors in Canada. Hum Resour Health 2013;11 (01):25

- 10 Lilly M. Medical versus social work-places: constructing and compensating the personal support worker across health care settings in Ontario, Canada. Gend Place Cult 2008;15(03): 285–299
- 11 Arain MA, Deutschlander S, Charland P. Are healthcare aides underused in long-term care? A cross-sectional study on continuing care facilities in Canada. BMJ Open 2017;7(05):e015521
- 12 Estabrooks CA, Squires JE, Carleton HL, Cummings GG, Norton PG. Who is looking after mom and dad? Unregulated workers in Canadian long-term care homes. Can J Aging 2015;34(01):47–59
- 13 White DE, Jackson K, Besner J, Norris JM. The examination of nursing work through a role accountability framework. J Nurs Manag 2015;23(05):604–612
- 14 Franzosa E, Tsui EK, Baron S. Home health aides' perceptions of quality care: goals, challenges, and implications for a rapidly changing industry. New Solut 2018;27(04):629–647
- 15 Saari M, Patterson E, Killackey T, Raffaghello J, Rowe A, Tourangeau AE. Home-based care: barriers and facilitators to expanded personal support worker roles in Ontario, Canada. Home Health Care Serv Q 2017;36(3,4):127–144
- 16 Kroezen M, Schäfer W, Sermeus W, Hansen J, Batenburg R. Healthcare assistants in EU member states: an overview. Health Policy 2018;122(10):1109–1117
- 17 Janet L, Jennifer S, Alvin Y. Ontario personal support workers in home and community care: CRNCC/PSNO survey results. Accessed March 20, 2021 at: https://www.ryerson.ca/content/ dam/crncc/knowledge/infocus/factsheets/InFocus-Ontario PSWs in Home and Community Care.pdf
- 18 Booi L, Sixsmith J, Chaudhury H, O'Connor D, Young M, Sixsmith A. 'I wouldn't choose this work again': perspectives and experiences of care aides in long-term residential care. J Adv Nurs 2021; 77(09):3842–3852
- 19 Kelly C, Bourgeault IL. The personal support worker program standard in Ontario: an alternative to self-regulation? Healthc Policy 2015;11(02):20–26
- 20 National Alliance for Caregiving. Caregiving in the U.S.. 2020. Accessed January 21, 2022 at: https://www.caregiving.org/caregiving-in-the-us-2020/
- 21 Statistics Canada. Care counts: care receivers in Canada, 2018. Accessed February 1, 2021 at: https://www150.statcan.gc.ca/n1/ pub/11-627-m/11-627-m2020002-eng.htm
- 22 Government of Canada. Government of Canada announces funding to train 4,000 personal support worker interns, support sector. Accessed May 18, 2021 at: https://www.canada.ca/en/ employment-social-development/news/2020/12/governmentof-canada-announces-funding-to-train-4000-personal-supportworker-interns-support-sector.html
- 23 CBC News. Ontario's PSW shortage is impacting people living with disabilities in Waterloo region. Accessed May 18, 2021 at: https://www.cbc.ca/news/canada/kitchener-waterloo/pswshortage-ontario-waterloo-region-impacts-1.5953472
- 24 Muir T. Measuring social protection for long-term care. Accessed January 21, 2022 at: https://www.oecd-ilibrary.org/socialissues-migration-health/measuring-social-protection-for-longterm-care_a411500a-en
- 25 CIHI. Canada's seniors population outlook: uncharted territory. Accessed May 18, 2021 at: https://www.cihi.ca/en/infographiccanadas-seniors-population-outlook-uncharted-territory
- 26 United Nations Department of Economic and Social Affairs PD. World population prospects 2019: highlights. Accessed January 21, 2022 at: https://population.un.org/wpp/Publications/ Files/WPP2019_Highlights.pdf
- 27 Brophy JT, Keith MM, Hurley M, McArthur JE. Sacrificed: Ontario healthcare workers in the time of COVID-19. New Solut 2021;30 (04):267–281
- 28 Drennan VM, Ross F. Global nurse shortages-the facts, the impact and action for change. Br Med Bull 2019;130(01):25–37

- 29 Estabrooks CA, Straus SE, Flood CM, et al. Restoring trust: COVID-19 and the future of long-term care in Canada. Facets 2020;5 (01):651–691
- 30 Baker SB, Xiang W, Atkinson I. Internet of things for smart healthcare: technologies, challenges, and opportunities. IEEE Access 2017;5:26521–26544
- 31 Tyagi RK, Cook L, Olson J, Belohlav J. Healthcare technologies, quality improvement programs and hospital organizational culture in Canadian hospitals. BMC Health Serv Res 2013;13 (01):413
- 32 Boric-Lubecke O, Gao X, Yavari E, Baboli M, Singh A, Lubecke VM. E-healthcare: Remote monitoring, privacy, and security. In: 2014 IEEE MTT-S International Microwave Symposium (IMS2014). IEEE; 2014:1–3; Tampa, FL
- 33 Report: 2020 national survey of Canadian nurses: use of digital health technology in practice. Accessed January 21, 2022 at: https://www.infoway-inforoute.ca/en/component/edocman/ 3812-2020-national-survey-of-canadian-nurses-use-of-digitalhealth-technology-in-practice/view-document?Itemid=0
- 34 Forum on Aging, Disability, and Independence; Board on Health Sciences Policy; Division on Behavioral and Social Sciences and Education; Institute of Medicine; National Research Council. The Future of Home Health Care: Workshop Summary. Washington, DC: National Academies Press; 2015
- 35 Katalinic O, Young A, Doolan D. Case study: the interact home telehealth project. J Telemed Telecare 2013;19(07):418–424
- 36 Lindeman DA, Kim KK, Gladstone C, Apesoa-Varano EC. Technology and caregiving: emerging interventions and directions for research. Gerontologist 2020;60(Suppl 1):S41–S49
- 37 Sterling MR, Dell N, Tseng E, et al. Home care workers caring for adults with heart failure need better access to training and technology: a role for implementation science. J Clin Transl Sci 2020;4(03):224–228
- 38 King S, Liu L, Stroulia E, Nikolaidis I. Using simulations to integrate technology into health care aides' workflow. Int J Adv Corp Learn 2013;6(02):28
- 39 Wildevuur SE, Simonse LW. Information and communication technology-enabled person-centered care for the "big five" chronic conditions: scoping review. J Med Internet Res 2015; 17(03):e77
- 40 Petersen HV, Foged S, Madsen AL, Andersen O, Nørholm V. Nurses' perception of how an e-message system influences cross-sectoral communication: a qualitative study. J Nurs Manag 2018;26(05):509–517
- 41 Aslani N, Ahmadi M, Samadbeik M. A systematic review of the attributes of electronic personal health Records for Patients with multiple sclerosis. Health Technol (Berl) 2020;10(03):587–599
- 42 Steele Gray C, Barnsley J, Gagnon D, et al. Using information communication technology in models of integrated communitybased primary health care: learning from the iCOACH case studies. Implement Sci 2018;13(01):87
- 43 Sterling MR, Silva AF, Leung PBK, et al. "It's like they forget that the word 'health' is in 'home health aide'": understanding the perspectives of home care workers who care for adults with heart failure. J Am Heart Assoc 2018;7(23):e010134
- 44 Falconer E, Kho D, Docherty JP. Use of technology for care coordination initiatives for patients with mental health issues: a systematic literature review. Neuropsychiatr Dis Treat 2018; 14:2337–2349
- 45 Rahimi B, Nadri H, Lotfnezhad Afshar H, Timpka T. A systematic review of the technology acceptance model in health informatics. Appl Clin Inform 2018;9(03):604–634
- 46 Alhuwail D, Koru G. Identifying home care clinicians' information needs for managing fall risks. Appl Clin Inform 2016;7(02): 211–226
- 47 Daudt HM, van Mossel C, Scott SJ. Enhancing the scoping study methodology: a large, inter-professional team's experience with

Arksey and O'Malley's framework. BMC Med Res Methodol 2013; 13(01):48

- 48 Arksey H, O'Malley L. Scoping studies: towards a methodological framework. Int J Soc Res Methodol 2005;8(01):19–32
- 49 Robinson KA, Saldanha IJ, McKoy NA. Development of a framework to identify research gaps from systematic reviews. J Clin Epidemiol 2011;64(12):1325–1330
- 50 Belur J, Tompson L, Thornton A, Simon M. Interrater reliability in systematic review methodology: exploring variation in coder decision-making. Sociol Methods Res 2021;50(02):837–865
- 51 Brown RN, Ruggiano N, Page TF, et al. CareHeroes web and android apps for dementia caregivers. Res Gerontol Nurs 2016; 9(04):193–203
- 52 Kaunda-Khangamwa BN, Steinhardt LC, Rowe AK, et al. The effect of mobile phone text message reminders on health workers' adherence to case management guidelines for malaria and other diseases in Malawi: lessons from qualitative data from a clusterrandomized trial. Malar J 2018;17(01):481
- 53 Gars U, Skov H. The TECS model leads to active use of technology in home care. Nord J Nurs Res 2017;37(01):51–58
- 54 Stroulia E, Nikolaidisa I, Liua L, King S, Lessard L. Home care and technology: a case study. Stud Health Technol Inform 2012; 182:142–152
- 55 Sassen B, Kok G, Schepers J, Vanhees L. Supporting health care professionals to improve the processes of shared decision making and self-management in a web-based intervention: randomized controlled trial. J Med Internet Res 2014;16(10):e211
- 56 Nilsson L, Fagerström C. Decision-makers and mediators in a home healthcare digitisation process: nurses' experiences of implementation and use of a decision support system. Contemp Nurse 2018;54(4,5):511–521
- 57 Gund A, Sjöqvist BA, Wigert H, Hentz E, Lindecrantz K, Bry K. A randomized controlled study about the use of eHealth in the home health care of premature infants. BMC Med Inform Decis Mak 2013;13(01):22
- 58 Andersen GR, Bendal S, Westgaard RH. Work demands and health consequences of organizational and technological measures introduced to enhance the quality of home care services–a subgroup analysis. Appl Ergon 2015;51:172–179
- 59 Alhuwail D, Koru G. Leveraging health information technology for fall-risk management in home care: a qualitative exploration of clinicians' perspectives. Home Health Care Manage Pract 2016;28(04):241–249
- 60 Danilovich MK, Diaz L, Saberbein G, Healey WE, Huber G, Corcos DM. Design and development of a mobile exercise application for home care aides and older adult medicaid home and community-based clients. Home Health Care Serv Q 2017;36(3-4): 196–210
- 61 Lexis M, Everink I, Van Der Heide L, Spreeuwenberg M, Willems C, De Witte L. Activity monitoring technology to support homecare delivery to frail and psychogeriatric elderly persons living at home alone. Technol Disabil 2013;25(03):189–197
- 62 Westra BL, Oancea C, Savik K, Marek KD. The feasibility of integrating the Omaha system data across home care agencies and vendors. Comput Inform Nurs 2010;28(03):162–171
- 63 Yang Y, Bass EJ, Bowles KH, Sockolow PS. Impact of home care admission nurses' goals on electronic health record documentation strategies at the point of care. Comput Inform Nurs 2019;37 (01):39–46
- 64 Tapper L, Quinn H, Kerry J, Brown KG. Introducing handheld computers into home care. Can Nurse 2012;108(01):28–32
- 65 Sockolow PS, Bowles KH, Adelsberger MC, Chittams JL, Liao C. Impact of homecare electronic health record on timeliness of clinical documentation, reimbursement, and patient outcomes. Appl Clin Inform 2014;5(02):445–462
- 66 Han S, Juell-Skielse G, Smedberg Å, Aasi P, Nilsson AG. Benefits of mobile reporting systems in social home care: the case of seven

Swedish municipalities. Int J Technol Assess Health Care 2014;30 (04):409–415

- 67 Gjevjon ER, Hellesø R The quality of home care nurses' documentation in new electronic patient records. J Clin Nurs 2010;19 (1,2):100–108
- 68 Bercovitz A, Sengupta M, Jamison P. Electronic medical record adoption and use in home health and hospice. NCHS Data Brief 2010;(45):1–8
- 69 De Vliegher K, Paquay L, Vernieuwe S, Van Gansbeke H. The experience of home nurses with an electronic nursing health record. Int Nurs Rev 2010;57(04):508–513
- 70 Alhuwail D, Koru G, Mills ME. Supporting the information domains of fall-risk management in home care via health information technology. Home Health Care Serv Q 2016;35 (3,4):155–171
- 71 Vasalampi A. Adoption and use of a mobile system at home care. Stud Health Technol Inform 2017;242:1042–1046
- 72 Santi SM, Hinton S, Berg K, Stolee P. Bridging the information divide: health information sharing in home care. Can J Nurs Res 2013;45(01):16–35
- 73 Göransson C, Eriksson I, Ziegert K, et al. Testing an app for reporting health concerns-Experiences from older people and home care nurses. Int J Older People Nurs 2018;13(02):e12181
- 74 Dean KM, Hatfield LA, Jena AB, et al. Preliminary data on a care coordination program for home care recipients. J Am Geriatr Soc 2016;64(09):1900–1903
- 75 Stutzel MC, Filippo MP, Sztajnberg A, et al. Multi-part quality evaluation of a customized mobile application for monitoring elderly patients with functional loss and helping caregivers. BMC Med Inform Decis Mak 2019;19(01):1–18
- 76 Farsjø C, Kluge A, Moen A. Using a tablet application about nutrition in home care-Experiences and perspectives of healthcare professionals. Health Soc Care Community 2019;27(03):683–692
- 77 Bastide R, Bardy P, Borrel B, et al. Plas'O'Soins: a software platform for modeling, planning and monitoring homecare activities. IRBM 2014;35(02):82–87
- 78 Arbaje Al, Hughes A, Werner N, et al. Information management goals and process failures during home visits for middle-aged and older adults receiving skilled home healthcare services after hospital discharge: a multisite, qualitative study. BMJ Qual Saf 2019;28(02):111–120
- 79 Dowding DW, Russell D, Onorato N, Merrill JA. Technology solutions to support care continuity in home care: a focus group study. J Healthc Qual 2018;40(04):236–246
- 80 Lyngstad M, Hellesø R Electronic communication experiences of home health care nurses and general practitioners: a crosssectional study. Stud Health Technol Inform 2014;201:388–394
- 81 Lyngstad M, Grimsmo A, Hofoss D, Hellesø R Home care nurses' experiences with using electronic messaging in their communication with general practitioners. J Clin Nurs 2014;23 (23,24):3424–3433
- 82 Gentles SJ, Lokker C, McKibbon KA. Health information technology to facilitate communication involving health care providers, caregivers, and pediatric patients: a scoping review. J Med Internet Res 2010;12(02):e22
- 83 Bossen C, Christensen LR, Grönvall E, Vestergaard LS. CareCoor: augmenting the coordination of cooperative home care work. Int J Med Inform 2013;82(05):e189–e199
- 84 Mangwi Ayiasi R, Kolsteren P, Batwala V, Criel B, Orach CG. Effect of village health team home visits and mobile phone consultations on maternal and newborn care practices in Masindi and Kiryandongo, Uganda: a community-intervention trial. PLoS One 2016;11(04):e0153051
- 85 Little A, Medhanyie A, Yebyo H, Spigt M, Dinant GJ, Blanco R. Meeting community health worker needs for maternal health care service delivery using appropriate mobile technologies in Ethiopia. PLoS One 2013;8(10):e77563

- 86 Lluch M. Healthcare professionals' organisational barriers to health information technologies-a literature review. Int J Med Inform 2011;80(12):849–862
- 87 de Souza CHA, Morbeck RA, Steinman M, et al. Barriers and benefits in telemedicine arising between a high-technology hospital service provider and remote public healthcare units: a qualitative study in Brazil. Telemed J E Health 2017;23(06): 527–532
- 88 Alaboudi A, Atkins A, Sharp B, Balkhair A, Alzahrani M, Sunbul T. Barriers and challenges in adopting Saudi telemedicine network: the perceptions of decision makers of healthcare facilities in Saudi Arabia. J Infect Public Health 2016;9(06):725–733
- 89 Kaye R, Kokia E, Shalev V, Idar D, Chinitz D. Barriers and success factors in health information technology: a practitioner's perspective. J Manag Mark Healthc 2010;3(02):163–175
- 90 Irinoye OO, Ayandiran EO, Fakunle I, Mtshali N. Nurses' perception and barriers to use of information communication technology in a teaching hospital in Nigeria. Comput Inform Nurs 2013; 31(08):394–400
- 91 Schoen J, Mallett JW, Grossman-Kahn R, Brentani A, Kaselitz E, Heisler M. Perspectives and experiences of community health workers in Brazilian primary care centers using m-health tools in home visits with community members. Hum Resour Health 2017;15(01):71
- 92 Eley R, Fallon T, Soar J, Buikstra E, Hegney D. Barriers to use of information and computer technology by Australia's nurses: a national survey. J Clin Nurs 2009;18(08):1151–1158
- 93 Ramsey A, Lord S, Torrey J, Marsch L, Lardiere M. Paving the way to successful implementation: identifying key barriers to use of technology-based therapeutic tools for behavioral health care. J Behav Health Serv Res 2016;43(01):54–70
- 94 O' Connor Y, O' Connor S, Heavin C, Gallagher J, O' Donoghue J. Sociocultural and technological barriers across all phases of implementation for mobile health in developing countries. In: Al-Jumeily D, Hissain A, Mallucci C, Oliver C, eds. Applied Computing in Medicine and Health. Waltham, MA: Elsevier; 2016:212–230
- 95 Shah SGS, Robinson I. Benefits of and barriers to involving users in medical device technology development and evaluation. Int J Technol Assess Health Care 2007;23(01):131–137
- 96 Hassan AYI. Challenges and recommendations for the deployment of information and communication technology solutions for informal caregivers: scoping review. JMIR Aging 2020;3(02): e20310
- 97 Omboni S, Caserini M, Coronetti C. Telemedicine and M-health in hypertension management: technologies, applications and clinical evidence. High Blood Press Cardiovasc Prev 2016;23(03):187–196
- 98 Braun R, Catalani C, Wimbush J, Israelski D. Community health workers and mobile technology: a systematic review of the literature. PLoS One 2013;8(06):e65772
- 99 Early J, Gonzalez C, Gordon-Dseagu V, Robles-Calderon L. Use of mobile health (mHealth) technologies and interventions among community health workers globally: a scoping review. Health Promot Pract 2019;20(06):805–817
- 100 Thakkar M, Davis DC. Risks, barriers, and benefits of EHR systems: a comparative study based on size of hospital. Perspect Health Inf Manag 2006;3:5
- 101 Lindberg B, Nilsson C, Zotterman D, Söderberg S, Skär L. Using information and communication technology in home care for communication between patients, family members, and healthcare professionals: a systematic review. Int J Telemed Appl 2013; 2013:461829
- 102 Choukou MA, Maddahi A, Polyvyana A, Monnin C. Digital health technology for Indigenous older adults: a scoping review. Int J Med Inform 2021;148:104408
- 103 Koo BM, Vizer LM. Examining mobile technologies to support older adults with dementia through the lens of personhood and

human needs: scoping review. JMIR Mhealth Uhealth 2019;7 (11):e15122

- Brown P, Oliver E, Dening KH. Increasing need for telehealth services for families affected by dementia as a result of Covid-19.
 J Community Nurs 2020;34(05):59–64
- 105 Wosik J, Fudim M, Cameron B, et al. Telehealth transformation: COVID-19 and the rise of virtual care. J Am Med Inform Assoc 2020;27(06):957–962
- 106 Alamoodi AH, Garfan S, Zaidan BB, et al. A systematic review into the assessment of medical apps: motivations, challenges, recommendations and methodological aspect. Health Technol (Berl) 2020;10(05):1045–1061
- 107 Abelson JS, Kaufman E, Symer M, Peters A, Charlson M, Yeo H. Barriers and benefits to using mobile health technology after operation: a qualitative study. Surgery 2017;162(03):605–611
- 108 Jha AK, DesRoches CM, Campbell EG, et al. Use of electronic health records in U.S. hospitals. N Engl J Med 2009;360(16):1628–1638
- 109 Bandini J, Rollison J, Feistel K, Whitaker L, Bialas A, Etchegaray J. Home care aide safety concerns and job challenges during the COVID-19 pandemic. New Solut 2021;31(01):20–29
- 110 Yao W, Chu C-H, Li Z. The adoption and implementation of RFID technologies in healthcare: a literature review. J Med Syst 2012; 36(06):3507–3525
- 111 Scott Kruse C, Karem P, Shifflett K, Vegi L, Ravi K, Brooks M. Evaluating barriers to adopting telemedicine worldwide: a systematic review. J Telemed Telecare 2018;24(01):4–12
- 112 Tieu L, Sarkar U, Schillinger D, et al. Barriers and facilitators to online portal use among patients and caregivers in a safety net health care system: a qualitative study. J Med Internet Res 2015; 17(12):e275
- 113 Venkatesh M, Davis D. User acceptance of information technology: toward a unified view. Manage Inf Syst Q 2003;27(03):425
- 114 Davis FD. Perceived usefulness, perceived ease of use, and user acceptance of information technology. Manage Inf Syst Q 1989; 13(03):319
- 115 Venkatesh V, Davis FD. A theoretical extension of the technology acceptance model: four longitudinal field studies. Manage Sci 2000;46(02):186–204
- 116 Tavares J, Goulão A, Oliveira T. Electronic health record portals adoption: empirical model based on UTAUT2. Inform Health Soc Care 2018;43(02):109–125
- 117 Vanneste D, Vermeulen B, Declercq A. Healthcare professionals' acceptance of BelRAI, a web-based system enabling personcentred recording and data sharing across care settings with interRAI instruments: a UTAUT analysis. BMC Med Inform Decis Mak 2013;13(01):129
- 118 An J-Y, Hayman LL, Panniers T, Carty B. Theory development in nursing and healthcare informatics: a model explaining and predicting information and communication technology acceptance by healthcare consumers. ANS Adv Nurs Sci 2007;30(03): E37–E49
- 119 Jimenez G, Spinazze P, Matchar D, et al. Digital health competencies for primary healthcare professionals: a scoping review. Int J Med Inform 2020;143:104260
- 120 Tsekleves E, Darby A, Whicher A, Swiatek P. Co-designing design fictions: a new approach for debating and priming future healthcare technologies and services. Arch Des Res 2017;30(02):5–21
- 121 Aryana B, Brewster L, Nocera JA. Design for mobile mental health: an exploratory review. Health Technol (Berl) 2019;9 (04):401-424
- 122 Holden RJ, Karanam YLP, Cavalcanti LH, et al. Health information management practices in informal caregiving: an artifacts analysis and implications for IT design. Int J Med Inform 2018; 120:31–41
- 123 Afzal A, Stolee P, Heckman GA, Boscart VM, Sanyal C. The role of unregulated care providers in Canada-A scoping review. Int J Older People Nurs 2018;13(03):e12190

- 124 Peek STM, Wouters EJM, van Hoof J, Luijkx KG, Boeije HR, Vrijhoef HJM. Factors influencing acceptance of technology for aging in place: a systematic review. Int J Med Inform 2014;83(04): 235–248
- 125 Moloney J, Abbas R, Beecham S, Ahmad B, Richardson I. Challenges and Opportunities for Caregiving through Information and Communication Technology. In: Proceedings of the 13th International Joint Conference on Biomedical Engineering Systems and Technologies. SCITEPRESS Science and Technology Publications; 2020:329–336
- 126 Innes A, Morgan D, Kosteniuk J. Dementia care in rural and remote settings: a systematic review of informal/family caregiving. Maturitas 2011;68(01):34–46
- 127 Benefield LE, Beck C. Reducing the distance in distance-caregiving by technology innovation. Clin Interv Aging 2007;2(02): 267–272
- 128 Tassé MJ, Wagner JB, Kim M. Using technology and remote support services to promote independent living of adults with intellectual disability and related developmental disabilities. J Appl Res Intellect Disabil 2020;33(03):640–647
- 129 Ikeda Y, Kobayakawa M, Nakao H, Iseki F, Matsushita H. Information Technology/Artificial Intelligence Innovations Needed for Better Quality of Life in Caregiving Homes. In: Health Informatics Matsushita H, ed. Switzerland: Springer Nature; 2021:37–58

- 130 Merkel S, Kucharski A. Participatory design in gerontechnology: a systematic literature review. Gerontologist 2019;59(01):e16–e25
- 131 Neumann-Podczaska A, Seostianin M, Madejczyk K, et al. An experimental education project for consultations of older adults during the pandemic and healthcare lockdown. Healthcare (Basel) 2021;9(04):425
- 132 Tuijt R, Rait G, Frost R, Wilcock J, Manthorpe J, Walters K. Remote primary care consultations for people living with dementia during the COVID-19 pandemic: experiences of people living with dementia and their carers. Br J Gen Pract 2021;71(709):e574–e582
- 133 Greenberg NE, Wallick A, Brown LM. Impact of COVID-19 pandemic restrictions on community-dwelling caregivers and persons with dementia. Psychol Trauma 2020;12(S1):S220–S221
- 134 Shang J, Chastain AM, Perera UGE, et al. COVID-19 preparedness in US home health care agencies. J Am Med Dir Assoc 2020;21 (07):924–927
- 135 Egan K. Digital technology, health and well-being and the COVID-19 pandemic: it's time to call forward informal carers from the back of the queue. Semin Oncol Nurs 2020;36(06):151088
- 136 Goodman-Casanova JM, Dura-Perez E, Guzman-Parra J, Cuesta-Vargas A, Mayoral-Cleries F. Telehealth home support during COVID-19 confinement for community-dwelling older adults with mild cognitive impairment or mild dementia: survey study. J Med Internet Res 2020;22(05):e19434