



COVID-19 Induced Coagulopathy (CIC): Thrombotic Manifestations of Viral Infection

Swati Sharma¹ Aastha Mishra² Zahid Ashraf¹

¹Department of Biotechnology, Jamia Millia Islamia, New Delhi, India
²CSIR-Institute of Genomics and Integrative Biology, Delhi, India

Address for correspondence Zahid Ashraf, PhD, Department of Biotechnology, Jamia Millia Islamia, New Delhi 110025, India (e-mail: zashraf@jmi.ac.in; mohammadzashraf@gmail.com).

TH Open 2022;6:e70–e79.

Abstract

Coronavirus disease 2019 (COVID-19) is caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and may result in an overactive coagulative system, thereby resulting in serious cardiovascular consequences in critically affected patients. The respiratory tract is a primary target for COVID-19 infection, which is manifested as acute lung injury in the most severe form of the viral infection, leading to respiratory failure. A proportion of infected patients may progress to serious systemic disease including dysfunction of multiple organs, acute respiratory distress syndrome (ARDS), and coagulation abnormalities, all of which are associated with increased mortality, additionally depending on age and compromised immunity. Coagulation abnormalities associated with COVID-19 mimic other systemic coagulopathies otherwise involved in other severe infections, such as disseminated intravascular coagulation (DIC) and may be termed COVID-19 induced coagulopathy (CIC). There is substantial evidence that patients with severe COVID-19 exhibiting CIC can develop venous and arterial thromboembolic complications. In the initial stages of CIC, significant elevation of D-dimer and fibrin/fibrinogen degradation products is observed. Alteration in prothrombin time, activated partial thromboplastin time, and platelet counts are less common in the early phase of the disease. In patients admitted to intensive care units (ICUs), coagulation test screening involving the measurement of D-dimer and fibrinogen levels, has been recommended. Prior established protocols for thromboembolic prophylaxis are also followed for CIC, including the use of heparin and other standard supportive care measures. In the present review, we summarize the characteristics of CIC and its implications for thrombosis, clinical findings of coagulation parameters in SARS-CoV-2 infected patients with incidences of thromboembolic events and plausible therapeutic measures.

Keywords

- ▶ covid-19
- ▶ coagulopathy
- ▶ thrombosis
- ▶ inflammation

Introduction

Coronavirus disease of 2019 (COVID-19) is a viral infection caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).^{1–4} Because of its highly contagious nature and global spread, it was declared a pandemic by WHO since early March 2020.⁵ SARS-CoV-2 comprises of

positive-sense single-stranded RNA genome harboring a surface glycoprotein known as spike protein, or S proteins. These proteins are believed to be responsible for the tropism toward the specific receptors present on the cell surface of the target organism. The virus specifically targets respiratory epithelium via angiotensin converting enzyme 2 (ACE2)

received
October 29, 2021
accepted after revision
November 18, 2021

DOI <https://doi.org/10.1055/s-0042-1744185>.
ISSN 2512-9465.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

receptor, thereby making its entry into the host cells.⁶ ACE2 receptors are highly expressed in many different cell types, including lung alveolar cells, cardiac myocytes, and vascular endothelium.⁷ Primarily, SARS-CoV-2 is transmitted by inhaling viral particles, facilitating their entry into the respiratory tract.¹ Additionally, the virus disseminates through fomite transmission, depending on the different surfaces and varied time intervals in which the virus can persist.⁸ The initial presentation of COVID-19 overlaps with that of other viral syndromes and includes fever, cough, fatigue, shortness of breath, headache, diarrhea and myalgias.^{10,11,26} However, the respiratory distress syndrome accompanying COVID-19 may differ from traditional acute respiratory distress syndrome caused by other viruses of the same family. Despite the extremity of hypoxemia in this infection, there is ongoing lung damage, marked by increased respiratory compliance and shunt fraction along with heightened recognition of systemic features of a hypercoagulable state. The alteration in coagulation parameters caused by SARS-CoV-2 infection can be termed 'COVID-19 induced coagulopathy' (CIC), which may cause various adverse cardiovascular complications, ultimately leading to death in some patients.^{12,13} CIC is different from the classic DIC observed in the case of sepsis. Coagulation changes in the COVID-19 patients though mimic that of DIC but are not identical. In the patients with COVID-19 infection, a strong local pulmonary thrombotic microangiopathy along with the direct endothelial cell infection by the viral particles is induced that inflict the coagulopathic response. The elevated plasma D-dimer level is the most remarkable abnormal coagulation feature in the case of severe COVID-19 patients. Whereas, in sepsis-associated DIC, a more profound thrombocytopenia is reported. In addition, DIC patients exhibit much lower levels of clotting factors and significant decrease in plasma concentrations of coagulation inhibitors like antithrombin and protein C. These features are not observed in CIC.¹⁴

Furthermore, patients who have certain co-morbidities, in particular as associated with cardiovascular disease, are speculated to have the worst prognosis among COVID-19 patients. Amidst the cardiovascular comorbid conditions, patients with diabetes are more likely to have severe infection. Other comorbidities, such as chronic obstructive pulmonary disease, liver and renal diseases, have also been associated with poor disease progression. Several studies, as well as case reports, suggest patients with severe infection present with coagulation dysfunction, as evident from a rise in blood D-dimer, lactate dehydrogenase and total bilirubin levels. Standard coagulation parameters, like partial thromboplastin time (PTT) or activated partial thromboplastin time (aPTT) and platelet count show slight or no change in the initial presentation of the disease. Only 5% of the reported cases exhibit prolonged PT and 6% prolonged aPTT as an initial manifestation of the disease.¹⁵ However, with progression of disease and severity there is progressive prolongation of PT, whereas aPTT appears to remain largely unchanged in non-fatal COVID-19 cases vs fatal cases, with no significant correlation with disease severity or mortality.¹⁶ Initial clinical studies showed patients with elevated

level of troponin indicative of cardiac injury, but due to lack of imaging data such as cardiac magnetic resonance imaging (MRI) or echocardiography, the mechanism remained unclear. Significantly higher deaths are reported in patients with cardiac injury (51.2%) as compared with those without any cardiovascular complication (4.5%).¹⁷ Also, there is speculation that the virus may act through direct cell damage upon ACE2 receptor binding, causing a systemic inflammatory response with an arising cytokine storm, destabilizing coronary plaques, and aggravating hypoxia, which represent risk factors for thrombotic events.¹⁶

SARS-CoV-2 may have a direct effect on hemostasis or it may act indirectly by creating pathological effects such as hypoxia and inflammation that may predispose COVID-19 patients to thromboembolic events. Preliminary data are suggestive that hemostatic abnormalities such as CIC, occur in COVID-19 patients.^{16,18} Also, the heightened inflammatory response causing cytokine storm, severe illness, and underlying traditional risk factors may all predispose an infected patient to thrombotic events or other cardiovascular manifestations (►Fig. 1).¹⁹ Therefore, an understanding of CIC mediated thrombotic complications in COVID-19 is of utmost importance for better management of the severe cases of the infection. In the present review we summarize all the interpretations of the characteristics of CIC till date with its implications for thrombosis and its management. We also present clinical findings related to coagulation parameters in infected patients with incidences of thromboembolic events and probable therapeutic measures for the effective management of the disease.

Thrombotic Incidences in COVID-19 Patients Exhibiting CIC

Up to November 2021, more than 253.1 million cases of COVID-19 have been reported globally accounting for ~5.1 million deaths.²⁰ There are also reports of asymptomatic carriers^{15,16,21,22}). The association of abnormal coagulopathy with infection was first observed in the initial reports from Wuhan, China showing 6% elevation in aPTT, 5% in PT and 36% in D-dimer in the first 99 patients admitted to a hospital.²³ These patients showed increased inflammatory markers, including interleukin-6 (IL-6), erythrocyte sedimentation rate (ESR), and C-reactive protein (CRP).¹⁵ The initial reports were suggestive of high risk of thromboembolic complications in these patients. With the progression of the pandemic, numerous other reports have revealed prolonged PT, high D-dimers, along with high levels of fibrinogen and high CRP, lymphocytopenia and mild thrombocytopenia.^{8,18} A systemic review consisting a total of 36 studies including the patients requiring ICU placement, presented the pooled incidence rate of 28% (95% CI, 22–34%) for venous thromboembolism (VTE).²⁴ Micro-thrombosis formation was noted in 80% of lung autopsies from fatal COVID-19 cases.²⁵ High incidence of VTE (31%) resulting in complications such as pulmonary embolism (PE) (80%), as well as arterial thrombosis (3.7%), has been reported in COVID-19 patients by Klok et al.²⁶ In ►Fig. 2,^{16,26–36} we




Fig. 1 Depiction of effect and manifestation of COVID-19 and associated cardiovascular implications.

have summarized the incidences of venous thromboembolism (VTE) reported in COVID-19 patients, according to different studies.

Molecular Mechanism Resulting in CIC Mediated Thrombosis

Endotheliitis and Hyperinflammation

Endotheliitis and hyperinflammation initiates CIC. The main site of viral infection in humans is lung alveolar type II epithelium, where the spike protein S1 attaches itself to the ACE-2 receptor.^{6,9} The virus replicates in the infected cells causing inflammatory response resulting in pyroptosis.

The receptors of innate immunity like TLR (Toll-like receptors) detect danger-associated molecular patterns (DAMPs) such as ATP and DNA released after pyroptosis and evoke intense inflammatory responses along with release of proinflammatory chemokines and cytokines from neighboring cells³⁷ (→ Fig. 3). Also, TLR-7 and TLR-8 recognizes the multiple single-stranded RNA fragments of SARS-CoV-2, thus contributing to innate immune hyper activation and causing a cytokine storm.³⁸ Thus, upon infection, the vascular endothelium, which is a guardian of vascular integrity, gets hijacked by the virus and then exposed to the milieu of proinflammatory cytokines elicited by the innate immunity.³⁹ All the studies conducted have hypothesized that the entry of the virus via endothelial-expressed ACE-2 paves the way by which the virus can enter and infect other tissues.

ACE-2 is also highly expressed in various extrapulmonary sites, including blood vessels, heart, kidney, and intestine.^{40,41} Clinical data from infected patients have revealed increased levels of IL-6, IL-1 β , MCP-1 (monocyte chemo-attractant protein 1), MIP (macrophage inflammatory protein), IFN (interferon)- γ , and IP10 (CXCL10).¹ Disruption of endothelial function and integrity by these proinflammatory cytokines leads to release of von Willebrand factor (VWF), upregulation of adhesion molecules such as ICAM (intercellular adhesion molecule)-1, integrin $\alpha\beta$ 3, P-selectin, E-selectin and endothelial tissue factor (TF, CD142).^{42,43} The sudden activation of endothelial cells causes elevated VWF antigen (VWF:Ag) levels. The function of VWF is mediated through proteolysis guided by von Willebrand factor-cleaving protease, which is also known as a disintegrin and metalloprotease with thrombospondin type 1 motif, member 13 (ADAMTS-13). It is hypothesized that in the case of COVID-19 infection, ADAMTS-13 regulation of VWF multimer distribution is impaired resulting in




Fig. 2 Thromboembolic complications in COVID-19 patients.^{16,26-36}




Fig. 3 Mechanism of CIC induced Thrombosis: The viral particles enter the host through ACE-2 receptors found on endothelial cells (EC) of respiratory pathway resulting in endothelial activation and dysfunction that leads to pyroptosis and cytokine storm. This in turn causes platelet activation and upregulation of tissue factor (TF) and fragment (1 + 2). The spiked inflammatory response along with downregulation of anticoagulant and fibrinolytic pathway leads to formation of thrombus.

procoagulant state.⁴⁴ Ilaria et al reported a significant change in the VWF-ADAMTS13 axis in infected patients, with an increase in VWF:Ag to ADAMTS13 activity ratio, which is strongly associated with disease severity. This imbalance in the ratio of VWF level to ADAMTS13 increases the hypercoagulable state of COVID-19 patients and thereby risk of microthrombosis.⁴⁵ Thus, this cascade of events cause endothelium to adopt a procoagulant phenotype, which is then supportive of platelet and leukocyte recruitment and marks the initiation of CIC.^{46,47}

IL-6, a crucial cytokine that is markedly increased in severe COVID-19, is an activator of coagulation that induces TF expression along with increased production of fibrinogen and platelets.⁴⁸⁻⁵⁰ Increased Angiotensin II (ATII) expression, along with proinflammatory cytokines and antiphospholipid antibodies are responsible for TF activation and overexpression in the infected patients. Thus, it is likely that the overexpression of TF in COVID-19 is related to the pathogenesis of the disease.⁵¹ The researchers have looked for correlation between TF expression and the severity of COVID-19 and found that TF expression both in monocytes and extracellular vesicles is associated with severity of disease.^{52,53} This is because TF can increase its expression during localized hypoxia of lungs in the case of COVID-19 infection and generate a thrombotic phenotype, which is observed to be regulated by extracellular RNA activated Toll-like receptor 3-activated protein 1 signaling.⁵⁴ This in turn initiates extrinsic cascade of coagulation and spiked inflammatory response. This inflammation further causes endothelitis in CIC.

These observations are in alignment with clinical autopsy of COVID-19 patients, as viral inclusion within the endothelial cells is seen in sections of lungs and kidney, thus highlighting the ability of virus to cause endothelial injury after infection. These findings are augmented with histological evidence of endothelial inflammation and cell death by viral load.³⁹ Together, these findings suggest a direct role of SARS-CoV-2 in promoting systemic microvascular dysfunction causing a prothrombotic milieu.⁴⁸

Role of Platelets in CIC Mediated Thrombosis

In severe cases of COVID-19, one of the common findings is thrombocytopenia. A recent meta-analysis reveals association of thrombocytopenia with 5-fold increase in risk of severe infection.⁴⁹ The recent studies demonstrated that severe COVID-19 cases often have cardiovascular comorbidities suggestive of thrombocytopenia or platelet apoptosis as a major contributor of virus pathogenesis.^{50,55} Other molecular events involved in SARS-CoV-2 pathogenesis, like inflammation, hypoxia, immune system activation, endothelial activation and dysfunction can lead to platelet apoptosis as well as activation, resulting in increased thrombotic events.⁵⁵⁻⁵⁸ Patient autopsies from those with severe SARS-CoV-2 infection show platelet-rich thrombotic microangiopathy.^{59,61} The presence of arterial and venous thrombus through-out the lungs is suggestive of CIC influenced platelet hyperactivation that in turn increases the severity of the disease by exacerbating inflammation. Platelets express a wide repertoire of adhesion as well as immune receptors, like

FcγRIIa, CXC/CCL receptors and TLRs along with an ability to engulf virions. SARS-CoV-2 has been demonstrated to activate platelets via TLR-7 and the FcγRIIa receptors, which evokes an array of functional responses that support thrombus formation, including formation of platelet leukocyte aggregates, platelet degranulation, and so on.^{62–65} Activated platelets expressing CD40L and P-selectin interact with neutrophils and release α-granules along with complement C3 and different cytokines like CC-chemokine ligand 2 (CCL2), CCL3, CCL7, IL-7, IL-8, IL-1β.^{66,67} These findings correlate with the clinical data of COVID-19 patients, as the level of these cytokines have been reported to be significantly elevated in these patients compared with healthy controls.⁶⁸ Previous studies with dengue virus show that platelets release IL-1β that results in increased permeability.⁶⁹ The release of IL-1β and other cytokines by platelets in response to SARS-CoV-2 possibly causes COVID-19 associated ARDS. As seen in COVID-19 pathology, parallel to cytokine release, another important event in ARDS, is recruitment of neutrophils to the pulmonary vasculature.⁷⁰ The molecular mechanism by which activated platelets binds to neutrophils and lead to rolling of platelet-bound neutrophils on the endothelium, is termed as 'secondary capture'. This plays a pivotal role in the initiation of immune-thrombosis.⁷¹ The transmigration of the platelets to the alveolar lumen is mediated by binding of activated platelets to neutrophils and results in the formation of edematous lungs, which leads to further platelet activation.⁷⁰ This series of events leads to formation of neutrophil extracellular traps (NETs). Recent reports suggest that NETs trigger thrombo-inflammation in COVID-19 patients,⁷⁰ causing vascular thrombosis and high risk of mortality.⁷² Taken together, platelet activation along with apoptosis are contributors to the severity of pathology of COVID-19, which includes thrombosis and the cytokine storm.

Along with mediating T cell function and leukocyte aggregation, platelets also increase B cell immunoglobulin (Ig) G1, IgG2, and IgG3 production.⁷³ This finding is significant in the light of COVID-19 as formation of antiphospholipid antibodies, like anti-β2 glycoprotein IgG antibodies, are associated with thrombosis in patients with COVID-19.⁷⁴ A cohort studied from France has demonstrated that a large percent of severe COVID-19 patients exhibits a detectable lupus anticoagulant.⁷⁵ However, it still remains to be established whether the presence of antiphospholipid antibodies in SARS-CoV-2 patients plays a role in infection-associated thrombosis or represents merely an association. Nevertheless, deciphering the ways platelets modulate the adaptive immune response to COVID-19 and the role played by adaptive immunity mediating thrombosis will be crucial in uncovering novel therapeutic strategies to combat the disease.

Prediction of CIC Severity and Occurrence of Thrombotic Events

Analysis of Coagulation and Inflammatory Parameters

There are 6 baseline coagulation tests that should be performed to measure the extent of CIC in infected patients.

These are PT, aPTT, D-dimer, fibrinogen, fibrin degradation products (FDP) and platelet count. These tests represent traditional tests used to help assess bleeding or thrombotic risk. Additional laboratory measures to evaluate CIC associated inflammation are C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), ferritin, procalcitonin, and high-sensitivity cardiac troponin. Inflammatory markers such as CRP, ESR are elevated in COVID-19 patients. These markers are also associated with the occurrence of thrombosis. We discuss below the potential utility of the 6 baseline coagulation tests that can be used to help assess the severity of disease and predict the future course of medical intervention.

- PT: In the initial stage of the disease, PT is normal or near-normal in most COVID-19 patients. Only 5% cases showed prolonged PT during the initial manifestation of the disease.¹⁶ However, there is significant change in the PT profile in critically ill COVID-19 patients having CIC.^{70,71} Critically ill patients show an average increase 1.9s in PT as compared with non-fatal or critical cases. Also, as the severity of the disease progress, ~48% of critically ill patient exhibit marked and progressive increase in PTs.¹⁹ Thus, an upward trending PT can add clinical evaluation in monitoring progression of the disease, in particular fatal or critical cases. That is, progressive increase in PTs can be considered a warning sign and a predictor of increased risk of mortality.
- aPTT: the aPTT profile is often unaltered in COVID-19 patients, and only a fraction (6%) of patients shows prolonged aPTT.¹⁶ There is no significant difference in the average aPTT between fatal and non-fatal cases of infection. Also, no significant correlation was observed between aPTT and disease severity or mortality.¹⁹ Therefore, the aPTT cannot be considered as reliable indicator of CIC and disease progression in COVID-19. Also, the aPTT cannot be associated with prevalence of VTE in COVID-19 cases.
- D-dimer: quantitative D-dimer is used routinely as a biomarker for exclusion/diagnosis and prediction of recurrence of VTE.^{76,77} 36% of reported cases have showed elevated levels of D-dimer (approximately, 0.9 mg/L).^{12,16} An elevated D-dimer level is observed in fatal cases as compared with non-fatal cases (mean level of 2.4 vs 0.5 mg/L) and is inversely correlated with the survival of patient.^{12,78} Additionally, non-survivors have shown steady and progressive elevation in D-dimer level, whereas in COVID-19 survivors the levels remain constant or improves.⁷⁹ Endothelial dysfunction caused by viral entry results in elevated levels of D-dimer along with thrombin and FDPs, inflammation and hypoxia resulting in CIC which can lead to pulmonary congestion mediated by thrombosis.⁷⁹ These findings suggest D-dimer to be highly prognostic in COVID-19. Evaluation of D-dimer level may also aid in identifying patients that could benefit from anticoagulant therapy.
- Fibrinogen: Most COVID-19 patients show elevated (4.55 g/L) fibrinogen in the initial stage of the disease.

The degree of elevation correlates with increased IL-6 levels but not with mortality.^{17,79} IL-6 is an important mediator of inflammation and in COVID-19, inflammation precedes coagulation. As the disease progresses, there may be a progressive decrease in the fibrinogen level, and at a late course of the disease is strongly associated with mortality. Approximately 29% of critically ill patients had fibrinogen <1 g/L, but this happens at the last stage of the disease.¹⁷ Thus, fibrinogen levels cannot be used for prognosis of the initial phase of COVID-19, but progressive deterioration may reflect a poor prognosis.

- FDPs: FDP levels are relatively constant in both mild cases and initial presentation of the disease. However, critically ill patients admitted to ICU showed significantly higher levels of FDP. The approximate levels of FDP in survivors are 4 µg/mL vs 7.6 µg/mL in the non-survivors. The elevation of D-dimer is followed by elevation of FSP. Though D-dimer is more sensitive marker for coagulopathy during the initial phase of the disease, FSP can also be considered prognostic for the COVID-19 infection as progressive elevation of FSP level correlates with mortality.¹⁷
- Platelet count: Platelet count is mostly unaltered or slightly reduced in most cases of COVID-19 and thrombocytopenia is reported in only 12–36% of total cases.^{12,16,19} Platelets play a crucial role in progression of coagulopathy and in the development of thrombosis (explained in the earlier section of the review). Severe thrombocytopenia correlates with progression of disease as more than 55% of critically ill COVID-19 patients have a platelet count of <100 × 10⁹/L.

Therapeutic Perspectives

The lack of specific therapy against SARS-CoV2 and significant association of thrombotic complications with severe COVID-19 has led to exploration of anti-thrombotics as a choice of drug for infected patients. Antithrombotic drugs

having anti-inflammatory and/or antiviral properties are also under clinical trial. Anticoagulants are first choice of antithrombotic therapy. VTE prophylaxis with either low molecular weight heparin or unfractionated heparin is commonly applied to COVID-19 patients.^{79,80} Initial clinical reports suggested that administration of low molecular weight heparin reduces the death rate in COVID-19 patients with an elevated sepsis-induced coagulopathy score or an elevated D-dimer level.^{81,82} Heparins, apart from being effective anticoagulants seem to have pleiotropic effects by way of their ability to bind to DAMPs, such as HMGB-1, and proinflammatory cytokines that may be additionally therapeutic in the context of viral infection, including anti-inflammatory effects.^{82,83} Recent findings are suggestive of other links between heparin and SARS-CoV-2, for example that the antiviral properties of heparin can be attributed to its ability to bind with the spike protein of the SARS-CoV-2 virus directly.⁸⁴ Similar results have been shown after docking of a major protease (PDB id 6y2e) of COVID-19 with heparin. It showed binding with affinity -11.8 Kcal/mol (–Fig. 4). But the use of heparins in COVID-19 cases is limited to individual case reports because of the potential for heparin therapeutics to cause heparin-induced thrombocytopenia.⁸⁵

Another therapeutic approach is the use of inhibitors of the contact factor activation pathway. Inhibition of thrombo-inflammatory response by inhibition of FXII also appears to be an attractive therapeutic target. Activation of FXII by negatively charged molecules like NETs and platelet derived PolyP initiates activation of contact pathway. The level of NETs and PolyP is reported to be upregulated in severe COVID-19 patients.⁸⁶ Activation of contact pathway causes thrombin generation and bradykinin activation. This in turn causes downstream complement activation and production of inflammatory cytokines.⁸⁷ It has been demonstrated in animal models of thrombosis that inhibition of FXIIa gives protection from occlusive thrombus formation without




Fig. 4 Binding of heparin with COVID-19 main protease: (A). Ligplot view of blind docking of main protease of COVID-19, (PDB: 6y2e) with heparin (Pubchem: 772). For a given protein-ligand docked PDB file, LigPlot automatically generates schematic diagrams as represented by hydrogen bonds and by hydrophobic contacts. Green dashed lines between the atoms involved represent the hydrogen bonds and an arc with orange spikes pointing toward the ligand atom they contact with represents the hydrophobic contacts. (B). Hydrophobicity surface view of protease bound to heparin: image as created by chimera represents “hydrophobicity surface” that is dodger blue for the most hydrophilic, to white, to orange red for the most hydrophobic.




Fig. 5 Flow chart for management of COVID-19 associated VTE. Abbreviations: Prothrombin time (PT), Platelet count (PC), Intensive care unit (ICU), Low molecular weight heparin (LMWH), Un-fractionated heparin (UFH).

impeding hemostasis.⁸⁸ The potential safety of FXII inhibitors is attributed to the fact that administered individuals display no bleeding disorder, and there is no reported effect on immune function.⁸⁹ Therefore, a FXII inhibitory antibody can be a promising novel combination of antithrombotic and anti-inflammatory drugs that can be used in COVID-19. Another therapeutic approach that may help in COVID-19 includes fibrinolytic agents such as tPA (tissue-type plasminogen activator). This interest in tPA has emerged from the concept that ARDS is marked by significant local inflammatory reaction along with a hypofibrinolytic state. As described earlier, severe COVID-19 may cause ARDS that can result in microvascular thrombosis, extensive thrombin generation and fibrin formation, because of mass upregulation of inflammatory cytokines and leukocytes.⁹⁰ In reports of its use in severe infection case, an initial improvement was observed after tPA infusion.⁹¹⁻⁹³ Antiplatelet therapies are also potentially promising in managing COVID-19 associated thrombosis. The majority of clinical findings associating antiplatelet therapy with improved outcomes relates to aspirin.⁹⁴ Recent experimental data has shown that aspirin prevents neutrophil mediated microvascular thrombosis and intravascular coagulation by targeting platelets in a mouse model of bacterial sepsis.⁹⁵ Still, the effect of aspirin in assisting in COVID-19 symptoms remains to be estab-

lished. Various other antiplatelet therapeutics, including dipyridamole and nafamostat, are currently being accessed for their potential to reduce the severity of COVID-19. Recently it has been reported that dipyridamole suppresses SARS-CoV-2 replication in vitro. The earlier studies with same drug suggest that use of adjunct dipyridamole may improve the clinical course in severe cases of COVID-19.⁹⁶ Nafamostat, a serine protease inhibitor that is marketed in Asia for the treatment of DIC and pancreatitis is being evaluated for its role in helping with COVID-19 associated coagulopathy because of its antiplatelet effect.⁹⁷ **Fig. 5** depicts a flow chart for management of COVID-19 associated VTE. With all the available evidence-based guidelines for the treatment of COVID-19, it can be said that the current therapies are inadequate as they do not substantially reduce morbidity and mortality. However, with the new mechanistic insights emerging from the concerted efforts of biomedical research, more opportunities will arise to identify potentially safe and effective therapeutic strategies for COVID-19.

Conclusion

This pandemic has jeopardized the security and stability of the entire human race for an unknown future duration,

plunging it into uncharted territory, leaving all of us feeling powerless in the face of an infectious and invisible threat. The emergence of the virus has presented unprecedented health-care and economic challenges globally. The decoding of pathological manifestations of this virus becomes paramount in the management of this disease. A crucial clinical feature of severe COVID-19 infection is presence of prothrombotic milieu, which is associated with increased rates of arterial, venous and microvascular thrombosis, along with adverse clinical outcomes. Emerging evidence from the experimental findings reveal that SARS-CoV-2 can infect endothelial cells and invoke immune response. This results in activation of inflammatory pathways causing endothelium dysregulation, leukocyte activation, NET generation, complement deposition, and thrombocytopenia. All these events together conspire to unleash a prothrombotic state (immune-thrombosis) that leads to significant thrombotic complications. Nevertheless, uncovering the precise mechanisms of SARS-CoV-2 infection and associated thrombotic complications is required to infer novel therapeutic approaches. Thus, in the context of this pandemic, the intersection of adaptive and innate immunity with inflammation and thrombosis has been thrust into the global spotlight. With the growing experimental evidences and understanding of thrombotic complications in SARS-CoV-2 infection, more rigorous diagnostic approaches resulting in early detection of thrombotic events can be developed. Also, the use of antithrombotic therapy for the prevention and treatment of COVID-19-associated thrombosis will help in improving disease outcomes in COVID-19 patients.

Funding

Ministry of Ayush, Government of India: S-14013/2018.

Conflict of Interest

None declared.

Acknowledgments

The authors would like to thank Jamia Millia Islamia, India for providing necessary support.

References

- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395(10223):497–506
- World Health Organization. Coronavirus Disease 2019 (COVID-19) Situation report - 46. (Accessed on March 12 2020) Online: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200306-sitrep-46-covid-19.pdf?sfvrsn=96b04adf_2
- Xiong TY, Redwood S, Prendergast B, Chen M. Coronaviruses and the cardiovascular system: acute and long-term implications. *Eur Heart J* 2020;41(19):1798–1800
- Zhu N, Zhang D, Wang W, et al; China Novel Coronavirus Investigating and Research Team. A novel coronavirus from patients with pneumonia in China. 2019. *N Engl J Med* 2020;382(08):727–733. Doi: 10.1056/NEJMoa2001017
- Cucinotta D, Vanelli M. WHO declares COVID-19 a Pandemic. *Acta Biomed* 2020;91(01):157–160. Doi: 10.23750/abm.v91i1.93973
- Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* 2020;181(02):281–292.e6
- Zhang H, Pennington JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* 2020;46(04):586–590
- van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med* 2020;382(16):1564–1567
- Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579(7798):270–273
- Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*; 2020
- Guan WJ, Ni ZY, Hu Y, et al; China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382(18):1708–1720
- Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the Coronavirus Disease 2019 (COVID-19) Pandemic. *J Am Coll Cardiol* 2020; 12;75(18):2352–2371
- Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential Effects of Coronaviruses on the Cardiovascular System: A Review. *JAMA Cardiol* 2020;5(07):831–840
- Levi M, Iba T. COVID-19 coagulopathy: is it disseminated intravascular coagulation? *Intern Emerg Med* 2021;16(02):309–312. Doi: 10.1007/s11739-020-02601-y
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;395(10223):507–513
- Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost* 2020;18(04):844–847
- Shi S, Qin M, Shen B, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiol* 2020;5(07):802–810 [epub ahead of print]. Doi: 10.1001/jamacardio.2020.0950
- Fan BE, Chong VCL, Chan SSW, et al. Hematologic parameters in patients with COVID-19 infection. *Am J Hematol* 2020
- Lew TW, Kwek TK, Tai D, et al. Acute respiratory distress syndrome in critically ill patients with severe acute respiratory syndrome. *JAMA* 2003;290(03):374–380
- <https://covid19.who.int/>
- Yang J, Zheng Y, Gou X, et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. *Int J Infect Dis* 2020;94:91–95. Doi: 10.1016/j.ijid.2020.03.017
- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395(10229):1054–1062
- Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020;180(07):934–943
- Boonyawat K, Chantrathammachart P, Numthavaj P, et al. Incidence of thromboembolism in patients with COVID-19: a systematic review and meta-analysis. *Thromb J* 2020;18(01):34. Doi: 10.1186/s12959-020-00248-5
- Dolhnikoff M, Duarte-Neto AN, de Almeida Monteiro RA, et al. Pathological evidence of pulmonary thrombotic phenomena in severe COVID-19. *J Thromb Haemost* 2020;18(06):1517–1519
- Klok FA, Kruijff MJHA, van der Meer NJM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020;191:145–147

- 27 Llitjos JF, Leclerc M, Chochois C, et al. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J Thromb Haemost* 2020;18(07):1743–1746
- 28 Middeldorp S, Coppens M, van Haaps TF, et al. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost* 2020;18(08):1995–2002
- 29 Marone EM, Rinaldi LF. Upsurge of deep venous thrombosis in patients affected by COVID-19: Preliminary data and possible explanations. *J Vasc Surg Venous Lymphat Disord* 2020;8(04):694–695
- 30 Wichmann D, Sperhake JP, Lütgehetmann M, et al. Autopsy Findings and Venous Thromboembolism in Patients With COVID-19: A Prospective Cohort Study. *Ann Intern Med* 2020;173(04):268–277
- 31 Zhang L, Feng X, Zhang D, et al. Deep Vein Thrombosis in Hospitalized Patients with Coronavirus Disease 2019 (COVID-19) in Wuhan, China: Prevalence, Risk Factors, and Outcome. *Circulation*; 2020
- 32 Ren B, Yan F, Deng Z, et al. Extremely High Incidence of Lower Extremity Deep Venous Thrombosis in 48 Patients With Severe COVID-19 in Wuhan. *Circulation* 2020;142(02):181–183
- 33 Stoneham SM, Milne KM, Nuttall E, et al. Thrombotic risk in COVID-19: a case series and case-control study. *Clin Med (Lond)* 2020;20(04):e76–e81
- 34 Voicu S, Bonnin P, Stépanian A, et al. High prevalence of deep vein thrombosis in mechanically ventilated COVID-19 patients. *J Am Coll Cardiol* 2020;76(04):480–482
- 35 Desborough MJR, Doyle AJ, Griffiths A, Retter A, Breen KA, Hunt BJ. Image-proven thromboembolism in patients with severe COVID-19 in a tertiary critical care unit in the United Kingdom. *Thromb Res* 2020;193:1–4
- 36 Fraissé M, Logre E, Pajot O, Mentec H, Plantefève G, Contou D. Thrombotic and hemorrhagic events in critically ill COVID-19 patients: a French monocenter retrospective study. *Crit Care* 2020;24(01):275
- 37 Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LFP. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol* 2020;20(06):363–374. Doi: 10.1038/s41577-020-0311-8
- 38 Moreno-Eutimio MA, López-Macías C, Pastelin-Palacios R. Bioinformatic analysis and identification of single-stranded RNA sequences recognized by TLR7/8 in the SARS-CoV-2, SARS-CoV, and MERS-CoV genomes. *Microbes Infect* 2020;22(4-5):226–229. Doi: 10.1016/j.micinf.2020.04.009
- 39 Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet* 2020;395(10234):1417–1418. Doi: 10.1016/S0140-6736(20)30937-5
- 40 Crackower MA, Sarao R, Oudit GY, et al. Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature* 2002;417(6891):822–828. Doi: 10.1038/nature00786
- 41 Danilczyk U, Penninger JM. Angiotensin-converting enzyme II in the heart and the kidney. *Circ Res* 2006;98(04):463–471. Doi: 10.1161/01.RES.0000205761.22353.5f
- 42 Kayal S, Jaïs JP, Aguiní N, Chaudière J, Labrousse J. Elevated circulating E-selectin, intercellular adhesion molecule 1, and von Willebrand factor in patients with severe infection. *Am J Respir Crit Care Med* 1998;157(3 Pt 1):776–784. Doi: 10.1164/ajrccm.157.3.9705034
- 43 Aird WC. The role of the endothelium in severe sepsis and multiple organ dysfunction syndrome. *Blood* 2003;101(10):3765–3777. Doi: 10.1182/blood-2002-06-1887
- 44 Ward SE, Fogarty H, Karampini E, et al; Irish COVID-19 Vasculopathy Study (iCVS) investigators. ADAMTS13 regulation of VWF multimer distribution in severe COVID-19. *J Thromb Haemost* 2021;19(08):1914–1921. Doi: 10.1111/jth.15409
- 45 Mancini I, Baronciani L, Artoni A, et al. The ADAMTS13-von Willebrand factor axis in COVID-19 patients. *J Thromb Haemost* 2021;19(02):513–521. Doi: 10.1111/jth.15191
- 46 Massberg S, Enders G, Leiderer R, et al. Platelet-endothelial cell interactions during ischemia/reperfusion: the role of P-selectin. *Blood* 1998;92(02):507–515
- 47 Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* 1994;76(02):301–314. Doi: 10.1016/0092-8674(94)90337-9
- 48 Monteil V, Kwon H, Prado P, et al. Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. *Cell* 2020;181(04):905–913.e7. Doi: 10.1016/j.cell.2020.04.004
- 49 Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A meta-analysis. *Clin Chim Acta* 2020;506:145–148. Doi: 10.1016/j.cca.2020.03.022
- 50 Yang X, Yang Q, Wang Y, et al. Thrombocytopenia and its association with mortality in patients with COVID-19. *J Thromb Haemost* 2020;18(06):1469–1472. Doi: 10.1111/jth.14848
- 51 Cañas CA, Cañas F, Bautista-Vargas M, Bonilla-Abadía F. Role of Tissue Factor in the Pathogenesis of COVID-19 and the Possible Ways to Inhibit It. *Clin Appl Thromb Hemost* 2021;27:10760296211003983. Doi: 10.1177/10760296211003983
- 52 Hottz ED, Azevedo-Quintanilha IG, Palhinha L, et al. Platelet activation and platelet-monocyte aggregate formation trigger tissue factor expression in patients with severe COVID-19. *Blood* 2020;136(11):1330–1341 Google Scholar | Crossref | Medline
- 53 Rosell A, Havervall S, von Meijenfeldt F, et al. Patients With COVID-19 Have Elevated Levels of Circulating Extracellular Vesicle Tissue Factor Activity That Is Associated With Severity and Mortality—Brief Report. *Arterioscler Thromb Vasc Biol* 2021;41(02):878–882
- 54 Bhagat S, Biswas I, Ahmed R, Khan GA. Hypoxia induced up-regulation of tissue factor is mediated through extracellular RNA activated Toll-like receptor 3-activated protein 1 signalling. *Blood Cells Mol Dis* 2020;84:102459
- 55 Tang WH, Martin KA, Hwa J. Aldose reductase, oxidative stress, and diabetic mellitus. *Front Pharmacol* 2012;3:87
- 56 Tang WH, Stitham J, Gleim S, et al. Glucose and collagen regulate human platelet activity through aldose reductase induction of thromboxane. *J Clin Invest* 2011;121(11):4462–4476
- 57 Tang WH, Stitham J, Jin Y, et al. Aldose reductase-mediated phosphorylation of p53 leads to mitochondrial dysfunction and damage in diabetic platelets. *Circulation* 2014;129(15):1598–1609
- 58 Sinauridze EI, Kireev DA, Popenko NY, et al. Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thromb Haemost* 2007;97(03):425–434
- 59 Ackermann M, Verleden SE, Kuehnel M, et al. Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in Covid-19. *N Engl J Med* 2020;383(02):120–128
- 60 Barton LM, Duval EJ, Stroberg E, Ghosh S, Mukhopadhyay S. COVID-19 Autopsies, Oklahoma, USA. *Am J Clin Pathol* 2020;153(06):725–733
- 61 Fox SE, Akmatbekov A, Harbert JL, Li G, Quincy Brown J, Vander Heide RS. Pulmonary and cardiac pathology in African American patients with COVID-19: an autopsy series from New Orleans. *Lancet Respir Med* 2020;8(07):681–686
- 62 Koupenova M, Clancy L, Corkrey HA, Freedman JE. Circulating platelets as mediators of immunity, inflammation, and thrombosis. *Circ Res* 2018;122(02):337–351. Doi: 10.1161/CIRCRESAHA.117.310795
- 63 Assinger A. Platelets and infection - an emerging role of platelets in viral infection. *Front Immunol* 2014;5:649. Doi: 10.3389/fimmu.2014.00649
- 64 Koupenova M, Corkrey HA, Vitseva O, et al. The role of platelets in mediating a response to human influenza infection. *Nat Commun* 2019;10(01):1780. Doi: 10.1038/s41467-019-09607-x
- 65 Koupenova M, Vitseva O, MacKay CR, et al. Platelet-TLR7 mediates host survival and platelet count during viral infection in the

- absence of platelet-dependent thrombosis. *Blood* 2014;124(05):791–802. Doi: 10.1182/blood-2013-11-536003
- 66 Fitch-Tewfik JL, Flaumenhaft R. Platelet granule exocytosis: a comparison with chromaffin cells. *Front Endocrinol* 2013;4:77
- 67 Sut C, Tariket S, Aubron C, et al. The non-hemostatic aspects of transfused platelets. *Front Med (Lausanne)* 2018;5:42
- 68 Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19. *J Infect* 2020;80(06):607–613
- 69 Hottz ED, Lopes JF, Freitas C, et al. Platelets mediate increased endothelium permeability in dengue through NLRP3-inflammatory activation. *Blood* 2013;122(20):3405–3414
- 70 Middleton EA, He XY, Denorme F, et al. Neutrophil extracellular traps contribute to immunothrombosis in COVID-19 acute respiratory distress syndrome. *Blood* 2020;136(10):1169–1179
- 71 Zarbock A, Polanowska-Grabowska RK, Ley K. Platelet-neutrophil-interactions: linking hemostasis and inflammation. *Blood Rev* 2007;21(02):99–111
- 72 Wang T, Chen R, Liu C, et al. Attention should be paid to venous thromboembolism prophylaxis in the management of COVID-19. *Lancet Haematol* 2020;7(05):e362–e363
- 73 Cognasse F, Hamzeh-Cognasse H, Lafarge S, et al. Human platelets can activate peripheral blood B cells and increase production of immunoglobulins. *Exp Hematol* 2007;35(09):1376–1387. Doi: 10.1016/j.exphem.2007.05.021
- 74 Favaloro EJ, Henry BM, Lippi G. Is Lupus Anticoagulant a Significant Feature of COVID-19? A Critical Appraisal of the Literature. *Semin Thromb Hemost* 2022;48(01):55–71
- 75 Helms J, Tacquard C, Severac F, et al; CRICS TRIGGERSEP Group (Clinical Research in Intensive Care and Sepsis Trial Group for Global Evaluation and Research in Sepsis) High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med* 2020;46(06):1089–1098. Doi: 10.1007/s00134-020-06062-
- 76 Pulivarthi S, Gurrani MK. Effectiveness of d-dimer as a screening test for venous thromboembolism: an update. *N Am J Med Sci* 2014;6(10):491–499
- 77 Palareti G, Cosmi B, Legnani C, et al; DULCIS (D-dimer and Ultrasonography in Combination Italian Study) Investigators. D-dimer to guide the duration of anticoagulation in patients with venous thromboembolism: a management study. *Blood* 2014;124(02):196–203
- 78 Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020;18(05):1094–1099. Doi: 10.1111/jth.14817
- 79 Ranucci M, Ballotta A, Di Dedda U, et al. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost* 2020;18(07):1747–1751
- 80 Bikdeli B, Madhavan MV, Jimenez D, et al; Global COVID-19 Thrombosis Collaborative Group, Endorsed by the ISTH, NATAF, ESVM, and the IUA, Supported by the ESC Working Group on Pulmonary Circulation and Right Ventricular Function. COVID-19 and thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy, and follow-up: JACC State-of-the-art review. *J Am Coll Cardiol* 2020;75(23):2950–2973. Doi: 10.1016/j.jacc.2020.04.031
- 81 Connors JM, Levy JH. COVID-19 and its implications for thrombosis and anticoagulation. *Blood* 2020;135(23):2033–2040. Doi: 10.1182/blood.202006000
- 82 Wool GD, Miller JL. The Impact of COVID-19 Disease on Platelets and Coagulation. *Pathobiology* 2021;88(01):15–27. Doi: 10.1159/000512007
- 83 Arnold K, Xu Y, Sparkenbaugh EM, et al. Design of anti-inflammatory heparan sulfate to protect against acetaminophen-induced acute liver failure. *Sci Transl Med* 2020;12(535):eaav8075. Doi: 10.1126/scitranslmed.aav8075
- 84 Poterucha TJ, Libby P, Goldhaber SZ. More than an anticoagulant: Do heparins have direct anti-inflammatory effects? *Thromb Haemost* 2017;117(03):437–444. Doi: 10.1160/TH16-08-0620
- 85 Mycroft-West C, Su D, Elli S, et al. The 2019 coronavirus (SARS-cov-2) surface protein (spike) s1 receptor binding domain undergoes conformational change upon heparin binding. *bioRxiv* 2020:2020.2002.2029.971093
- 86 Riker RR, May TL, Fraser GL, Gagnon DJ, Bandara M, Zemrak W, SederDB. Heparin-induced thrombocytopenia with thrombosis in covid-19 adult respiratory distress syndrome. *Research and Practice in Thrombosis and Haemostasis*. n/a
- 87 Maas C, Renné T. Coagulation factor XII in thrombosis and inflammation. *Blood* 2018;131(17):1903–1909. Doi: 10.1182/blood-2017-04-569111
- 88 Renné T, Stavrou EX. Roles of factor XII in innate immunity. *Front Immunol* 2019;10:2011. Doi: 10.3389/fimmu.2019.02011
- 89 Larsson M, Rayzman V, Nolte MW, et al. A factor XIIa inhibitory antibody provides thromboprotection in extracorporeal circulation without increasing bleeding risk. *Sci Transl Med* 2014;6(222):222ra17. Doi: 10.1126/scitranslmed.3006804
- 90 Renné T, Schmaier AH, Nickel KF, Blombäck M, Maas C. In vivo roles of factor XII. *Blood* 2012;120(22):4296–4303. Doi: 10.1182/blood-2012-07-292094
- 91 Whyte CS, Morrow GB, Mitchell JL, Chowdary P, Mutch NJ. Fibrinolytic abnormalities in acute respiratory distress syndrome (ARDS) and versatility of thrombolytic drugs to treat COVID-19. *J Thromb Haemost* 2020;18(07):1548–1555. Doi: 10.1111/jth.14872
- 92 Wang J, Hajizadeh N, Moore EE, et al. Tissue plasminogen activator (tPA) treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): A case series. *J Thromb Haemost* 2020;18(07):1752–1755. Doi: 10.1111/jth.14828
- 93 Liu C, Ma Y, Su Z, et al. Meta-analysis of preclinical studies of fibrinolytic therapy for acute lung injury. *Front Immunol* 2018;9:1898. Doi: 10.3389/fimmu.2018.01898
- 94 Hardaway RM, Harke H, Tyroch AH, Williams CH, Vazquez Y, Krause GF. Treatment of severe acute respiratory distress syndrome: a final report on a phase I study. *Am Surg* 2001;67(04):377–382
- 95 Ouyang Y, Wang Y, Liu B, Ma X, Ding R. Effects of antiplatelet therapy on the mortality rate of patients with sepsis: A meta-analysis. *J Crit Care* 2019;50:162–168. Doi: 10.1016/j.jcrc.2018.12.004
- 96 Carestia A, Davis RP, Grosjean H, Lau MW, Jenne CN. Acetylsalicylic acid inhibits intravascular coagulation during *Staphylococcus aureus*-induced sepsis in mice. *Blood* 2020;135(15):1281–1286. Doi: 10.1182/blood.2019002783
- 97 Liu X, Li Z, Liu S, et al. Potential therapeutic effects of dipyrindamole in the severely ill patients with COVID-19. *Acta Pharm Sin B* 2020;10(07):1205–1215. Doi: 10.1016/j.apsb.2020.04.008