



# Prevalence of Proximal Contact Loss between Implant-Supported Prosthesis and Adjacent Natural Teeth: An Umbrella Review

Amirhossein Fathi<sup>1</sup> Ramin Mosharraf<sup>1</sup> Behnaz Ebadian<sup>2</sup> Mehdi Javan<sup>3</sup> Sabire Isler<sup>4</sup>  
Sara Nasrollahi Dezaki<sup>5</sup>

<sup>1</sup>Department of Prosthodontics, Dental Materials Research Center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>2</sup>Department of Prosthodontics, Dental Implants Research Center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

<sup>3</sup>Private Practice, Tehran, Iran

<sup>4</sup>Department of Prosthodontics, Istanbul University, Istanbul, Turkey

<sup>5</sup>Dental Students' Research Committee, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

**Address for correspondence** Amirhossein Fathi, DMD, MS, MSD, Dental Prosthodontics Department, Dental Materials Research Center, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran (e-mail: Amir\_alty@yahoo.com).

Eur J Dent 2022;16:742–748.

## Abstract

Contact loss between the implant prosthesis and adjacent natural teeth is a clinical complication whose overall prevalence is uncertain. Therefore, the main purpose of this umbrella study was to evaluate the extent of contact loss between implant prostheses and adjacent natural teeth. Electronic database of MEDLINE/PubMed, Cochrane, and Google Scholar was searched until August 2021 without considering language restrictions and according to Preferred Report Items for Systematic and Meta-Analysis guidelines (preferential reporting items for systematic review and meta-analysis). Inclusion criteria were systematic/meta-analysis review articles related to contact loss between implant prostheses and adjacent natural teeth. Inclusion criteria and risk of bias for the selected systematic/meta-analysis review studies were assessed by two or three qualified researchers, and the fourth researcher was used to resolve the ambiguities. From 43 eligible articles, five systematic/meta-analysis review studies were selected for this study. Important information such as the range of contact points, the prevalence, and the location of the contact loss was extracted. Three research studies had a low risk of bias and were considered clinical evidence. Analysis of low-risk studies showed that the superiority of open contact loss was excessive. Prevalence of proximal contact loss was more in mesial contact, especially in the mandibular arch. No significant differences were reported in sex or between the posterior and anterior regions.

## Keywords

- ▶ proximal contact loss
- ▶ implant-supported prostheses
- ▶ adjacent natural teeth

published online  
June 10, 2022

DOI <https://doi.org/10.1055/s-0042-1745771>.  
ISSN 1305-7456.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

## Introduction

The use of implant-supported protheses in patients suffering from edentulousness is a good treatment with a good prognosis.<sup>1,2</sup> Nevertheless, various biological and mechanical complications are reported in implant protheses<sup>3-7</sup>; one of the most important complications is interproximal contact loss (ICL) between the implant prosthesis and the adjacent natural tooth,<sup>8,9</sup> and numerous physiological factors play a role in the occurrence of this complication, including the type, location, and chewing forces. In fact, the most important issue regarding this complication is the mesial migration of natural teeth,<sup>10,11</sup> for which periodontal ligament is responsible.

Osseointegrated implant protheses are ankylosed and unmovable, which result in further contact loss.<sup>12,13</sup> However, there is no consensus on the underlying cause of ICL since it is not seen in all patients necessarily.<sup>14</sup> It is difficult to avoid ICL due to its various possible factors which may be progressive and require clinical intervention. To prevent the displacement of food particles, caries, and periodontal disease, modification of the prosthesis and restoration of adjacent natural teeth are suggested as ICL<sup>15-17</sup> treatment methods. Some studies have mentioned that ICL causes inflammation around the implant, resulting in marginal bone loss and implant failure.<sup>14,18</sup>

Considering the importance of the issue, this umbrella review investigates the incidence of ICL between implant protheses and nearby natural teeth. In this study, the null hypotheses were that “there is no correlation between implant-supported protheses and natural teeth adjacent to the ICL” and “there is no significant difference between the sex of the individual and the ICL position (mesial/distal, anterior/posterior, and maxilla/mandible).”

## Methods

### Search Strategy

Electronic searches were conducted in PubMed/MEDLINE, Cochrane, and Google Scholar until August 2021 without language restrictions. According to population, intervention, comparison, and outcome, the research question was “Is there a correlation between contact loss of the implant prosthesis and the nearby natural tooth?”. The “population” included patients with implant protheses. The “intervention” consisted of contact loss between implant protheses and adjacent natural teeth. “Comparison” was performed with individuals who had contact with the implant prosthesis and adjacent natural teeth. The primary “outcome” included the prevalence of ICL between implant-supported protheses and adjacent natural teeth; the secondary “outcome” included the incidence of ICL according to gender and positions. This review study was conducted using the Preferred Report Items for Systematic and Meta-Analysis guidelines.<sup>19</sup> The articles used included a systematic/meta-analysis review and resources that examined the incidence of contact loss among implants and adjacent natural teeth. The Assessment of Multiple Systematic Reviews (AMSR2) method was used to calculate the risk of systematic/meta-analysis review bias.<sup>20</sup> The database was searched based on medical

subject heading (mesh) and non-mesh keywords in simple or multiple conjunctions: (((dental implant [Title/Abstract]) OR (implant with support [Title/Abstract])) AND (loss of contact [Title/Abstract])) OR (open contact [Title/Abstract]) OR (adjacent natural teeth [Title/Abstract]).

### Inclusion and Exclusion Criteria in Screening

Inclusion criteria consisted of clinical studies in systematic/meta-analysis review and the existence of contact loss between implant-protheses and nearby natural teeth. Exclusion criteria included duplicate reviews, comments, and editorials. The studies were confirmed following receipt of the full text and observation of the inclusion and exclusion criteria.

### Data Collection Process

Two independent reviewers (R.M. and B.E.) qualified the eligible articles for review (1.0 Kappa). One researcher (R.M.) was responsible for extracting qualitative or quantitative data from the studies, and the second researcher (B.E.) was responsible for reviewing all collected data. Collected information included the author’s name, year and type of the study, the number of contacts, the incidence of contact loss, location, and type of prosthesis. Ambiguity and incompatibility were solved by resolving discussions. If a problem were unresolved, the third researcher helped. The initial search yielded 43 articles, of which 17 remained after removing duplicates and irrelevant ones by consensus. Five studies were found eligible eventually.

### Bias Risk Assessment

Based on the risk of bias assessment, to assess the quality of systematic/meta-analysis review studies, we used 16 questions of AMSR2<sup>20</sup> (► **Table 1**). In the end, each article received a score that indicated the risk of bias in that study. With eight to eleven positive responses, the risk of bias decreased; if four to seven questions were answered positively, the risk of bias was moderate and if fewer than three questions received a positive response, the risk of bias was considered as high.<sup>21</sup> Three qualified investigators assessed the articles (kappa = 0.9). Ambiguity and incompatibility were followed by resolving discussions. If a problem remained unresolved, the fourth researcher assisted.

## Results

### Screening of Systematic/Meta-analysis Reviews

A search in PubMed/MEDLINE databases (17 articles), Embase (three articles), Google Scholar (zero), and the Cochrane Library (23 articles) resulted in finding 43 articles. After removing duplicate sources, 41 studies remained for reviewing the titles and abstracts. After carefully studying the titles/abstracts of the articles, 12 articles met the eligibility criteria. Seven of them<sup>15-17,22-25</sup> were excluded due to the reasons indicated in ► **Table 2**, and a total of five studies were selected eventually,<sup>14,26-29</sup> which included 73 articles published between 2021 and 2016. Details of the research strategy are in ► **Fig. 1**. ► **Table 3** summarizes the most important features of these studies.

**Table 1** The Assessment of Multiple Systematic Reviews (AMSR2) tool

Systematic articles	1	1	3	5	6	7	8	9	11	14	2	4	10	12	13	15	16
	2	1	3	5	6	8	11	14	4	7	2	9	10	12	13	15	16
	3	1	2	3	5	6	10	11	13	14	16	4	7	8	9	12	15
	4	1	2	3	5	6	9	11	12	13	14	16	4	7	8	10	15
	5	3	16	1	2	4	5	6	7	8	9	10	14	11	12	13	15
AMSR2 items																	
Yes			Partial yes					No					NMC				

Abbreviation: NMC, no meta-analysis conducted

Note: Overall methodological quality: low: 0–5, moderate: 5–10, and high: 11–16.

Note: Criteria for AMSTAR analysis according to positives answers: low risk (8–11), moderate risk (4–7), and high risk ( $\leq 3$ ).

**Table 2** Excluded studies and reasons for exclusion

	References	Reasons for exclusion
22	Kim et al 2019	Clinical method guide
16	Liu et al 2019	Clinic report
17	Sfondouris and Prestipino 2019	Clinic report
23	Zeng et al 2018	Case control study
24	Luo et al 2016	Non-English
25	Ren et al 2016	Non-systematic review
15	Wat et al 2017	Clinic report

### Risk of Bias Assessment

The risk of bias was measured using the AMSR2 tool. This tool uses for a variety of different studies. Based on the number of correct responses, the level of bias in the study was reported as high, medium, or low (–Table 1). In this study, the risk of bias was either moderate (including one systematic/meta-analysis review<sup>27</sup>) or low (including three systematic/meta-analysis reviews<sup>26,28,29</sup>). Systematic/meta-analysis review studies with a low risk of bias were considered as clinical evidence. The low-risk systematic/meta-analysis review accounted for 72.6% of the study volume (–Table 3).

### Characteristics of Systematic Reviews

General information on each systematic/meta-analysis review is presented in –Table 3 (authors and year of publication, number, and type of studies, type of analysis, research period, interventions, outcomes, risk of bias, and main results).

### General Sample Analysis

Generally, we analyzed five review studies that consisted of 73 articles (43 retrospective, 16 prospective, and 14 RTC articles) and 29,765 (501 to 12370) proximal contact points in total. All reviews<sup>14,26–29</sup> reported a high prevalence of ICL; Oh et al<sup>28</sup> showed that the ratio of ICL in the implants was 2.5 times higher than that of the teeth and was increased in proximal space in follow-up periods; Manicone et al<sup>27</sup> showed that ICL is a common problem occurring in 29% of contact points associated with an increase in inflammation

of adjacent teeth. One of the reviews<sup>26</sup> declared that the posterior/anterior regions and gender did not affect the prevalence. However, most reviews<sup>26,28,29</sup> stated high heterogeneity and the need to perform further randomized control trials and blinded observations.

Further examination of the studies showed that the incidence of ICL in the maxillary areas was similar to that of the mandible.<sup>30–37</sup> Besides, all studies presented more damage in the mesial regions. Some articles<sup>30,32–34,36</sup> compared ICL in the anterior and posterior regions and concluded that ICL mostly occurs in the posterior regions.<sup>32–34</sup> Most studies evaluated age, sex, and implant site and did not find any significant correlation between appearance of ICL and these factors.


### Discussion

This umbrella study examined the incidence of ICL between implant protheses and nearby natural teeth. The null hypotheses were not accepted because the results showed more ICL between implant-supported protheses and nearby natural teeth. There was also a significant difference in the position of the ICL (mesial/distal).

The main result of this umbrella review was a high incidence of ICL among natural teeth and implant protheses. Studies have also concluded that the prevalence of ICL increases more than 80% after 5 years.<sup>38</sup>

ICL is a common complication. According to Manicone et al, it occurs in approximately 30% of contact points. Open contact (OC) is annoying to the patient, causes more inflammation in nearby tissues, and can increase the risk of new defects.<sup>27</sup> The prevalence of proximal OC varied between studies.<sup>9,15,33,36,37</sup> Various studies have reported the first OC between 1 and 123 months after the restoration.<sup>33,36,37</sup> Wong et al indicated that the incidence of OC was similar among protheses repaired with screw or cement. As time goes on, the size of the space between the teeth and the implant restoration may increase<sup>15,33</sup> and the number of OC will increase over time.<sup>9,33,36,37</sup>

Studies have reported that the prevalence of ICL among different sexes and ages is almost the same, indicating that ICL is not limited to a specific age or gender.<sup>39</sup> These studies indicate that 50  $\mu$ m metal shim and flossing can affect the



**Fig. 1** Flow charts for the studies were identified, displayed, and included in the study.

prevalence of ICL. Possible causes of ICL include dental migration, artificial crown-related factors, and bone growth factors.<sup>31,33,34,37</sup>

The results of this umbrella study show that the mesial ICL is more prevalent than the distal. The possible cause of this condition is faster wear in the mesial than distal. Thus, the mesial naturally compensate for this wear through displacement. The clinical causes of ICL in the distal region are not well understood. In general, the results of studies indicate that osseointegrated implants are at risk of infra-occlusion and mesial/distal ICL obstruction due to the destruction of nearby teeth and bone growth in the facial region.<sup>40</sup>

Some studies have reported a higher prevalence of ICL in the mandible than in the maxilla.<sup>14,37</sup> This may be due to the tendency of the mandibular teeth to mesial and, thus, increase the likelihood of developing ICL.<sup>33</sup> Studies have reported a similar incidence of ICL in the anterior/posterior regions. In patients with a high Frankfort-mandibular plane angle, the anterior force components are high and increase the prevalence of ICL. According to the anterior force theory, when a force applies to the posterior teeth, this force travels through the proximal contacts to reach the midline. There-

fore, this force transmission causes wear in the proximal contact of the anterior and posterior teeth.

ICL can be a high-risk complication in implant-supported restorations because it causes marginal bone resorption.<sup>30,32</sup> The results suggest that patients should be aware of the increased risk of ICL among natural teeth and implant-supported protheses over time.<sup>32,34</sup> The causes of ICL are not fully understood, so more research is needed. One of the limitations of this umbrella study is the high heterogeneity between articles, such as the type of study, non-randomness, different methods, and durations of ICL evaluation which makes it difficult to determine the prevalence and evaluate the adverse effects of ICL.

## Conclusion

Based on the findings of this umbrella study, the following general results are included.

ICL is very common and often occurs at the contact points in mesial and in the mandibular arch, and no significant differences were reported between ICL and gender or posterior/anterior regions.

**Table 3** Baseline characteristics of systematic reviews assessing the prevalence of proximal contact loss between implant-supported prostheses and adjacent natural teeth

Author (y)	Types/No. of studies included	Number of contact loss	Method of analysis	Search period	Population	Interventions	Comparison	Outcomes assessed	Risk of bias	Main results
Bento et al 2021 <sup>26</sup>	8 retrospective Two prospective	6,473	Systematic review and meta-analysis	Up to September 2020	Patients rehabilitated with implant-supported prostheses	The proximal contact loss between the implant-supported prostheses and adjacent natural teeth	Situations in which no proximal contact was lost between the implant-supported prostheses and adjacent natural teeth	The prevalence of proximal contact loss between implant prostheses and adjacent natural teeth, the prevalence of proximal contact loss in terms of sex and location	Low risk	The prevalence of proximal contact loss was high, occurring more frequently with the mesial contact and the mandibular arch. Significant differences were not found concerning sex or between the posterior and anterior region
Manicone et al 2021 <sup>27</sup>	10 retrospective Five prospective	12,370	Systematic review and meta-analysis	November 2020	Patients who had received or were scheduled to receive a single implant restoration had been rehabilitated or were scheduled to be rehabilitated with implant-supported fixed partial dentures	The overall prevalence of PCL determines the distribution and clinical features	Prevalence, the condition, context, and population framework	The number of cases of PCL that occurred, the number of PCL that occurred at the mesial contact point, at the distal contact point, in the mandible, and the maxilla	Moderate risk	PCL is a frequent complication. Approximately 29% of contact points develop this condition, which may cause food impaction and damage to the interproximal tissues
Oh et al, 2020 <sup>28</sup>	11 retrospective Five prospective	2,757	Systematic review and meta-analysis	March 2020	Patients with implant-supported prostheses or tooth-supported prostheses	Open proximal contact with implant-supported fixed prostheses compared with tooth-supported fixed prostheses	Odds of developing OPC with implant-supported prostheses or tooth-supported prostheses	Odds of OPC between implant-supported fixed prostheses or adjacent teeth	Low risk	The odds of developing OPC were significantly higher with implant-supported prostheses than with tooth-supported prostheses
Papageorgio et al 2018 <sup>29</sup>	Nine RCTs 14 retrospective Four prospective	7,664	Systematic review and meta-analysis	June 2018	Human patients of any age, sex, or ethnicity with at least one osseointegrated dental implant placed (including its restoration) among natural teeth	Frequency of infra-position and missing contact points in implant-supported restorations within natural dentitions over time	The influence of various patient, implant, or study-related characteristics	The IIP of the osseointegrated implant (and its supra-structure) compared with adjacent teeth, to loss of the PCP of the implant crown with the adjacent natural tooth	Low risk	Long-term adverse effects of dental implants among natural teeth can be observed in terms of IIP and PCP loss to the adjacent teeth
Greenstein et al 2016 <sup>14</sup>	Five RCTs	501	Review	2015	Patients with implant-supported prostheses or tooth-supported prostheses	Evaluate the potential causes, clinical significance, and treatment of open contacts between dental implant restorations and adjacent natural teeth	Percentage of restored dental implants manifesting open proximal contact areas adjacent to natural teeth	The incidence of open contacts that develop after implant restorations occurs next to teeth	High risk	The occurrence of an interproximal separation next to an implant restoration was greater than anticipated

Abbreviations: IIP, implant infra-position; OPC, open proximal contact; PCL, proximal contact loss; PCP, proximal contact point; RCT, randomized control trial.

## Conflict of Interest

None declared.

## References

- 1 Pjetursson BE, Brägger U, Lang NP, Zwahlen M. Comparison of survival and complication rates of tooth-supported fixed dental prostheses (FDPs) and implant-supported FDPs and single crowns (SCs). *Clin Oral Implants Res* 2007;18(Suppl 3):97–113
- 2 Blanes RJ, Bernard JP, Blanes ZM, Belser UC. A 10-year prospective study of ITI dental implants placed in the posterior region. II: influence of the crown-to-implant ratio and different prosthetic treatment modalities on crestal bone loss. *Clin Oral Implants Res* 2007;18(06):707–714
- 3 Pjetursson BE, Lang NP. Prosthetic treatment planning on the basis of scientific evidence. *J Oral Rehabil* 2008;35(Suppl 1):72–79
- 4 Kreissl ME, Gerds T, Muche R, Heydecke G, Strub JR. Technical complications of implant-supported fixed partial dentures in partially edentulous cases after an average observation period of 5 years. *Clin Oral Implants Res* 2007;18(06):720–726
- 5 Kim Y, Oh TJ, Misch CE, Wang HL. Occlusal considerations in implant therapy: clinical guidelines with biomechanical rationale. *Clin Oral Implants Res* 2005;16(01):26–35
- 6 Tallarico M, Caneva M, Baldini N, et al. Patient-centered rehabilitation of single, partial, and complete edentulism with cemented- or screw-retained fixed dental prosthesis: the First Osstem Advanced Dental Implant Research and Education Center Consensus Conference 2017. *Eur J Dent* 2018;12(04):617–626
- 7 Ferreira PW, Nogueira PJ, de Araújo Nobre MA, Guedes CM, Salvado F. Impact of mechanical complications on success of dental implant treatments: a case-control study. *Eur J Dent* 2022;16(01):179–187
- 8 Jo D-W, Kwon M-J, Kim J-H, Kim Y-K, Yi Y-J. Evaluation of adjacent tooth displacement in the posterior implant restoration with proximal contact loss by superimposition of digital models. *J Adv Prosthodont* 2019;11(02):88–94
- 9 Wong AT, Wat PY, Pow EH, Leung KC. Proximal contact loss between implant-supported prostheses and adjacent natural teeth: a retrospective study. *Clin Oral Implants Res* 2015;26(04):e68–e71
- 10 Herber R-P, Fong J, Lucas SA, Ho SP. Imaging an adapted dentoalveolar complex. *Anat Res Int* 2012;2012:782571
- 11 Wolpoff MH. Interstitial wear. *Am J Phys Anthropol* 1971;34(02):205–227
- 12 Odman J, Gröndahl K, Lekholm U, Thilander B. The effect of osseointegrated implants on the dento-alveolar development. A clinical and radiographic study in growing pigs. *Eur J Orthod* 1991;13(04):279–286
- 13 Bernard JP, Schatz JP, Christou P, Belser U, Kiliaridis S. Long-term vertical changes of the anterior maxillary teeth adjacent to single implants in young and mature adults. A retrospective study. *J Clin Periodontol* 2004;31(11):1024–1028
- 14 Greenstein G, Carpentieri J, Cavallaro J. Open contacts adjacent to dental implant restorations: etiology, incidence, consequences, and correction. *J Am Dent Assoc* 2016;147(01):28–34
- 15 Wat PY, Wong AT, Leung KC, Pow EH. Proximal contact loss between implant-supported prostheses and adjacent natural teeth: a clinical report. *J Prosthet Dent* 2011;105(01):1–4
- 16 Liu X, Liu J, Zhou J, Tan J. Closing open contacts adjacent to an implant-supported restoration. *J Dent Sci* 2019;14(02):216–218
- 17 Sfondouris T, Prestipino V. Chairside management of an open proximal contact on an implant-supported ceramic crown using direct composite resin. *J Prosthet Dent* 2019;122(01):1–4
- 18 Varthis S, Tarnow DP, Randi A. Interproximal open contacts between implant restorations and adjacent teeth. Prevalence—causes—possible solutions. *J Prosthodont* 2019;28(02):e806–e810
- 19 Asar S, Jalalpour S, Ayoubi F, Rahmani M, Rezaeian M. PRISMA; preferred reporting items for systematic reviews and meta-analyses. *Majallah-i Ilmi-i Danishgah-i Ulum-i Pizishki-i Rafsanjan* 2016;15(01):68–80
- 20 Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007;7(01):10
- 21 Silva V, Grande AJ, Carvalho AP, Martimbiano AL, Riera R. Overview of systematic reviews—a new type of study. Part II. *Sao Paulo Med J* 2015;133(03):206–217
- 22 Kim J-Y, Lim Y-J, Heo Y-K. Modification of framework design for an implant-retained fixed restoration helps when proximal contact loss occurs. *J Dent Sci* 2019;14(02):213–215
- 23 Zeng BJ, Guo Y, Yu RY. [Effect of the vacuum-formed retainer on preventing the proximal contact loss between implant supported crown and adjacent natural teeth]. *Beijing Da Xue Xue Bao* 2018;50(03):553–559
- 24 Luo Q, Ding Q, Zhang L, Peng D, Xie QF. [The loss of interproximal contact between posterior fixed implant prostheses and adjacent teeth: a retrospective study]. *Zhonghua Kou Qiang Yi Xue Za Zhi* 2016;51(01):15–19
- 25 Ren S, Lin Y, Hu X, Wang Y. Changes in proximal contact tightness between fixed implant prostheses and adjacent teeth: a 1-year prospective study. *J Prosthet Dent* 2016;115(04):437–440
- 26 Bento VA, Gomes JM, Lemos CA, Limirio JP, Rosa CD, Pellizzer EP. Prevalence of proximal contact loss between implant-supported prostheses and adjacent natural teeth: a systematic review and meta-analysis. *J Prosthet Dent* 2021;S0022-3913(21):00333–00334
- 27 Manicone PF, De Angelis P, Rella E, Papetti L, D'Addona A. Proximal contact loss in implant-supported restorations: a systematic review and meta-analysis of prevalence. *J Prosthodont* 2022;31(03):201–209
- 28 Oh W-S, Oh J, Valcanaia AJ. Open proximal contact with implant-supported fixed prostheses compared with tooth-supported fixed prostheses: a systematic review and meta-analysis. *Int J Oral Maxillofac Implants* 2020;30(06):e99–e108
- 29 Papageorgiou SN, Eliades T, Hämmerle CHF. Frequency of infraposition and missing contact points in implant-supported restorations within natural dentitions over time: a systematic review with meta-analysis. *Clin Oral Implants Res* 2018;29(Suppl 18):309–325
- 30 Saber A, Chakar C, Mokbel N, Nohra J. Prevalence of interproximal contact loss between implant-supported fixed prostheses and adjacent teeth and its impact on marginal bone loss: a retrospective study. *Int J Oral Maxillofac Implants* 2020;35(03):625–630
- 31 Pang NS, Suh CS, Kim KD, Park W, Jung BY. Prevalence of proximal contact loss between implant-supported fixed prostheses and adjacent natural teeth and its associated factors: a 7-year prospective study. *Clin Oral Implants Res* 2017;28(12):1501–1508
- 32 Varthis S, Randi A, Tarnow DP. Prevalence of interproximal open contacts between single-implant restorations and adjacent teeth. *Int J Oral Maxillofac Implants* 2016;31(05):1089–1092
- 33 Wei H, Tomotake Y, Nagao K, Ichikawa T. Implant prostheses and adjacent tooth migration: preliminary retrospective survey using 3-dimensional occlusal analysis. *Int J Prosthodont* 2008;21(04):302–304
- 34 French D, Naito M, Linke B. Interproximal contact loss in a retrospective cross-sectional study of 4325 implants: distribution and incidence and the effect on bone loss and peri-implant soft tissue. *J Prosthet Dent* 2019;122(02):108–114
- 35 Shi J-Y, Zhu Y, Gu YX, Lai HC. Proximal contact alterations between implant-supported restorations and adjacent natural teeth in the posterior region: a 1-year preliminary study. *Int J Oral Maxillofac Implants* 2019;34(01):165–168

- 36 Byun SJ, Heo SM, Ahn SG, Chang M. Analysis of proximal contact loss between implant-supported fixed dental protheses and adjacent teeth in relation to influential factors and effects. A cross-sectional study. *Clin Oral Implants Res* 2015;26(06):709–714
- 37 Koori H, Morimoto K, Tsukiyama Y, Koyano K. Statistical analysis of the diachronic loss of interproximal contact between fixed implant protheses and adjacent teeth. *Int J Prosthodont* 2010;23(06):535–540
- 38 Liang CH, Nien CY, Chen YL, Hsu KW. The prevalence and associated factors of proximal contact loss between implant restoration and adjacent tooth after function: a retrospective study. *Clin Implant Dent Relat Res* 2020;22(03):351–358
- 39 Oesterle LJ, Cronin RJ Jr. Adult growth, aging, and the single-tooth implant. *Int J Oral Maxillofac Implants* 2000;15(02):252–260
- 40 Chang M, Wennström JL. Longitudinal changes in tooth/single-implant relationship and bone topography: an 8-year retrospective analysis. *Clin Implant Dent Relat Res* 2012;14(03):388–394