



When to Suspect DRUJ's Instability in Children? Case Report of a Rare Presentation of Distal Forearm Fractures

Claire-Anne Saugy¹ Aline Bourgeois Bregou²

¹Service d'Orthopédie, Hôpital de zone de Morges, Morges, Switzerland

²Unité Pédiatrique de Chirurgie Orthopédique et Traumatologique, CHUV, Lausanne, Vaud, Switzerland

Address for correspondence Claire-Anne Saugy, M.D., Service d'Orthopédie, Hôpital de zone de Morges, Chemin du Crêt 2, 1110 Morges, Switzerland (e-mail: claire-anne.saugy@ehc.vd.ch).

European J Pediatr Surg Rep 2022;10:e73–e75.

Abstract

Pediatric displaced distal metaphyseal ulnar fractures and distal radial buckle fractures are common. However, to the best of our knowledge, their association has never been specifically reported. Thus, classification and management of this pattern remain challenging especially in young children. Distal radioulnar joint (DRUJ)'s assessment is difficult. A Galeazzi-equivalent injury should be suspected. We report the case of a 2-year-old boy who presented the above-mentioned association of forearm fractures and compare our management with actual recommendations. We would recommend a low-suspicion threshold for DRUJ's instability in young children presenting with displaced distal metaphyseal ulnar fracture associated with distal radial buckle fracture. That suspicion should raise the necessity of appropriate treatment and follow-up.

Keywords

- ▶ Forearm fracture
- ▶ Children
- ▶ distal radioulnar joint
- ▶ Galeazzi-equivalent injury

New Insights & the Importance for the Pediatric Surgeon

DRUJ's assessment is required in young children presenting unusual distal forearm fractures (e.g., displaced distal metaphyseal ulnar fracture associated with distal radial buckle fracture). It may impact treatment and follow-up.

Introduction

Distal forearm fractures are the most common fractures among children.^{1,2} Several injury patterns are possible: torus/buckle, greenstick, complete, displaced or undisplaced metaphyseal fractures, physeal fractures, and Galeazzi-equivalent injuries.^{3,4} The radius is predominantly concerned.² Depending on the residual growth and the remodeling capacity of the growing bone, treatment modalities include cast immobilization with or without closed reduction, and closed or open reduction with fixation in the most severe injuries.¹ However, to the best of our knowledge, the following pattern has never been described: displaced meta-

physeal ulnar fracture associated with distal radial buckle fracture. Thus, classification and management of this pattern remain challenging. Stability of distal radioulnar joint (DRUJ) must be maintained.

Case Report

We report the case of a 2-year-old boy who fell from the height of 30 cm and sustained a right forearm trauma. No adult witnessed the fall. Initial workup showed a painful deformation of the right wrist without neurovascular disorder and skin lesion. X-rays revealed a displaced metaphyseal ulnar fracture with a 30-degree posterior tilt associated with

received
September 15, 2021
accepted after revision
March 2, 2022

DOI <https://doi.org/10.1055/s-0042-1748317>.
ISSN 2194-7619.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany




Fig. 1 Serial X-rays of the patient.

a distal radial buckle fracture (→Fig. 1). Pediatric orthopaedic surgeons performed a closed reduction under general anesthesia and an above elbow plaster cast immobilization, with small palmar flexion and forearm neutral position (→Fig. 1). It remained for 4 weeks and was then replaced by a wrist brace for 4 more weeks.

Clinical and radiological evolution were good (→Fig. 1). Two months after surgery, the child had no residual pain, no range of motion limitation, and no DRUJ's instability. Activities with risk of falling were then allowed. Follow-up was discontinued 5 months after surgery.

Discussion

Distal radius fractures are very common during childhood.^{5,6} They mostly result from a fall onto an outstretched hand.⁶ And they are frequently associated with ulnar fractures (diaphysis, metaphysis, or ulnar styloid process).^{2,6} Isolated ulnar fracture is uncommon.²

One specific injury pattern combines distal radius fracture with DRUJ's disruption. It includes true ulnar dislocation (called Galeazzi injury) or ulnar epiphyseal avulsion associated with displacement (called Galeazzi-equivalent injury).^{3,5} Because of the relative weakness of bone in comparison with ligaments in skeletally immature children, Galeazzi-equivalent injury occurs exclusively in this population and predominates among young teenagers.^{3-5,7} However, this pattern seems infrequent although probably underestimated.^{7,8} The initial clinical assessment of the DRUJ could indeed be impossible in young children suffering from pain. One study showed that displaced Salter Harris 2 fractures of the radius and nonphyseal distal radius fractures were the most common injuries associated with DRUJ's instability in a population of 85 patients (mean age at trauma: 14 years, range: 6.7–17.8). Time between trauma and DRUJ's instability diagnosis ranged from 0 to 18 years (mean: 3 years).⁹

Galeazzi-equivalent injuries are classified according to the Letts and Rowhani classification^{3,10} (→Table 1).

Unlike in adults, most of the Galeazzi-equivalent injuries in children are treated by closed reduction and immobilization in an above elbow plaster cast for 4 to 6 weeks.^{3,7,11} The

forearm should be placed in full supination.^{10,11} Major residual instability, irreducible fracture, and dislocation are treated with open reduction.⁴ Reported results are good.^{7,12} However, there is a paucity of data on long-term results. One retrospective study including 10 children (mean age: 13.7 years, range: 11–16) with a mean follow-up of 6 years showed ulnar length discrepancy, bony deformation, or joint incongruence in the majority of the patients.¹² Misdiagnosed injuries or improper treatment could also compromise the DRUJ' stability, the wrist and forearm range of motion, and generate chronic pain.

To the best of our knowledge, our patient's fractures pattern has never been specifically described in the literature and is unusual for his age. It combines a displaced distal metaphyseal ulnar fracture and a distal radial buckle fracture. Each of these fractures is quite common, but their association is infrequent. Our initial differential diagnosis included the Galeazzi-equivalent injury versus "simple" metaphyseal fractures without DRUJ's instability. The DRUJ' stability could not be assessed in the emergency unit because of pain and swelling. Initial X-rays showed a certain displacement between both bones. However, no age-specific diagnostic radiographic measurements are described in literature for DRUJ's instability. An ulnar epiphyseal injury could also not be excluded because of the nonossified nature of the epiphysis in young children.

Our treatment included closed reduction under general anesthesia and immobilization in an above elbow plaster cast with the forearm in neutral alignment. Immobilization remained for a total of 8 weeks. Regarding the healing ability of young patients, we could question the length of our immobilization. Prognosis of radial buckle fracture is excellent because of the lack of significant displacement and retained residual stability.¹³ It allows short immobilization (2–4 weeks).¹⁴ Displaced metaphyseal fractures in children usually need longer immobilization (4–6 weeks).¹³ In up to 30% of complete radial metaphyseal fractures, loss of reduction occurs after closed reduction and immobilization.^{1,13,15} However, malunion, nonunion, and refracture are rarely observed.¹ No such data are available for complete ulnar metaphyseal fractures.

Table 1 Letts and Rowhani classification^{3,10}

Type		Description
A		Fracture of the radius at the junction of the middle and distal thirds +
	1	Dorsal dislocation of the ulna
	2	Epiphyseal fracture of the distal ulna with dorsal displacement of ulnar metaphysis
B		Fracture of the distal third of the radius +
	1	Dorsal dislocation of the ulna
	2	Epiphyseal fracture of the distal ulna with dorsal displacement of ulnar metaphysis
C		Greenstick fracture of the radius with dorsal bowing +
	1	Dorsal dislocation of distal ulna
	2	Epiphyseal fracture of distal ulna with displacement of ulnar metaphysis
D		Fracture of distal radius with volar bowing +
	1	Volar dislocation of the ulna
	2	Epiphyseal fracture of distal ulna with volar displacement of ulnar metaphysis

The injury in our patient healed without sequela. However, we recommend a low-suspicion threshold for DRUJ's instability in young children presenting with displaced distal metaphyseal ulnar fracture associated with distal radial buckle fracture. That suspicion should raise the necessity of appropriate treatment and follow-up. Delay between trauma and DRUJ's instability diagnosis can indeed be long.

Conflict of Interest

None.

References

- Schoenecker JG, Bae DS. Fracture of the distal radius and ulna. In: Flynn JM, Skaggs DL, Waters PM, eds. *Rockwood & Wilkins' Fractures in Children*. Eighth edition. Philadelphia: Wolters Kluwer Health; 2015:349–411
- Joeris A, Lutz N, Blumenthal A, Slongo T, Audigé L. The AO Pediatric Comprehensive Classification of Long Bone Fractures (PCCF): Part I: Location and morphology of 2,292 upper extremity fractures in children and adolescents. *Acta Orthop* 2017;88(02):123–128
- Rodríguez-Merchán EC. Pediatric fractures of the forearm. *Clin Orthop Relat Res* 2005;1(432):65–72
- Pannu GS, Herman M. Distal radius-ulna fractures in children. *Orthop Clin North Am* 2015;46(02):235–248
- Little JT, Klionsky NB, Chaturvedi A, Soral A, Chaturvedi A. Pediatric distal forearm and wrist injury: an imaging review. *Radiographics* 2014;34(02):472–490
- Otayek S, Ramanoudjame M, Fitoussi F. [Distal radius fractures in children]. *Hand Surg Rehabil* 2016;35S:S150–S155
- Eberl R, Singer G, Schalamon J, Petnehazy T, Hoellwarth ME. Galeazzi lesions in children and adolescents: treatment and outcome. *Clin Orthop Relat Res* 2008;466(07):1705–1709
- Walsh HP, McLaren CA, Owen R. Galeazzi fractures in children. *J Bone Joint Surg Br* 1987;69(05):730–733
- Andersson JK, Lindau T, Karlsson J, Fridén J. Distal radio-ulnar joint instability in children and adolescents after wrist trauma. *J Hand Surg Eur Vol* 2014;39(06):653–661
- Letts M, Rowhani N. Galeazzi-equivalent injuries of the wrist in children. *J Pediatr Orthop* 1993;13(05):561–566
- Imatani J, Hashizume H, Nishida K, Morito Y, Inoue H. The Galeazzi-equivalent lesion in children revisited. *J Hand Surg [Br]* 1996;21(04):455–457
- Cha SM, Shin HD, Jeon JH. Long-term results of Galeazzi-equivalent injuries in adolescents—open reduction and internal fixation of the ulna. *J Pediatr Orthop B* 2016;25(02):174–182
- Schweisch P. Distal forearm fractures in children: Initial management - UpToDate [Internet]. Accessed April 1, 2022 at: https://www.uptodate.com/contents/distal-forearm-fractures-in-children-initial-management?search=distal%20forearm%20fracture&source=search_result&selectedTitle=1~150&usage_type=default&display_rank=1
- Williams KG, Smith G, Luhmann SJ, Mao J, Gunn JD III, Luhmann JD. A randomized controlled trial of cast versus splint for distal radial buckle fracture: an evaluation of satisfaction, convenience, and preference. *Pediatr Emerg Care* 2013;29(05):555–559
- Hennrikus WL, Mehlman CT. The community orthopedic surgeon taking trauma call: pediatric distal radius and ulna fracture pearls and pitfalls. *J Orthop Trauma* 2019;33(Suppl 8):S6–S11