



Delayed Aspiration of Air Emboli from the Central Venous Catheter in a Case of Suspected Massive Venous Air Embolism: A Therapeutic Success

Ankur Khandelwal¹ Sudhansu S. Nayak² Anshil Bhatia¹ Amit K. Shrivastava³ Vikas Bhardwaj³

¹Department of Anaesthesiology and Critical Care, Sharda University School of Medical Sciences and Research, Greater Noida, Uttar Pradesh, India

²Department of Anesthesiology, Atal Bihari Vajpayee Institute of Medical Sciences and Ram Manohar Lohia Hospital, New Delhi, India

³Department of Neurosurgery, Neurosciences Centre, Sharda University School of Medical Sciences and Research, Greater Noida, Uttar Pradesh, India

Address for correspondence Ankur Khandelwal, MD, DM, Department of Anaesthesiology and Critical Care, Sharda University School of Medical Sciences and Research, Greater Noida, Uttar Pradesh 201310, India (e-mail: ankurchintus@gmail.com).

J Neuroanaesthesiol Crit Care 2022;9:198–200.

Abstract

Venous air embolism (VAE), though, clinically benign in majority of cases, the significant ones can lead to life-threatening cardiopulmonary and neurological consequences. Though studies mention the success rate of only 6 to 16% in aspirating air from the central venous catheter (CVC) during VAE, the technique is very specific for diagnosing VAE and has high therapeutic significance. We report a case in which delayed aspiration of air emboli from the CVC in suspected massive VAE during decompressive craniectomy resulted in rapid resolution of hemodynamic instability. If not inserted previously, CVC may be considered in a hemodynamically unstable patient with suspected VAE.

Keywords

- ▶ air embolism
- ▶ central venous catheters
- ▶ hemodynamic
- ▶ neurosurgery

Introduction

Venous air embolism (VAE) has been reported widely during neurosurgical procedures with an incidence between 16 and 86%.¹ The incidence is variable owing to the difference in surgical and anesthetic techniques, patient's position during surgery, and diagnostic methods used. The two most important factors that affect the clinical impact of VAE are the volume and the rate of air entrainment.² While studies mention the success rate of only 6 to 16% in aspirating air from the central venous catheter (CVC) during VAE, the technique is very specific for diagnosing VAE and has high

therapeutic significance.³ We report a case in which delayed aspiration of air emboli from the CVC in suspected massive VAE during decompressive craniectomy resulted in rapid resolution of hemodynamic instability. Written informed consent was taken from the patient for reporting this case for publication.

Case Report

A 40 year male weighing 70 kg encountered a road traffic accident and suffered head injury. He was taken to a nearby hospital where a noncontrast computed tomography of brain

article published online
July 20, 2022

DOI <https://doi.org/10.1055/s-0042-1748840>.
ISSN 2348-0548.

© 2022. Indian Society of Neuroanaesthesiology and Critical Care. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

was done that showed hemorrhagic contusions in right fronto-temporo-parietal (FTP) lobe with midline shift and mass effect. His initial Glasgow Coma Scale (GCS) on admission was E2V3M5 (E: eye opening, V: verbal response, M: motor response). Intubation of trachea was done and he was treated conservatively for 7 days in the same hospital. However, following deterioration of GCS on 8th day and expansion of hematoma, he was advised to undergo right FTP decompressive craniectomy and hematoma evacuation. The patient's family refused for any further intervention in the same hospital and shifted the patient to our hospital. His GCS on admission to our hospital was E1VTM4. We immediately scheduled the patient for right FTP decompressive craniectomy. In the operation theater, his endotracheal tube was replaced with a new sterile one of same size (8 mm ID). His preoperative vital parameters included blood pressure (BP)—129/83 mm Hg, heart rate (HR)—102 beats/min, and percentage of oxygen saturation (SpO₂)—98% (fraction of inspired concentration, FiO₂ = 0.4). We secured two wide bore 16 G intravenous (IV) cannula. Left radial artery was cannulated for beat-to-beat BP monitoring. CVC was not inserted. Induction of anesthesia was done with IV fentanyl—150 µg and IV propofol—120 mg. A loading dose of 8 mg cisatracurium was administered for muscle relaxation. Oxygen:air mixture (FiO₂: 50%) along with total IV anesthesia with fentanyl (75 µg/h), cisatracurium (2 mg/h), and propofol titrated maintain bispectral index of 40 to 50 that was incorporated to maintain intraoperative anesthesia. A small degree of reverse Trendelenburg positioning was done to facilitate decrease in intracranial pressure (ICP) and optimizing surgical visualization. The initial part of the surgery was uneventful. However, during the process of removal of the bone flap, there was injury to superior sagittal sinus (SSS) following which there was a sudden and sustained decline in end-tidal carbon dioxide (EtCO₂) from 32 to 22 mm Hg, BP from 126/75 to 64/38 mm Hg, and rapid progressive decline of SpO₂ from 98 to 72%. His HR increased from 92 to 130 beats/min associated with premature ventricular complexes (PVCs). There was associated blood loss of ~200 mL. An arterial blood gas analysis was done that showed partial pressure of carbon dioxide (PaCO₂) and EtCO₂ gradient of 15 mm Hg. Suspecting it to be VAE, the surgeon was informed immediately. The surgical field was flooded with normal saline and bony edges were sealed with wax. Simultaneously, we ventilated the lungs with 100% oxygen, lowered the head-end of the bed, infused IV fluids, and gave intermittent dose (3 mg) of mephentermine to avoid prolonged hypotension. After ~5 minutes of resuscitation, the EtCO₂ increased to 29 mm Hg, BP increased to 110/65 mm Hg, and HR decreased to 110/min. However, the SpO₂ increased only up to 95% at FiO₂ = 1 and occasional PVCs. The surgery was resumed. The FiO₂ was kept at 0.8 targeted to SpO₂ ≥ 94%. Another similar episode occurred during dural repair that was managed in the same way. However, after the second event, the BP remained in a lower range of 100/68 mm Hg and SpO₂ of 91 to 92% at FiO₂ = 0.8 and frequent PVCs. The surgery completed in ~150 minutes. During the surgery, the total blood loss amounted to around



Fig. 1 Aspiration of air emboli from central venous catheter (arrow).

600 mL. We transfused two units of packed red blood cells and two units of fresh frozen plasma. Considering the possibility of ongoing VAE (due to low BP and SpO₂), we inserted Electrocardiogram (ECG) guided CVC (7F, triple lumen) in the right subclavian vein immediately after the completion of the surgery. The tip of the catheter was targeted 2 cm below the superior vena cava (SVC) and the right atrium (RA) junction. To our surprise, 80 mL of air could be aspirated from the CVC (→ Fig. 1). The BP and SpO₂ started to increase within next 20 minutes and within an hour, the patient was hemodynamically stable with SpO₂ of 97% at FiO₂ = 0.5. The patient was electively ventilated after the surgery. On 4th postoperative day (POD 4), tracheostomy was done. Gradually, he started to improve and on POD 9, the patient was E3VTM6. The patient was gradually weaned off from the mechanical ventilation. Decannulation of the tracheostomy tube was done on POD 17 and the patient was discharged from the hospital on POD 20.

Discussion

In this case, the predominant inciting event for the occurrence of VAE was injury to the SSS. Open and noncollapsible veins form the major predisposing factor for entrainment of atmospheric air.⁴ Moreover, the slight head up position in our case probably aggravated the air entrainment. The above presentation could be attributed to manifestation of systemic hypotension due to blood loss as a result of SSS injury. However, there was sudden and sustained fall in EtCO₂.

Moreover, the oxygen desaturation (from 98% to 72%), ECG changes of PVCs, PaCO₂, and EtCO₂ gradient of 15 mm Hg, and the requirement of high FiO₂ (0.8) so as to maintain SpO₂ ≥ 94% even postresuscitation excluded the possibility of systemic hypotension as the primary cause of the event. Other probable causes could be trigeminocardiac reflex and raised ICP. However, in both cases, bradycardia is a predominant feature. Moreover, the timing of SSS injury and consequent decrease in EtCO₂ along with hypotension and oxygen desaturation show a definite association with VAE. Again, aspiration of air from the CVC is highly suggestive of the diagnosis of VAE.

This case report primarily brings forth the importance of CVC in the management of massive or significant VAE. Previous literatures have described air aspiration from the CVC leading to prompt resolution of hemodynamic perturbations in suspected VAE.^{5,6} In these reports, CVC was in situ during the time of VAE episode. Delayed aspiration of air via CVC is uncommon. In our case, the CVC was inserted at the end of the surgery due to persistence of hemodynamic instability. The CVC was inserted in our case as per the recommendation, that is, tip of the catheter was targeted 2 cm below the SVC and the RA junction that improves the success rate of aspiration of air emboli.³ In the delayed phase of VAE, cardiovascular support and frequent neurological monitoring are warranted till hemodynamic parameters and neurological integrity are restored.

ECG-guided central venous catheterization has been reported to be accurate and successful. The success rate has been reported to be around 96%.^{7,8} The technique is easy and inexpensive. Fluoroscopy/C-arm-guided CVC tip placement is also an alternative; however, it involves radiation. Use of transesophageal echocardiography (TEE) has been shown to have 100% accuracy, but the technique is invasive and it is not available in all centers.⁹

Early detection and management of VAE are the sine qua non to prevent potentially fatal complications. A battery of monitors is in use for its early detection before hemodynamic changes occur. TEE and Doppler ultrasound are highly sensitive. However, both have their own limitations and drawbacks. EtCO₂ tension monitoring, though intermediate in sensitivity, has the advantage of being noninvasive and

semiquantitative. Capnography detects clinically significant VAE with reasonable precision.¹⁰

Conclusions

The significant VAE episodes can lead to poorer outcome, albeit the fact that majority of the VAE episodes are clinically innocuous. Considering the diagnostic and therapeutic effectiveness of CVC, it may be inserted in a hemodynamically unstable patient with suspected VAE, if hitherto not introduced.

Conflict of Interest

None declared.

References

- Giraldo M, Lopera LM, Arango M. Venous air embolism in neurosurgery. *Rev Colomb Anesthesiol* 2015;43:40–44
- Mirski MA, Lele AV, Fitzsimmons L, Toung TJ. Diagnosis and treatment of vascular air embolism. *Anesthesiology* 2007;106(01):164–177
- Hanna PG, Gravenstein N, Pashayan AG. In vitro comparison of central venous catheters for aspiration of venous air embolism: effect of catheter type, catheter tip position, and cardiac inclination. *J Clin Anesth* 1991;3(04):290–294
- Bithal P, Dash HH, Vishnoi N, Chaturvedi A. Venous air embolism: does the site of embolism influence the hemodynamic changes? *Neurol India* 2003;51(03):370–372
- Khandelwal A, Gupta S, Prabhakar H, Burman S. Venous air embolism following application of Mayfield head clamp: a word of caution. *Indian J Anaesth* 2018;62(12):1002–1003
- Dube SK, Sharma VB, Kumar S, Rath GP. Venous air embolism during scalp dissection in a case of cerebellar haemangioblastoma. *J Neuroanaesth Crit Care* 2014;1:61–62
- Lee JH, Bahk JH, Ryu HG, Jung CW, Jeon Y. Comparison of the bedside central venous catheter placement techniques: landmark vs electrocardiogram guidance. *Br J Anaesth* 2009;102(05):662–666
- Gebhard RE, Szmuk P, Pivalizza EG, Melnikov V, Vogt C, Wartens RD. The accuracy of electrocardiogram-controlled central line placement. *Anesth Analg* 2007;104(01):65–70
- Venugopal AN, Koshy RC, Koshy SM. Role of chest X-ray in citing central venous catheter tip: a few case reports with a brief review of the literature. *J Anaesthesiol Clin Pharmacol* 2013;29(03):397–400
- Pandia MP, Bithal PK, Dash HH, Chaturvedi A. Comparative incidence of cardiovascular changes during venous air embolism as detected by transesophageal echocardiography alone or in combination with end tidal carbon dioxide tension monitoring. *J Clin Neurosci* 2011;18(09):1206–1209