




Emicizumab-induced photosensitivity

Pedro Asensi Cantó¹ Mercedes Rodríguez Serna² Pilar Lloret Madrid¹ Jürgen Solís Ruiz¹
Ana Rosa Cid Haro¹ Santiago Bonanad Boix¹ Saturnino Haya Guaita¹

¹Unidad de Hemostasia y Trombosis, Servicio de Hematología, Hospital La Fe, Valencia, Spain

²Servicio de Dermatología, Hospital La Fe, Valencia, Spain.

Address for correspondence Pedro A. Cantó, Graduated in Medicine, Department of Hematology, Hospital La Fe, Avinguda de Fernando Abril Martorell, 106, 46026 València, Spain (e-mail: asensi_ped@gva.es).

TH Open 2022;06:e154–e155.

Abstract

Keywords

- ▶ hemophilia therapy
- ▶ hemostasis
- ▶ hemophilia A/B

Emicizumab constitutes a novel and effective prophylaxis for hemophilia A patients with and without inhibitors. In this case report, we describe an emicizumab-induced photosensitivity that forced permanent sun-exposure suppression. To the best of our knowledge, this side effect had not been communicated until present.

A 35 year-old-man diagnosed with hemophilia A and inhibitors started prophylaxis with emicizumab. Six days after the first loading dose of 3 mg/kg (195 mg), he was exposed to the sun for 45 minutes without photoprotection, and 3 hours after he developed a cutaneous erythema in photoexposed areas (▶ **Fig. 1A and B**). Mucosa was not involved. With the exception of emicizumab, the patient had not used any new topic or systemic photosensitizing agent. A phototest (▶ **Fig. 1C**) evidenced erythema and edema starting from 12.5 mJ/cm² of ultraviolet B and an aberrant response to ultraviolet A starting from 7.5J/cm², coherent with photosensitivity. Autoimmunity blood tests, including antinuclear antibodies immunofluorescence assay, rheumatoid factor,

and erythrocyte sedimentation rate, they did not show any significant alteration.

Extended-spectrum photoprotection with a sun protection factor 50+ and avoidance of prolonged sunlight exposure permitted continuation of emicizumab and the regression of lesions in 2 weeks.

Emicizumab constitutes a novel and effective prophylaxis for hemophilia A patients with and without inhibitors. This report describes the first case of photosensitivity associated with emicizumab therapy. Photoprotection is only partially effective to prevent skin lesions produced by ultraviolet radiation. Therefore, sunlight avoidance, especially in maximal ultraviolet irradiation hours, is still necessary. Prolonged sun-exposure suppression may be a quality of life–deteriorating side effect.

received
December 3, 2021
accepted
April 11, 2022

DOI <https://doi.org/10.1055/s-0042-1749092>.
ISSN 2512-9465.

© 2022. The Author(s).
This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



Fig. 1 (A, B) Appearance of cutaneous erythema in photoexposed areas. (C) Phototest revealing erythema and edema.