



Biomechanical Assessment of the Prepatellar Quadriceps Continuation

Análisis biomecánico de la continuación prepatelar del cuádriceps

Carlos Infante^{1,2} Rony Silvestre³ Magaly Iñiguez¹ Roberto Negrin¹ Maximiliano Barahona²

¹Orthopedics Department, Clínica las Condes, Santiago, Chile

²Orthopedics Department, Hospital Clínico Universidad de Chile, Santiago, Chile

³Biomechanical Department, Clínica MEDS, Santiago, Chile

Address for correspondence Maximiliano Barahona, MD, Departamento de Ortopedia, Hospital Clínico Universidad de Chile, Santos Dumont 999/ 351, Independencia, Santiago, Chile (e-mail: maxbarahonavasquez@gmail.com).

Rev Chil Ortop Traumatol 2022;63(2):e83–e86.

Abstract

Objective To determine the ultimate tensile strength (UTS) of the prepatellar quadriceps continuation (PQC).

Materials and Methods A human cadaveric study was performed. Ten fresh-frozen specimens were used. The proximal and medial thirds of the anterior cortex of the patella were used to assess the mechanical properties of the PQC. In each specimen, transverse section areas measuring 0.2 cm² (A1) and 1 cm² (A2) were studied. A gradual load was applied to determine the UTS.

Results The median UTS of A1 was of 232.56 N (range: 141.23 N to 295.33 N), and that of A2 was of 335.30 N (range: 216.45 N to 371.40 N). The increment in UTS was significant between the 2 areas ($p = 0.006$).

Conclusions The clinical significance of the present study lies in the fact that it shows that the PQC is a strong tissue that can be a safe anchor for reconstruction around the patella. A relatively small area supports at least 140 N, and, as the area grows, the UTS increases as well.

Level of evidence Biomechanical study.

Keywords

- ▶ knee biomechanics
- ▶ prepatellar quadriceps continuation
- ▶ ultimate tensile strength

Resumen

Objetivo Determinar la resistencia máxima a la tracción (RMT) de la continuación prepatelar del cuádriceps (CPC).

Materiales y Métodos Se realizó un estudio en cadáveres humanos. Fueron incluidos diez especímenes, en los cuales se utilizaron los tercios proximal y medial de la cortical anterior de la patela para evaluar las propiedades mecánicas de la CPC. En cada espécimen, se estudió un área de sección transversal de 0,2 cm² (A1) y 1 cm² (A2). Se aplicó una carga gradual para determinar la RMT.

Resultados La mediana de la RMT en el A1 fue de 232,56 N (rango: 141,23 N a 295,33 N) y en el A2 fue de 335,30 N (rango: 216,45 N a 371,40 N). El incremento en la TMR fue significativo entre las 2 áreas ($p = 0,006$).

Palabras clave

- ▶ biomecánica de la rodilla
- ▶ continuación prepatelar del cuádriceps
- ▶ resistencia máxima a la tracción

received

March 28, 2021

accepted

May 10, 2022

published online

July 11, 2022

DOI <https://doi.org/10.1055/s-0042-1750364>.

ISSN 0716-4548.

© 2022. Sociedad Chilena de Ortopedia y Traumatología. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Conclusión El significado clínico de este estudio es que la CPC es un tejido fuerte que puede servir de anclaje seguro para reconstrucciones alrededor de la patela. Un área relativamente pequeña tolera al menos 140 N y, a medida que crece el área, también aumenta la RMT.

Nivel de evidencia Estudio biomecánico.

Introduction

The extensor mechanism has a complex anatomy, and its indemnity is crucial for the activities of daily life.¹ The prepatellar quadriceps continuation (PQC) corresponds to the fibers of the rectus femoris tendon that connect the quadriceps and patellar tendon.² The PQC is attached to the anterior cortex of the patella by a chondroapophyseal type of junction in which the tendon is gradually transformed into fibrocartilage.³ During the embryonic process of formation of the patella, the quadriceps tendon superficially embeds the patella, which will subsequently give rise to the PQC.⁴ In the developing patella, there is no periosteum or perichondrium; instead, a fibrocartilaginous attachment has been described. All sesamoid bones form this type of attachment in the surface that it is adjacent to a tendon.⁵

The PQC is an essential structure of the extensor mechanisms, and it must withstand friction force during flexion-extension of the knee, being in high risk of overuse injuries like prepatellar friction syndrome.⁶ Moreover, PQC indemnity is crucial for active knee extension: damage to this structure, although infrequent, causes severe functional deficit.^{2,7}

Macroscopic, histological and imaging descriptions of the PQC have been published.⁸ Nevertheless, no biomechanical assessment of this tissue has been conducted. Surgical techniques for patellar instability have been described using the PQC as a fixation site,^{9,10} so a proper understanding of its biomechanical properties is needed.

The purpose of the present study was to determine the ultimate tensile strength (UTS) of the PQC. Our hypothesis is that a small area of the PQC withstands more than 100 N, and, as the area increases, the resistance of this tissue does so as well.

Methods

Ten fresh-frozen human cadaveric knees without evidence of previous injury, abnormalities or surgery were used in the present study. The median age of the specimens was of 59 years (range: 36 to 70 years), and a total of 5 female and 5 male right knees were tested. The specimens were procured from Innoved Institute LLC (Rosemont, IL, United States) after approval from the institutional ethics in research board. Each specimen was thawed 24 hours before use.

An anterior longitudinal incision was made 10 cm above the upper edge of the patella to the tibial tubercle. The quadriceps tendon, prepatellar continuation, and patellar tendon were carefully exposed. The proximal and medial

thirds of the patella were used to assess the mechanical properties of the PQC. In all ten specimens, a simple and a double stich were performed, spanning an area of 0.2 cm² (A1) and 1 cm² (A2) respectively. In five specimens, a simple stich was placed on the proximal third, and a double stich, on the middle third of the patella. Meanwhile, in the other five specimens, the double stich was placed proximally, and the simple stich, on the middle third. A non-absorbable ultra-high molecular weight polyethylene and polydioxanone suture was used (Orthocord #2, Ethicon, DePuy Synthes, Raynham, MA, United States).

The simple stich started 5 mm laterally to the antero-medial edge of the patella. It was performed as deep as palpating the anterior cortex and advancing 1 cm toward the lateral side. A quintuple knot was made to fix the point. For the double stich, after passing the first point 6 mm distally, a parallel 1-cm suture pass in the opposite direction was made, which was also knotted with a quintuple knot. Hence, the simple stich suture group test was for a 0.2-cm² transverse section area of tissue, and the double stich suture group test, for a 1-cm² transverse section area of PQC tissue. (► Fig. 1)

A stress-strain device was used (► Fig. 2), which, in one side, had a load cell with the capacity to measure up to 1,000 N (Model SSM-AJ-500; Interface, Scottsdale, AZ, United States). A sensor – a linear potentiometer for length measurement (range 300mm, reading 0.01 mm) – to measure




Fig. 1 Scheme of the suture distribution used in five specimens. In the other five, the double stich was placed in the proximal third of the patella, and the simple stich, in the middle third.




Fig. 2 A stress-strain device was used to measure the UTS (A). A cadaveric patella mounted from the suture in the device (B). A gradual load was applied to measure the PQC deformation from which the stress and strain were determined.

linear displacement was also used (Digitalanzeige Digital Linear Scale Linear Skala Externe Fernanzeige, Sauter, Basel, Switzerland). They were later exported to the Matlab (The MathWorks, Inc., Natick, MA, United States) software to build the force-displacement curve. A gradual load was applied to measure the PQC deformation from which the stress and strain were determined. The UTS was recorded. The load was applied proximally, along the longitudinal axis of the patella, emulating the tensile force of the quadriceps tendon. Failure was defined as rupture or detachment of the PQC.

The median and range of the UTS were reported. A non-parametric unpaired median test was used to compare the groups. A significance level of 5% was used, and data were processed using the Stata (StataCorp LLC, College Station, TX, United States) software, version 15.

Results

All ten specimens were tested. The median UTS of A1 was of 232.56 N (range: 141.23 N to 295.33 N), and the median UTS of A2 was of 335.30 N (range: 216.45 N to 371.40 N). The increase in UTS was significant between the two areas ($p = 0.006$). The distribution of the UTS is shown in **Fig. 3**.

Discussion

The main finding of the present study is that the PQC is a resistant tissue. A small area of 0.2 mm^2 offers a resistance of 140 N, and, as the section area grows, so does the resistance of the tissue.

The mean UTS of the medial patellofemoral ligament (MPFL) complex is of 140 N,¹¹ so it is equivalent to the




Fig. 3 Distribution of the UTS.

minimum UTS found for group A1 of the present study. In MPFL reconstruction, soft-tissue fixation in the femoral side has achieved good clinical results, and no significant increase in rate of rerupture.¹² Moreover, the mean ultimate load of an MPFL reconstruction using a double tunnel technique in the patella was of 213 N,¹³ which is lower than a 1-cm² attachment in the PQC found in the present study (335 N). Given the aforementioned information, it does not seem disproportionate to use the PQC as an anchor for the reconstruction of the MPFL, which would help to reduce the costs and complications related to tunnels in the patella. However, studies designed for this purpose should be performed first.

The Green¹⁴ and the Madigan et al.¹⁰ quadricepsplasties are surgical procedures described for patellar instability, in which a distal and lateral transfer of the insertion the vastus medialis is performed.^{10,14} This transfer depends on the

fixation in the PQC. Despite the high rate of dislocation after surgery, failure has been attributed to vector forces and not PQC attachment-related failures.¹⁵ Further, after the procedure, the patients tend to experience an increase in patellar height due to the vastus medialis pull.¹⁶ Hence, this adds to the fact that the PQC is a strong tissue to anchor an MPFL reconstruction, which would have a more anatomic vector to stop the patella from lateral instability.

The PQC has also been used to transfer the vastus medialis in patellar instability after total knee arthroplasty. Kelly et al.¹⁷ described a modification to the originally Insall technique, in which the vastus medialis is transferred laterally, anchoring it both in the quadriceps tendon and in the CPC.¹⁷ It is a rescue procedure, and its main indication is when a patient with chronic patellar dislocation undergoes total knee replacement.⁹

The limitation of the present study is that only the UTS was measured, and tissue resistance to cyclic forces was not determined. Nevertheless, the PQC has been used as an attachment for decades, and no complications related to it have been reported. The present study should encourage the assessment of the resistance of an MPFL reconstruction fixed to the PQC.

Conclusion

The PQC is a strong tissue that can be a safe attachment for reconstruction around the patella. A relatively small area withstands at least 140 N, and, as the area grows, the UTS increases as well.

Availability of Data

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Author's contributions

CI: contributed with the original idea, conceived the study, drafted the paper, and performed a critical review of the final paper. RS: performed the biomechanical testing. MI: coordinated and supervised the management of the cadaver specimens. RN: performed a critical review of the final paper. MB: performed the data analysis, literature review and drafted the manuscript. All authors have read and approved the final submitted manuscript.

Funding

The authors declare that they have received no funding pertaining to the performance of the present study.

Conflict of Interests

The authors have no conflict of interests to declare.

Ethical Approval

The present study was approved by the Ethics in Research Board at Clínica Las Condes, Santiago, Chile.

References

- Waligora AC, Johanson NA, Hirsch BE. Clinical anatomy of the quadriceps femoris and extensor apparatus of the knee. *Clin Orthop Relat Res* 2009;467(12):3297–3306
- Majeed H, dos Remedios I, Datta P, Griffiths D. Prepatellar continuation rupture: Report of an unusual case. *Knee* 2014;21(05):979–981
- Evans EJ, Benjamin M, Pemberton DJ. Fibrocartilage in the attachment zones of the quadriceps tendon and patellar ligament of man. *J Anat* 1990;171:155–162
- Eyal S, Blitz E, Shwartz Y, Akiyama H, Schweitzer R, Zelzer E. On the development of the patella. *Development* 2015;142(10):1831–1839
- Dwek JR. The periosteum: what is it, where is it, and what mimics it in its absence? *Skeletal Radiol* 2010;39(04):319–323
- Claes T, Claes S, De Roeck J, Claes T. Prepatellar Friction Syndrome: a common cause of knee pain in the elite cyclist. *Acta Orthop Belg* 2015;81(04):614–619
- Teixeira PA, Lecocq S, Moisei A, Chanson A, Louis M, Blum A. Sleeve-like avulsion fracture of the superior pole of the patella in a healthy adult. *Diagn Interv Imaging* 2013;94(01):108–111
- Wangwinyuvirat M, Dirim B, Pastore D, et al. Prepatellar quadriceps continuation: MRI of cadavers with gross anatomic and histologic correlation. *AJR Am J Roentgenol* 2009;192(03):W111–6
- Dao Q, Chen DB, Scott RD. Proximal patellar quadricepsplasty realignment during total knee arthroplasty for irreducible congenital dislocation of the patella: a report of two cases. *J Bone Joint Surg Am* 2010;92(14):2457–2461
- Madigan R, Wissinger HA, Donaldson WF. Preliminary experience with a method of quadricepsplasty in recurrent subluxation of the patella. *J Bone Joint Surg Am* 1975;57(05):600–607
- Criscenti G, De Maria C, Sebastiani E, et al. Material and structural tensile properties of the human medial patello-femoral ligament. *J Mech Behav Biomed Mater* 2016;54:141–148
- Marot V, Sanchis-Alfonso V, Perelli S, et al. Isolated reconstruction of medial patellofemoral ligament with an elastic femoral fixation leads to excellent clinical results. *Knee Surg Sports Traumatol Arthrosc* 2021;29(03):800–805
- Placella G, Speziali A, Sebastiani E, Morello S, Tei MM, Cerulli G. Biomechanical evaluation of medial patello-femoral ligament reconstruction: comparison between a double-bundle converging tunnels technique versus a single-bundle technique. *Musculoskelet Surg* 2016;100(02):103–107
- Green W. Quadricepsplasty in Treatment of Recurrent Subluxation of the Patella. *Zit nach Madigan*.
- Vivod G, Verdonk P, Drobnič M. Long-term clinical and radiographic outcome of patello-femoral realignment procedures: a minimum of 15-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2014;22(11):2747–2755
- Vacariu A, Studer K, Rutz E, Camathias C. High failure rate 10.8 years after vastus medialis transfer and lateral release (Green's quadricepsplasty) for recurrent dislocation of the patella. *Arch Orthop Trauma Surg* 2020;140(10):1349–13579
- Kelly MA. Proximal realignment and medial tibial tubercle transfer. *Oper Tech Sports Med* 1999;7(02):76–80