



## Editorial

# Dual-Energy Computed Tomography Applications in the Abdomen

Rochita Venkataraman<sup>1</sup> Venkatesh Kasi Arunachalam<sup>2</sup><sup>1</sup>Department of Radiology, Advantage Imaging and Research, Chennai, Tamil Nadu, India<sup>2</sup>Department of Radiology, Kovai Medical Center and Hospital, Coimbatore, India

J Gastrointestinal Abdominal Radiol ISGAR 2022;5:75.

Among the recent developments in computed tomography (CT), dual-energy CT (DECT) is one of the most important recent advances. The challenges with single-energy CT (SECT) acquisition are tissue characterization and lesion differentiation. In SECT, the two different elements with the same attenuation depict a similar Hounsfield's unit. DECT can overcome this challenge. In DECT, tissues are imaged with two energy levels. The attenuation response to both energy levels is used to characterize the tissues further. This issue especially deals with the applications of DECT in abdominal imaging.

In the first article, Alavandar et al<sup>1</sup> have discussed the basic principles and available hardware in DECT. In the second article, Narappulan et al<sup>2</sup> have analyzed the role of virtual monoenergetic imaging, one of the essential image sets generated from dual-energy source images. They discuss its role in evaluating hypervascular focal lesions in the liver. In the following article, Marri and Madhusudhan<sup>3</sup> have explained the role of DECT in the evaluation of diffuse liver diseases like fat/iron deposition and fibrosis. In the subsequent article, Singh et al<sup>4</sup> have discussed the use of DECT in evaluating gall bladder pathologies. Further in the issue, Mroueh et al<sup>5</sup> have explained the role of DECT in pancreas imaging with applications in pathologies like pancreatitis, trauma, and pancreatic neoplasms. In the following article by Mehra,<sup>6</sup> DECT role in urolithiasis has been discussed extensively.

Lastly, in the article by Tripathy et al,<sup>7</sup> DECT applications in abdominal interventions are discussed. The role of calcium and bone subtraction images in evaluating vessels in atherosclerotic diseases and virtual noncontrast images/iodine

maps in evaluating residual tumors following locoregional treatment of HCC is explained very well in this article. This article also analyzes the evaluation of endoleaks in low monoenergetic images and metal artifacts reduction in high monoenergetic data sets.

We wish our readers an enjoyable and highly informative reading.

**Conflict of Interest**

None declared.

**References**

- 1 Alavandar E, Arunachalam VK, Narappulan N, et al. Principles and available hardware in DECT. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:76–84
- 2 Narappulan N, Arunachalam VK, Alavandar E, Selvaraja S, Renganathan R, Cherian M. Evaluation of hypervascular focal liver lesions utilizing virtual monoenergetic images from third-generation dual-source dual-energy computed tomography. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:85–93
- 3 Marri UK, Madhusudhan KS. Dual-energy computed tomography in diffuse liver diseases. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:94–106
- 4 Singh T, Gupta P. Role of dual-energy computed tomography in gallbladder disease: a review. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:107–113
- 5 Mroueh N, Cao J, Kambadakone A. Dual-energy CT in the pancreas. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:114–120
- 6 Mehra S. Role of dual-energy computed tomography in urolithiasis. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:121–126
- 7 Tripathy TP, Patel R, Chandel K, Mukund A. Utility of dual-energy CT in abdominal interventions. *J Gastrointestinal Abdominal Radiol ISGAR* 2022;5:127–134

**Address for correspondence**  
Venkatesh Kasi Arunachalam,  
Arunachalam, DMRD, DNB, FRCR,  
Department of Radiology, Kovai  
Medical Center and Hospital,  
Coimbatore, Tamil Nadu 641014,  
India (e-mail: radvenki79@gmail.  
com).

DOI <https://doi.org/10.1055/s-0042-1750724>.  
ISSN 2581-9933.

© 2022. Indian Society of Gastrointestinal and Abdominal Radiology. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India