

K. WIESNER* (UNIVERSITY OF NEW BRUNSWICK, FREDERICTON, CANADA)

The Total Synthesis of Racemic Talatisamine


Pure Appl. Chem. **1975**, *41*, 93–112, DOI: 10.1351/pac197541010093.

K. WIESNER*, T. Y. R. TSAI, K. HUBER, S. E. BOLTON, R. VLAHOV (UNIVERSITY OF NEW BRUNSWICK, FREDERICTON, CANADA AND BULGARIAN ACADEMY OF SCIENCES, SOFIA, BULGARIA)

Total Synthesis of Talatisamine, a Delphinine Type Alkaloid

J. Am. Chem. Soc. **1974**, *96*, 4990–4992, DOI: 10.1021/ja00822a048.

Total Synthesis of (±)-Talatisamine



Significance: (–)-Talatisamine is a potassium ion channel inhibitor isolated from an *aconitum* species. Wiesner and co-workers reported the first total synthesis of (±)-talatisamine. Its preparation was facilitated by synthetic studies on degradation products of natural (–)-talatisamine. Other diterpene alkaloids prepared by the Wiesner group include atisine and napelline.

Comment: The tricyclic core of (±)-talatisamine is assembled by a Diels-Alder reaction. Ring expansion from the cyclobutane in **E** via retro-aldol-aldol reaction furnishes the bicyclo[2.2.2]octane motif within **F**. The atisine-type scaffold **G** is converted into the delphinine-type skeleton **I** by Wagner-Meerwein rearrangement in analogy to Wiesner's biosynthetic proposal. An oxidation followed by aza-Prins reaction furnishes the hexacyclic 6/6/5/6/5/6-membered-ring scaffold of (±)-talatisamine.

SYNFACTS Contributors: Erick M. Carreira, Lukas J. Sprenger
Synfacts 2023, 19(06), 0539 Published online: 11.05.2023
DOI: 10.1055/s-0042-1752540; Reg-No.: C04223SF

Category

Synthesis of Natural Products

Key words

(±)-talatisamine

Diels-Alder reaction

[2+2] photocycloaddition

Lemieux-Johnson oxidation

retro-aldol-aldol reaction

Wagner-Meerwein rearrangement

aza-Prins reaction

Synfact Classic