



Bruxism as a Consequence of Stress and Movement Disorders: Brief Review

Artak Heboyan¹ Mohmed Isaqali Karobari^{2,3} Abdullah Hamed Alwadaani⁴ Anand Marya⁵
Muhammad Sohail Zafar^{6,7}

¹ Faculty of Stomatology, Department of Prosthodontics, Yerevan State Medical University after Mkhitar Heratsi, Str. Koryun 2, Yerevan, Armenia

² Faculty of Dentistry, Department of Restorative Dentistry & Endodontics, University of Puthisastra, Phnom Penh, Cambodia

³ Conservative Dentistry Unit, School of Dental Sciences, Universiti Sains Malaysia, Health Campus, Kota Bharu, Kelantan, Malaysia

⁴ Dental Complex, College of Dentistry, King Faisal University, Al-Ahsa, Kingdom of Saudi Arabia

⁵ Department of Orthodontics, Faculty of Dentistry, University of Puthisastra, Phnom Penh, Cambodia

⁶ Department of Restorative Dentistry, College of Dentistry, Taibah University, Al Madinah, Al Munawwarah, Saudi Arabia

⁷ Department of Dental Materials, Islamic International Dental College, Riphah International University, Islamabad, Pakistan

Address for correspondence Artak Heboyan, BDS, DMD, MSD, PhD, Department of Prosthodontics, Faculty of Stomatology, Yerevan State Medical University after Mkhitar Heratsi, Str. Koryun 2, Yerevan 0025, Armenia (e-mail: heboyan.artak@gmail.com).

Eur J Gen Dent 2022;11:81–83.

Abstract

Keywords

- ▶ teeth grinding
- ▶ parasomnia
- ▶ parafunction
- ▶ repetitive jaw movement
- ▶ mastication muscle activity

Bruxism is considered to be an unusual repetitive movement syndrome, which is described as involuntary gnashing of teeth. The condition is classified as sleep bruxism (SB) and awake bruxism (AB). The objective of the study was to review the current literature on bruxism and its potential relation to stress and movement disorders. Factors causing bruxism are numerous. It has been found that people suffering from stress are more prone to the condition than healthy subjects. Awake bruxism is frequently observed in subjects with hyperkinetic movement disorders and it can be associated with anxiety, which involves the limbic part of the basal ganglia. However, further studies should be conducted to interpret the link more thoroughly.

Review

Stress is identified by people as a “condition of being in long-term tension due to requirements that exceed their potential,” as presented by some researchers.¹ Extreme long-lasting stress may have perpetual effect on the reaction to stress, surpassing the natural ability of the human body to adjust.² Stress-related physiological and psychological disorders are defined by the type of stress stimulus³ and its ability to generate various functional changes, which include consequences observed in the maxillofacial region.⁴

Bruxism can be defined as a condition in which an abnormal recurrent jaw action of clenching and mandible thrusting are observed.^{5,6} Currently, the condition is classified as sleep bruxism (SB) and awake bruxism (AB). However, phenomenologically, there are significant differences observed between the two types. Semivoluntary AB, i.e., non-functional activity of gnashing that occurs during wakefulness is defined as caused by stress and anxiety-conditioned recurrent or continued teeth grinding or jaw thrusting.^{5,7} SB, which is a sleep-related stereotypic movement disorder

DOI <https://doi.org/10.1055/s-0042-1754374>.
ISSN 2320-4753.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

associated with both rhythmical and non-rhythmical mastication muscle activity, which is actually considered to be a form of parasomnia.^{5,7,8} SB has previously been considered as a dysfunctional movement or pathological condition, whereas it is now accepted as a centrally controlled condition with various systemic risk factors. It has been postulated that sleep bruxism may have a protective role during sleep, for example, in relation to airway maintenance or in stimulating saliva flow.⁹ Clinical manifestation of both awake and sleep bruxism depends on diverse individual behavioral patterns.⁵

Regular rotation pattern of jaw-opener and jaw-closer muscle activity is deteriorated in bruxism, resulting in activation of both muscles.¹⁰ Bruxism acts as a permanent motion mechanism. The severity of the symptoms is directly related to the malfunctioning of the body. Thus, negative emotions, depression, and stressful conditions can bring an increase in muscle tone performance and teeth gnashing.¹¹ Stress and mental conditions are commonly recognized to cause the progression in parafunctional habits, as well as TMDs. Bruxism is known as a parafunctional habit, which has psychosocial, emotional, and psychological bases.¹² It is crucial for both treatment and prevention to diagnose the bruxism in its early stages.¹³

Bruxism is multifactorial in its etiological nature.^{14,15} The possible factors which contribute to the development of the condition include smoking,¹⁶ high alcohol consumption¹⁷ and coffee intake,¹⁸ gastroesophageal reflux disease,¹⁹ sleep apnea,²⁰ depression,²¹ and anxiety.²²⁻²⁵ These also include genetic predisposition and behavioral patterns.¹³ The condition can be also associated with anatomical and morphological abnormalities, such as dental malocclusion.²⁶

There are a few studies that have revealed the possible relation of bruxism to movement disorders and emotional alterations,²⁷ most of them being clinical reports. ► **Table 1** shows the number of bruxism cases in patients with various movement impairments.

AB is not commonly reported in patients with Parkinson's syndrome. This might be due to the fact that AB is not a specific symptom of the syndrome and is considered rather minor issue compared with other major motor as well as cognitive disorders. Obviously, the medications directed to dopaminergic pathways stimulate AB.²⁸ Moreover, continuous stimulation with apomorphine induce non-functional masticatory activity. On the contrary, although people with Parkinson's syndrome suffer from sleep disorders, SB is

Table 1 Cases of bruxism reported in patients with movement impairment²⁹

Movement impairments	Reported cases of bruxism
Parkinsonian syndromes	<i>n</i> = 175
Chorea	<i>n</i> = 7
Primary dystonia	<i>n</i> = 300
Psychomotor disorders	<i>n</i> = 57
Restless legs syndrome	<i>n</i> = 380
Secondary bruxism	<i>n</i> = 355

infrequent. It is mostly observed during NREM sleep and in the stage before alteration to REM sleep.²⁹

Bruxism is commonly reported among subjects with hyperkinetic syndromes including dystonia, levodopa-induced dyskinesia, and chorea. Pathologies in which AB is mostly observed and is found to be rather severe are stereotypic and cognitive impairments as Rett's and Down syndromes as well as autistic spectrum disorders.³⁰ Significant characteristic feature of AB is the relation of bruxism to anxiety. Personality traits have their impact on the symptoms observed in the general population,³¹ suggesting limbic system mechanisms, particularly basal ganglia being involved in the pathophysiology.

Amygdala, the hypothalamus, as well as other subcortical regions of the brain, such as red nucleus, the anterior pretectal nucleus, the cerebellum, the periaqueductal gray, the raphe nuclei and various parts of basal ganglia take their part in mastication. Inputs that are transmitted from the basal ganglia are received by the areas of the brain located in the frontal cerebral cortex via thalamocortical projection.^{31,32}

Regardless of the etiological factors, bruxism can be modulated by stress and various types of movement impairments.

For the management of the bruxism occlusal therapy can be performed, which will provide an opportunity to achieve harmonious relationship between occluding surfaces. Occlusal appliances such as hard splints are preferred over soft splints because they prevent inadvertent tooth movements.³³ Moreover, hard splints are more effective in reducing bruxism activity than the soft ones. Biofeedback can be applied for bruxism during wakefulness as well as for sleep bruxism. While awake, patients can be trained to control their jaw muscle activities through auditory. For SB, auditory, electrical, vibratory, and even taste stimuli can be used for feedback. Drugs that have paralytic effect on the muscles through an inhibition of acetylcholine release at the neuromuscular junction (botulinum toxin) decreases bruxism activity, especially in severe cases with comorbidities.¹⁵

Conclusion

Considering all above-mentioned data, it can be concluded that bruxism is related to stress. However, further studies should be performed to establish the definite association. Bruxism is more common in patients with movement disorders and the condition is subject to treatment by neurologist. Moreover, the management of patients with the condition requires clinical evaluation by various specialists. Bruxism management strategies include occlusal therapy with hard splints, biofeedback, and pharmacological therapy.

Funding
None.

Conflict of Interest
None declared.

References

- 1 American Institute of Stress. What is Stress? (2016). Available online at: Accessed July 12, 2020 <http://www.stress.org/what-is-stress>
- 2 Chrousos GP. Stress and disorders of the stress system. *Nat Rev Endocrinol* 2009;5(07):374–381
- 3 Crestani CC. Emotional stress and cardiovascular complications in animal models: a review of the influence of stress type. *Front Physiol* 2016;7:251
- 4 Saczuk K, Lapinska B, Wilmont P, Pawlak L, Lukomska-Szymanska M. Relationship between sleep bruxism, perceived stress, and coping strategies. *Int J Environ Res Public Health* 2019;16(17):3193
- 5 Lobbezoo F, Ahlberg J, Raphael KG, et al. International consensus on the assessment of bruxism: report of a work in progress. *J Oral Rehabil* 2018;45(11):837–844
- 6 Yap AU, Chua AP. Sleep bruxism: current knowledge and contemporary management. *J Conserv Dent* 2016;19(05):383–389
- 7 Reissmann DR, John MT, Aigner A, Schön G, Sierwald I, Schiffman EL. Interaction between awake and sleep bruxism is associated with increased presence of painful temporomandibular disorder. *J Oral Facial Pain Headache* 2017;31(04):299–305
- 8 Manfredini D, Winocur E, Guarda-Nardini L, Paesani D, Lobbezoo F. Epidemiology of bruxism in adults: a systematic review of the literature. *J Orofac Pain* 2013;27(02):99–110
- 9 Beddis H, Pemberton M, Davies S. Sleep bruxism: an overview for clinicians. *Br Dent J* 2018;225(06):497–501
- 10 Lavigne GJ, Khoury S, Abe S, Yamaguchi T, Raphael K. Bruxism physiology and pathology: an overview for clinicians. *J Oral Rehabil* 2008;35(07):476–494
- 11 American Academy of Sleep Medicine (AASM) International Classification of Sleep Disorders: Diagnostic and Coding Manual 2005:51–55
- 12 Wieckiewicz M, Paradowska-Stolarz A, Wieckiewicz W. Psychosocial aspects of bruxism: the most paramount factor influencing teeth grinding. *BioMed Res Int* 2014;2014:469187
- 13 Yamaguchi T, Abe S, Rompré PH, Manzini C, Lavigne GJ. Comparison of ambulatory and polysomnographic recording of jaw muscle activity during sleep in normal subjects. *J Oral Rehabil* 2012;39(01):2–10
- 14 Balanta-Melo J, Toro-Ibacache V, Kupczik K, Buvinic S. Mandibular bone loss after masticatory muscles intervention with botulinum toxin: an approach from basic research to clinical findings. *Toxins (Basel)* 2019;11(02):84
- 15 Shetty S, Pitti V, Satish Babu CL, Surendra Kumar GP, Deepthi BC. Bruxism: a literature review. *J Indian Prosthodont Soc* 2010;10(03):141–148
- 16 Rintakoski K, Ahlberg J, Hublin C, et al. Tobacco use and reported bruxism in young adults: a nationwide Finnish Twin Cohort Study. *Nicotine Tob Res* 2010;12(06):679–683
- 17 Rintakoski K, Kaprio J. Legal psychoactive substances as risk factors for sleep-related bruxism: a nationwide Finnish Twin Cohort study. *Alcohol* 2013;48(04):487–494
- 18 Kuhn M, Türp JC. Risk factors for bruxism. *Swiss Dent J* 2018;128(02):118–124
- 19 Machado NA, Fonseca RB, Branco CA, Barbosa GA, Fernandes Neto AJ, Soares CJ. Dental wear caused by association between bruxism and gastroesophageal reflux disease: a rehabilitation report. *J Appl Oral Sci* 2007;15(04):327–333. Doi: 10.1590/S1678-7752007000400016
- 20 Martynowicz H, Gac P, Brzecka A, et al. The relationship between sleep bruxism and obstructive sleep apnea based on polysomnographic findings. *J Clin Med* 2019;8(10):1653
- 21 Smardz J, Martynowicz H, Wojakowska A, Michalek-Zrabcowska M, Mazur G, Wieckiewicz M. Correlation between sleep bruxism, stress, and depression—a polysomnographic study. *J Clin Med* 2019;8(09):1344
- 22 Cruz-Fierro N, Martínez-Fierro M, Cerda-Flores RM, et al. The phenotype, psychotype and genotype of bruxism. *Biomed Rep* 2018;8(03):264–268
- 23 Castrolforio T, Bargellini A, Rossini G, Cugliari G, Rainoldi A, Deregis A. Risk factors related to sleep bruxism in children: a systematic literature review. *Arch Oral Biol* 2015;60(11):1618–1624
- 24 Soares-Silva L, Tavares-Silva C, Fonseca-Gonçalves A, Maia LC. Presence of oral habits and their association with the trait of anxiety in pediatric patients with possible sleep bruxism. *J Indian Soc Pedod Prev Dent* 2019;37(03):245–250
- 25 Ahlberg J, Lobbezoo F, Ahlberg K, et al. Self-reported bruxism mirrors anxiety and stress in adults. *Med Oral Patol Oral Cir Bucal* 2013;18(01):e7–e11
- 26 Lobbezoo F, Ahlberg J, Glaros AG, et al. Bruxism defined and graded: an international consensus. *J Oral Rehabil* 2013;40(01):2–4
- 27 Yap AU, Chua AP. Sleep bruxism: Current knowledge and contemporary management. *J Conserv Dent* 2016;19(05):383–389
- 28 Winocur E, Gavish A, Voikovitch M, Emodi-Perlman A, Eli I. Drugs and bruxism: a critical review. *J Orofac Pain* 2003;17(02):99–111
- 29 Ella B, Ghorayeb I, Burbaud P, Guehl D. Bruxism in movement disorders: a comprehensive review. *J Prosthodont* 2017;26(07):599–605
- 30 El Khatib AA, El Tekeya MM, El Tantawi MA, Omar T. Oral health status and behaviours of children with autism spectrum disorder: a case-control study. *Int J Paediatr Dent* 2014;24(04):314–323
- 31 Gungormus Z, Erciyas K. Evaluation of the relationship between anxiety and depression and bruxism. *J Int Med Res* 2009;37(02):547–550
- 32 Morquette P, Lavoie R, Fhima MD, Lamoureux X, Verdier D, Kolta A. Generation of the masticatory central pattern and its modulation by sensory feedback. *Prog Neurobiol* 2012;96(03):340–355
- 33 Alkan A, Bulut E, Arici S, Sato S. Evaluation of treatments in patients with nocturnal bruxism on bite force and occlusal contact area: a preliminary report. *Eur J Dent* 2008;2(04):276–282