

Urinary Nephrin is a Sensitive Marker to Predict Early Onset of Nephropathy in Type 2 Diabetes Mellitus

Ganesh Veluri¹ Murugan Mannangatti¹

¹Department of Biochemistry, Arupadai Veedu Medical College and Hospital, Vinayaka Missions Research Foundation (deemed to be university, Salem, Tamil Nadu), Kirumampakkam, Puducherry, India

Address for correspondence Murugan Mannangatti, PhD, Department of Biochemistry, Arupadai Veedu Medical College and Hospital, Vinayaka Mission University Research Centre, Pondicherry, 607402, India (e-mail: avmcbiochemistryedu@gmail.com).

J Lab Physicians 2022;14:497-504.

Abstract	Objective The present study has been designed to evaluate urinary nephrin that best predicts the occurrence of renal dysfunction in type 2 diabetes mellitus patients and its correlation with clinical parameters of nephropathy. Materials and Methods A total of 80 type 2 diabetes mellitus and 40 age- and gendermatched healthy controls were recruited. Biochemical and clinical parameters were analyzed in all the study participants. Analysis of variance was performed to compare				
	the differences between the groups. Pearson's correlation analysis was used to analyz the association of nephrin with clinical parameters of nephropathy. Receiver operatir characteristic curves were constructed to study the diagnostic accuracy of markers to identify disk stip and have athe				
	Results The levels of nephrin were significantly elevated in both groups of type 2 diabetes mellitus patients when compared with healthy individuals ($p = 0.0001$). The urinary nephrin was positively correlated with hemoglobin A1c and urinary albumin creatinine ratio ($r = 0.520$, 0.657, $p < 0.0001$) and negatively correlated with estimated glomerular filtration rate ($r = -0.539$, $p < 0.0001$). The diagnostic sensitivity and specificity of nephrin for nephropathy were 100 and 88%, respectively, and urinary				
 Keywords HbA1c type 2 diabetes mellitus urinary nephrin 	albumin creatinine ratio was 43 and 76%, respectively. Conclusion The study findings suggest that nephrin levels are strongly and positively associated with nephropathy in type 2 diabetes mellitus patients and it has a greater potential to be an early predictable marker of nephropathy than urinary albumin creatinine ratio.				

Introduction

Type 2 diabetes mellitus (T2DM) caused by hyperglycemia due to defect in insulin secretion from the β cells of pancreas and inactivation of insulin leads to insulin resistance.¹ T2DM

published online October 20, 2022 DOI https://doi.org/ 10.1055/s-0042-1757718. ISSN 0974-2727. is one of the leading disorders in the world, where 415 million people were affected by the year 2015 and it is estimated to reach 650 million by 2030; in the Indian scenario 65 million were affected by the year 2013 and it is estimated to reach 103 million by the year 2030.^{2,3}

 $\ensuremath{\mathbb{C}}$ 2022. The Indian Association of Laboratory Physicians. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-nc-nd/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

T2DM leads to many micro- and macrovascular complications, this in turn affect many organs like the kidney, retina of the eye, brain, lungs, and heart, etc. and most of the diabetic people are affected with nephropathy. Diabetic nephropathy (DN) is the major microvascular complication in patients with T2DM due to hyperglycemia in the blood and hypoglycemia in the skeletal muscles leading to excess mobilization of lipids and proteins resulting in the production of reactive oxygen species and reduced antioxidants.^{4–6} Currently, urinary albumin is used for the diagnosis of nephropathy in patients with T2DM. However, some of the recent researchers reported that microalbuminuria is not a gold standard, sensitive and specific, biomarker for diagnosis of nephropathy in patients with T2DM, because it also elevates in other disorders like obesity, hypertension, different types of kidney problems, etc. In addition to that some of the patients diagnosed with microalbuminuric stage are directly converting into normoalbuminuric stage due to treatment modalities.^{7–9}

Along with that some of the recent studies found podocyte proteins to play an a important role in the early prediction and progression of nephropathy, as such urinary nephrin is the one of the anchoring proteins of foot process of the podocytes in the kidney, urinary excretion nephrin levels are used for early detection of nephropathy in patients with T2DM. Nephrin is the transmembrane protein of the immunoglobulin super family and also it is a very essential component of slit diaphragm between the foot processes of the podocytes in the kidney.¹⁰ The physiological action of nephrin is to maintain the size selectivity of the slit diaphragm and form a network of interdigitating foot processes of the podocytes in the kidney to prevent leaking of proteins.¹¹ Dysregulation of nephrin due to hyperglycemia in podocytes lead to nephrinuria particularly in T2DM patients with normoalbuminuric patients, preceding microalbuminuria.¹² However, very few studies have been done on urinary nephrin and also examined the relationship between urinary nephrin, hemoglobin A1c (HbA1C), urinary albumin, and estimated glomerular filtration rate (eGFR) in South Indian population and a study is required to evaluate sensitive and specific biomarker (urinary albumin and nephrin) for early detection of nephropathy in T2DM patients. Hence, in the present study we aimed to investigate the levels of urinary nephrin and correlate with urinary albumin and eGFR to predict early onset and progression of nephropathy in T2DM.

Materials and Methods

This is a case–control study conducted in Basaveshwara Institute of Medical Sciences and Research Centre, Chitradurga, Karnataka, India, and Arupadai Veedu Medical College, Puducherry, India, from 2018 to 2021. A total 120 subjects with age group of 30 to 70 years were included in the study. Among these, 80 T2DM subjects were again divided into two groups: group 2: 40 T2DM patients with normoalbuminuria (duration of DM < 5 years, HbA1c: 6–7.5%, and albumin creatinine ratio [ACR]: < 30 mg/dL); group 3: 40 T2DM patients with micro-albuminuria (duration of DM 5–10 years, HbA1c: 7.6–10.5%, and ACR: 30–299 mg/dL) (**– Fig. 1**). Additionally, we recruited



Fig. 1 Selection of study subjects.

age- and gender-matched healthy controls considered as group 1 (n = 40). All the T2DM subjects included in the study were diagnosed with different stages of nephropathy according to the American Diabetic Association criteria and Kidney Disease Improving Global Outcomes criteria^{13,14}; patients smoking, alcoholism, liver diseases, hypertension, cardiovascular, cerebrovascular, thyroid, peripheral vascular diseases, and those in treatment on thiazolidine and anti-inflammative were excluded from the study.

Seven milliliters of fasting (overnight 8-12 hours) venous blood sample and 3 mL of postprandial blood sample were collected from all the subjects after obtaining informed consent from. Two milliliters of blood sample was transferred into tube containing anticoagulant sodium fluoride, 2 mL transferred into tube containing anticoagulant ethylenediaminetetraacetic acid, and remaining 3 mL transferred into plain tube. Anticoagulant tubes were separated immediately and plain tube was allowed to rest for 10 minutes at room temperature for clotting. All the samples were centrifuged at 3,000 revolutions per minute (RPM) for 10 minutes, after centrifugation separated samples (plasma and serum) were transferred into properly labeled aliquots and stored at deepfreeze at -80°C until biochemical analysis was done. Along with the blood sample urine sample was also collected from all the subjects, centrifuged at 3,000 RPM for 10 minutes, and after centrifugation of urine sample, 1 mL was separated into labeled aliquots, stored at deepfreeze at -50°C and the remaining urine sample was immediately processed for urinary albumin and creatinine.

Plasma fasting blood sugar (FBS), postprandial blood sugar (PPBS), glycosylated HbA1c, serum urea, and creatinine were analyzed by using laboratory standard methods, eGFR was calculated by using Chronic Kidney Disease-Epidemiology Collaboration formula, urinary ACR was analyzed by using immunoturbidimetric method, and urinary nephrin was analyzed by using enzyme-linked immunosorbent assay.

Statistical Analysis

The data were statistically analyzed by using Microsoft Excel spreadsheets, IBM Statiscal Package for the Social Sciences (SPSS) Version 20.0, and Medcalc (Version 12.1, Ostend, Belgium). First, the normality of the three groups of data distribution was tested using the Kolmogorov-Smirnov test. Continuous variables are expressed as mean \pm standard deviation. Comparisons between the three groups were analyzed by using one-way analysis of variance (ANOVA). Pearson's correlation analysis was done to test the correlation of urinary nephrin with other biochemical parameters. Receiver operating characteristic (ROC) curves were constructed to study the diagnostic accuracy of the markers to identify DN in the normoalbuminuric group when compared with healthy controls. A cutoff value with the best combination of sensitivity and specificity was determined from the ROC curve. Marker positivity in the groups studied were calculated based on values above the upper limit of the 75% confidence intervals of the median values of the markers and the ROC-derived cutoff values. A p-value of less than 0.05 was considered as statistically significant.

Results

- Table 1 shows the plasma FBS, plasma PPBS, serum urea, serum creatinine, HbA1c, comparison of the routine biochemical parameters, and urinary biomarkers across the study groups. There is statistically significant elevation of plasma FBS, PPBS, HbA1c, serum urea, creatinine, urinary ACR, and nephrin in patients with T2DM compared with healthy controls, respectively, *p*-value of 0.0001.

- Table 2 shows the comparison of the routine biochemical parameters and urinary biomarkers across the study groups by one-way ANOVA. There is statistically significant elevation of plasma FBS, PPBS, HbA1c, serum urea, creatinine, urinary ACR, and nephrin in the two groups of T2DM

subjects compared with healthy controls, respectively, *p*-value of 0.0001. It was also observed that T2DM subjects with microalbuminuria have high level of urinary nephrin then T2DM subjects with normoalbuminuria, respectively, *p*-value of 0.0001, which shows statistically highly significant relation.

• Table 3 illustrates significantly increased levels of urinary nephrin along with increasing age in T2DM patients, respectively (p < 0.001).

- Table 4 shows the correlation of urinary nephrin with other routine biochemical parameters of the subjects by using Pearson's correlation analysis. We observed that urinary nephrin was positively significantly correlated with HbA1c and urinary ACR (r=0.520, 0.657, respectively, p=0.0001) and negatively correlated with eGFR (r=-0.539, respectively, p < 0.0001).

- Fig. 2 shows the graphical representation of urinary nephrin distribution in all the three groups of study subjects. Significantly elevated level of urinary nephrin in T2DM subjects with normoalbuminuria and microalbuminuria was observed then in healthy controls.

- Fig. 3 shows the distribution of urinary ACR levels in different groups of study subjects. T2DM subjects with microalbuminuria have significantly increased levels of urinary albumin when compared with T2DM subjects with normoalbuminuria and healthy controls.

- Fig. 4 shows eGFR levels in the two groups of T2DM subjects and healthy controls and was compared between groups. Significantly decreased eGFR levels were observed in T2DM subjects with microalbuminuria then in T2DM subjects with normoalbuminuria and healthy controls.

Table 1 Comparison of means of the anthropometric, biochemical, and urinary nephrin data among T2DM patients and healthy controls

Parameter	Group 1 (n=40)	Group 2 (n = 80)	<i>p</i> -Value
Age (y)	48.10 ± 9.56	49.57 ± 7.58	0.431 ^a
Height (feet)	3.66 ± 0.55	3.40 ± 0.52	0.022 ^a
Weight (kg)	74.47±11.97	82.55 ± 14.01	0.028 ^a
BMI (kg/m²)	20.42 ± 2.09	24.61 ± 4.54	0.0001 ^b
FBS (mg/dL)	90.63±7.31	156.17 ± 31.76	0.0001 ^b
PPBS (mg/dL)	114.70 ± 13.39	225.68 ± 73.64	0.0001 ^b
Serum urea (mg/dL)	23.07 ± 9.71	43.08 ± 18.44	0.0001 ^b
Serum creatinine (mg/dL)	0.99 ± 0.19	3.54 ± 2.71	0.0001 ^b
HbA1c (%)	4.31 ± 0.76	7.06 ± 1.30	0.0001 ^b
eGFR (mL/min)	91.43±13.93	57.01 ± 40.91	0.0001 ^b
Urinary ACR (mg/dL)	14.34±4.73	116.21 ± 107.07	0.0001 ^b
Urinary nephrin (ng/mL)	0.52 ± 0.20	2.48 ± 1.21	0.0001 ^b

Abbreviations: ACR, albumin creatinine ratio; BMI, body mass index; eGFR, estimated glomerular filtration rate; FBS, fasting blood sugar; HbA1c, hemoglobin A1c; PPBS, postprandial blood sugar; T2DM, type 2 diabetes mellitus.

Note: Group 1 is the healthy control, group 2 has type 2 diabetes.

^aNot significant at the 0.05 probability level.

^bHighly significant at the 0.05 probability level.

Parameter	Group 1 (n = 40)	Group 2 (duration of $DM < 5 y [n = 40]$)	Group 3 (duration of DM 5–10 y [n=40])	<i>p</i> -Value
Age (y)	48.10 ± 9.56	46.03 ± 5.91	53.10 ± 7.49	0.002 ^a
Height (feet)	3.66 ± 0.55	3.36 ± 0.52	3.43 ± 0.53	0.004*
Weight (kg)	74.47 ± 11.97	77.13 ± 14.24	87.97 ± 11.65	0.078
BMI (kg/m ²)	20.42 ± 2.09	23.01 ± 2.95	26.20 ± 5.29	0.0001 ^b
FBS (mg/dL)	90.63 ± 7.31	135.80 ± 15.49	176.53 ± 30.89	0.0001 ^b
PPBS (mg/dL)	114.70 ± 13.39	175.37 ± 20.71	276.00 ± 73.24	0.0001 ^b
Serum urea (mg/dL)	23.07 ± 9.71	26.97 ± 9.06	59.20 ± 8.48	0.0001 ^b
Serum creatinine (mg/dL)	0.99 ± 0.19	0.97 ± 0.21	6.11 ± 1.13	0.0001 ^b
HbA1c (%)	4.31 ± 0.76	6.07 ± 0.47	8.04 ± 1.11	0.0001 ^b
eGFR (mL/min)	91.43 ± 13.93	95.09 ± 19.71	18.93 ± 4.05	0.0001 ^b
Urinary ACR (mg/dL)	14.34 ± 4.73	15.41 ± 6.69	217.02 ± 47.49	0.0001 ^b
Urinary nephrin (ng/mL)	0.52 ± 0.20	1.64 ± 0.53	3.34±1.12	0.0001 ^b

Table 2 Comparison of the routine biochemical parameters and urinary biomarkers across the study groups

Abbreviations: ACR, albumin creatinine ratio; BMI, body mass index; DM, diabetes mellitus; eGFR, estimated glomerular filtration rate; FBS, fasting blood sugar; HbA1c, hemoglobin A1c; PPBS, postprandial blood sugar.

^aSignificant at the 0.05 probability level.

^bHighly significant at the 0.05 probability level.

Table 3 Urinary nephrin concentrations in different age groups of T2DM patients

Parameter	31–40 y (n = 11)	41–50 y (n = 36)	51–60 y (n = 27)	61–70 y (n=6)	<i>p</i> -Value
Urinary nephrin (ng/mL)	1.57 ± 0.86	3.27 ± 0.78	3.04 ± 1.41	3.56 ± 1.22	0.001 ^a

Abbreviation: T2DM, type 2 diabetes mellitus.

^aSignificant at the 0.05 probability level.

Table 4 Shows the Pearson's correlation between the urinary nephrin with other clinical parameters of nephropathy

Parameter		HbA1c	Urinary ACR	eGFR
Urinary nephrin (ng/mL)	r	0.520	0.657	-0.539
	<i>p</i> -Value	0.0001ª	0.0001ª	0.0001 ^a

Abbreviations: ACR, albumin creatinine ratio; eGFR, estimated glomerular filtration rate; HbA1c, hemoglobin A1c. Note: r is the rho factor.

^aSignificant at the level of 0.05.

Discussion

Diabetic kidney disease (DKD) is one of the major microvascular complications in patients with T2DM and nowadays more than 40% of the DKD patients progress to end-stage renal disease.¹⁵ In this condition damage of glomerular filtration barrier and tripartite system have fenestrated endothelial cells of glomerular basement membrane (GBM) and podocytes of kidney. The physiological action of these barriers is selective, filtration of water and solutes, and impermeable to leaking of macromolecules like proteins.^{16,17} In T2DM patients, hyperglycemia in the blood and hypoglycemia in tissues trigger lipolysis and proteolysis resulting in the production of reactive oxygen species and advanced glycation end products. These will bind to the advanced glycation end product receptors and trigger downstream signaling facilitating generation of free radicals, activating inflammatory cells, increasing synthesis of angiotensin II, and production of growth factors like vascular endothelial growth factors and transforming growth factor beta, finally leading to proteinuria.^{18,19}

Urinary albumin estimation is used for testing the functional capacity of kidney, when patients show less than 30 mg/dL of albumin it is called normoalbuminuria, 30 to 299 mg/dL of albumin is microalbuminuria, and more than 300 mg/dL is macroalbuminuria, it depends on the duration of diabetes and further determined by eGFR.²⁰ Some of the recent studies reported that urinary microalbuminuria is not a



Fig. 2 Distribution of urinary nephrin levels in three groups of study subjects.



Fig. 3 Distribution of urinary albumin creatinine ratio (ACR) levels among the study subjects.



Fig. 4 Distribution of estimated glomerular filtration rate (eGFR) levels among the study subjects.

sensitive and specific biomarker for detection of nephropathy in T2DM patients because it has low sensitivity and larger variability. Our study has shown statistically significant elevation of urinary ACR levels in T2DM patients with microalbuminuria when compared with T2DM patients with normoalbuminuria and healthy controls, *p*-value less than 0.0001 (**-Table 2**). However, earlier studies have reported that mean arterial blood pressure, plasma concentration of FBSs, HbA1c, aldosterone, atrial natriuretic peptide, vasopressin, and angiotensin II are also risk factors for proteinuria. Additionally, recent studies revealed that T2DM patients with microalbuminuria reverted to normoalbuminuria.^{21–23}

If DN can be detected before the appearance of microalbuminuria, early intervention can be started to halt or reverse the process. Glomerular biomarker of renal injury is being studied as alternative markers for prediction of DN risk. Some of the recent studies found sensitive and specific biomarker for early detection of nephropathy in T2DM patients. Nephrin is a protein located in the podocytes of GBM of the kidney. Nephrin plays a vital role in podocytes and slit diaphragm to prevent excretion of protein in urine, but minimal structural alterations in podocytes lead to excretion of nephrin in urine prior to proteinuria. The studies are very limited and some of the studies found significantly elevated levels of urinary nephrin in T2DM and other studies reported the urinary nephrin levels were decreased in patient with T2DM.²⁴⁻²⁷ The present study also analyzed urinary nephrin, and significantly elevated levels were observed in patients in both groups of T2DM when compared with healthy controls and we also found significantly increased levels of urinary nephrin in T2DM patients with normoalbuminuria when compared with healthy controls. Similarly, other studies also reported podocytes are present in the GBM of kidney and play a good role in maintaining glycocalyx of glomerulus and maintained the slit diaphragm size by specific proteins like nephrin and they also suggested that elevated levels of urinary nephrin were useful for early detection of nephropathy in T2DM patients.

We also found that urinary nephrin was positively correlated with blood sugars, glycosylated hemoglobin, urea, creatinine, and urinary albumin (**-Table 4**). Similarly, another recent study also reported that association of proximal tubule dysfunction with podocytes damage and dysfunction of proximal convoluted tubule proteins could be validated as an early diagnosis and progression of nephropathy in patients with T2DM.^{28,29} Elevated levels of urinary nephrin in T2DM patients with normoalbuminuria stage and increased levels of more than two times in T2DM patients with microalbuminuria showed disease progression toward end-stage kidney disease.

The diagnostic utility of urinary ACR, eGFR, and urinary nephrin for the detection of early DN was compared between group 1 and group 2 by ROC curve analysis (**-Table 5**). Among these urinary ACR and eGFR there was no significant area under the curve (AUC), with sensitivity ranging from 43.33 to 46.67 and specificity 76.67 to 73.33, respectively, *p*-values 0.68 and 0.61 and the urinary nephrin showed a

Parameter	AUC	95% CI for AUC	Sensitivity (%) 95% Cl	Specificity (%) 95% Cl	p-Value
Urinary ACR (mg/dL)	0.532	0.398-0.662	43.33	76.67	0.683 ^a
eGFR (mL/min)	0.538	0.405-0.668	46.67	73.33	0.616 ^a
Urinary nephrin (ng/mL)	0.993	0.928-1.000	100.00	93.33	0.0001 ^b

Table 5 ROC curve analysis of urinary ACR, eGFR, and urinary nephrin

Abbreviations: ACR, albumin creatinine ratio; AUC, area under the curve; CI, confidence Interval; eGFR, estimated glomerular filtration rate; ROC, receiver operating characteristic.

^aNot significant.

^bHighly significant.

statistically significant AUC with sensitivity ranging from 100% and specificity from 93%, respectively, while *p*-value is less than 0.0001 (**~Fig. 5**).

Univariate linear regression analysis was done by using eGFR as dependent variable and urinary ACR and urinary nephrin as independent variables and showed a significant inverse association. Along with that we also analyzed urinary ACR as dependent variable and urinary nephrin as the independent variable and showed a direct association. The scatter plots showed a relationship between eGFR, urinary ACR, and urinary nephrin (r=0.87, 0.65, and 0.66, respectively, *p*-values < 0.001) (**-Fig. 6**). Hence, based on these findings urinary nephrin is identified as the best marker for diagnosis of DN then urinary ACR.



Fig. 5 Receiver operating characteristic (ROC) curve analysis of estimated glomerular filtration rate (eGFR), urinary albumin creatinine ratio (ACR), and urinary nephrin.



Fig. 6 Univariate linear regression analysis for early diagnosis of diabetic nephropathy.

Conclusion

The outcome of this study showed that urinary nephrin might be a better marker than urinary ACR for predicting early onset of nephropathy in patients with T2DM, because urinary nephrin levels elevated and there was no change in urinary ACR levels in T2DM patients with normoalbuminuria.

Authors' Contributions

V.G. contributed to study design, selection of subjects, collection of samples, processing, data analysis, and writing manuscript. M.M. contributed to assisting in the study design, selection of subjects, collection of samples, processing, data analysis, and writing manuscript. V.K.G. contributed to assisting in the study design, selection of subjects, collection of samples, processing, data analysis, and writing manuscript.

Ethical Approval

The subcommittee constituted by institutional ethical committee, Basaveshwara Medical College and Hospital, Chitradurga, Karnataka, India has reviewed and discussed my protocol on 28/03/2019 to conduct the research project (Ref. BMC&H/IEC/2018–2019/07).

Funding None.

Conflict of Interest None declared.

References

- 1 Badr A, Muhammad IU, Muhammad A, et al. Determination of serum adiponectin levels in type 2 diabetes patients of the Saudi population in the Al-Jouf region. Rom J Diabetes Nutr Metab Dis 2020;3:1–10
- 2 Salam RS, Laikangbam S, Tina D, et al. Adiponectina and body mass index in type 2 diabetes mellitus. Int J Contemp Med Res 2020;4:1–4
- 3 Raghuram N, Parul B, Akshay A, et al. Prevalence of diabetes and its determinants in the young adults Indian population-call for yoga intervention. Front Endocrinol 2020;11:1–9
- 4 Uwaezuoke SN. The role of novel biomarkers in predicting diabetic nephropathy: a review. Int J Nephrol Renovasc Dis 2017; 10:221–231
- 5 Vlad A, Vlad M, Petrica L, et al. Therapy with atorvastatin versus rosuvastatin reduces urinary podocytes, podocyte-associated molecules, and proximal tubule dysfunction biomarkers in patients with type 2 diabetes mellitus: a pilot study. Ren Fail 2017;39(01):112–119
- 6 Abdelbassit E, Maha Abdelmoneim B, Somia Abdelhameed B, et al. Urinary podocin level as a predictor of diabetic kidney disease. J Nephropathol 2019;8(03):26
- 7 Rigalleau V, Lasseur C, Raffaitin C, et al. Normoalbuminuric renalinsufficient diabetic patients: a lower-risk group. Diabetes Care 2007;30(08):2034–2039
- 8 Popovic D, Lalic K, Jotic A, et al. The inflammatory and hemostatic cardiovascular risk marke during acute hyperglycemic crisis in type 1 and type 2 diabetes. J Med Biochem 2019;38(02):126–133
- 9 Kostovska I, Tosheska-Trajkovska K, Topuzovska S, et al. Urinary nephrin is earlier, more sensitive and specific marker of diabetic nephropathy than microalbuminuria. J Med Biochem 2020;39 (01):83–90
- 10 Pettersson-Fernholm K, Forsblom C, Perola M, Groop PHFinn-Diane Study Group. Polymorphisms in the nephrin gene and diabetic nephropathy in type 1 diabetic patients. Kidney Int 2003;63(04):1205–1210

- 11 Jim B, Ghanta M, Qipo A, et al. Dysregulated nephrin in diabetic nephropathy of type 2 diabetes: a cross sectional study. PLoS One 2012;7(05):e36041
- 12 American Diabetes Association. Standards of medical care in diabetes. Diabetes Care 2017;40(01):S1–S135
- 13 Saran R, Robinson B, Abbott KC, et al. US renal data system 2016 annual data report: epidemiology of kidney disease in the United States. Am J Kidney Dis 2017;69(3, Suppl 1):A7–A8
- 14 National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. Am J Kidney Dis 2002;39(2, Suppl 1):S1–S266
- 15 Drummond K, Mauer MInternational Diabetic Nephropathy Study Group. The early natural history of nephropathy in type 1 diabetes: II. Early renal structural changes in type 1 diabetes. Diabetes 2002;51(05):1580–1587
- 16 Ponchiardi C, Mauer M, Najafian B. Temporal profile of diabetic nephropathy pathologic changes. Curr Diab Rep 2013;13(04): 592–599
- 17 Tan AL, Forbes JM, Cooper ME. AGE, RAGE, and ROS in diabetic nephropathy. Semin Nephrol 2007;27(02):130–143
- 18 Veron D, Bertuccio CA, Marlier A, et al. Podocyte vascular endothelial growth factor (Vegf₁₆₄) overexpression causes severe nodular glomerulosclerosis in a mouse model of type 1 diabetes. Diabetologia 2011;54(05):1227–1241
- 19 Kondapi K, Kumar NL, Moorthy S, Silambanan S. A study of association of urinary nephrin with albuminuria in patients with diabetic nephropathy. Indian J Nephrol 2021;31(02):142–148
- 20 Gross JL, de Azevedo MJ, Silveiro SP, Canani LH, Caramori ML, Zelmanovitz T. Diabetic nephropathy: diagnosis, prevention, and treatment. Diabetes Care 2005;28(01):164–176

- 21 Hong CY, Chia KS. Markers of diabetic nephropathy. J Diabetes Complications 1998;12(01):43–60
- 22 Kania K, Byrnes EA, Beilby JP, Webb SA, Strong KJ. Urinary proteases degrade albumin: implications for measurement of albuminuria in stored samples. Ann Clin Biochem 2010;47(Pt 2):151–157
- 23 Ma RX, Xu Y, Zhang J, Li YS, Liu LQ. Triptolide combined with irbesartan synergistically blocks podocyte injury in a type 2 diabetes rat model [in Chinese]. Zhonghua Nei Ke Za Zhi 2012; 51(02):117–122
- 24 Kondapi K, Santhi S, Swathy M. Diagnostic sensitivity of biomarkers in assessing the occurrence and progression of diabetic nephropathy. Eur J Mol Clin Med 2021;8(03):1–18
- 25 Yadav BK, Adhikari S, Gyawali P, Shrestha R, Poudel B, Khanal M. Use of protein: creatinine ratio in a random spot urine sample for predicting significant proteinuria in diabetes mellitus. Nepal Med Coll J 2010;12(02):100–105
- 26 Ng DP, Tai BC, Tan E, et al. Nephrinuria associates with multiple renal traits in type 2 diabetes. Nephrol Dial Transplant 2011;26 (08):2508–2514
- 27 Toyoda M, Suzuki D, Umezono T, et al. Expression of human nephrin mRNA in diabetic nephropathy. Nephrol Dial Transplant 2004;19(02):380–385
- 28 Petrica L, Vlad A, Gluhovschi G, et al. Proximal tubule dysfunction is associated with podocyte damage biomarkers nephrin and vascular endothelial growth factor in type 2 diabetes mellitus patients: a cross-sectional study. PLoS One 2014;9(11):e112538
- 29 Bobkova IN, Shchukina AA, Shestakova MV. Assessment of nephrin and podocin levels in the urine of patients with diabetes mellitus. Nephrology (Saint-Petersburg) 2017;21(02):33–40