



# Calcific Discitis Seems to be a Common Incidental Finding in Adults

Johannes Gossner<sup>1</sup>

<sup>1</sup>Department of Diagnostic and Interventional Radiology, Evangelisches Krankenhaus Göttingen-Weende, An der Lutter 24, Göttingen, Germany

Address for correspondence Johannes Gossner, MD, Department of Diagnostic and Interventional Radiology, Evangelisches Krankenhaus Göttingen-Weende, An der Lutter 24, 37074 Göttingen, Germany (e-mail: johannesgossner@gmx.de).

Indian J Radiol Imaging 2023;33:230–231.

## Abstract

Calcific discitis seems to be a rare cause of back pain in adults. Imaging shows a calcification of the nucleus pulposus with extension through the endplates on computed tomography. This can be accompanied by bone marrow edema on magnetic resonance imaging. In a retrospective review of 150 patients, 4 cases of calcific discitis were found (2.8%). None of the patients reported about back pain. Therefore, it seems that symptomatic cases of calcific discitis are a rare occurrence in contrast to the quite frequent incidental finding of asymptomatic cases on imaging. Knowledge of the various imaging appearances of calcific discitis is necessary for radiologists, because especially in acute cases with substantial bone marrow edema, this benign entity may be confused with infectious spondylodiscitis or malignancy.

## Keywords

- ▶ spinal imaging
- ▶ calcific discitis
- ▶ prevalence
- ▶ incidental finding

## Background

Calcific discitis seems to be a rare cause of back pain in adults, with around 40 published cases.<sup>1,2</sup> Imaging shows a calcification of the nucleus pulposus with extension through the endplates on computed tomography. This can be accompanied by bone marrow edema on magnetic resonance imaging. Calcific discitis is more prevalent in the thoracic spine.<sup>2</sup> The prognosis is favorable and pain usually resolves with symptomatic treatment. Sometimes follow-up imaging shows a resolution of the calcifications.<sup>2</sup> To the best of our knowledge, the prevalence of calcific discitis in adults has not been studied previously.

## Methodology


A retrospective review of chest computed tomography (CT) scans performed at our department during January 2022 was performed. All patients undergoing chest imaging on an 80-slice CT scanner (Canon Prime, Canon Medical Sys-

tems, Tokyo, Japan) were included in the study. Patients were scanned for different indications (such as cancer, pneumonia, interstitial lung diseases, or pulmonary embolism) and the use of contrast media was not mandatory. The sample consisted of 85 male and 65 female patients with a mean age of 69.5 years (range: 31–99 years). The reconstructed sagittal 3 mm slices of the thoracic and upper lumbar spine were transferred to a medical workstation and reviewed in a standard bone window setting using the departmental digital picture archive (SynedraView; Synedra Information Technology, Innsbruck, Austria). The diagnosis of calcific discitis was made if calcifications of the nucleus pulposus with extension through adjacent endplates were found. To rule out other causes of intervertebral disc calcifications (like degenerative changes or changes with ankylosing spinal disease), only cases with these typical imaging findings were excluded. All cases with fusion of the segment were excluded. In cases with calcific discitis, a chart review was performed.

article published online  
December 7, 2022

DOI <https://doi.org/10.1055/s-0042-1759482>.  
ISSN 0971-3026.


© 2022. Indian Radiological Association. All rights reserved.  
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)  
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



**Fig. 1** Two examples of calcific discitis of the thoracic spine showing the typical imaging finding of nucleus pulposus calcification with migration through the adjacent vertebral endplates.

## Results and Discussion

In our sample of 150 patients, 4 cases of calcific discitis were found (2.8%) (►Fig. 1). Calcific discitis was found in three women and one man; these patients had a mean age of 77 years. In two patients, the changes were multisegmental. In the retrospective review of the medical records, none of the cases spontaneously reported about back pain at admission. This small study observed a prevalence of 2.8% of calcific discitis on CT, which is contradictory to the low number of reported symptomatic cases in the literature. Given the obvious limitations of a retrospective chart review, none of the patients reported about back pain on admission. Therefore, it seems that symptomatic cases of calcific discitis are a rare occurrence in contrast to the quite frequent incidental finding of asymptomatic cases on imaging. This is not an unknown phenomenon on spinal medicine and known from symptomatic Schmorl nodules.<sup>3</sup> It may be speculated that the migration of the calcified nucleus pulposus into the endplates causes self-limiting pain in most patients, which impedes further workup and detection of symptomatic cases on imaging. Nonetheless, knowledge of the various imaging appearances of calcific discitis is necessary for radiologists, because especially in acute cases with substantial bone marrow edema, this benign entity may be confused with infectious spondylodiscitis or malignancy<sup>4</sup> (►Fig. 2). The pathogenesis of calcific discitis remains unclear. A more recent hypothesis suggests secondary interruption or diminished blood supply of the nucleus pulposus, for example, after trauma, inflammation, or fusion.<sup>5</sup> It has to be noted that there are other forms of intervertebral disc calcifications. Nucleus pulposus calcifications are often seen in patients with fused intervertebral segments (i.e., in ankylosing spondylitis



**Fig. 2** Imaging of a patient presenting with back pain and suspected malignancy on bone scintigraphy. Computed tomography (on the left) shows the typical imaging appearance of calcified discitis. Magnetic resonance imaging (on the right) shows reactive bone marrow edema due to the migration of the calcified nucleus pulposus through the endplates.

or diffuse idiopathic skeletal hyperostosis). However, in patients with degenerative changes, calcifications of the annulus fibrosus seem to be more common.<sup>6,7</sup>

## Conclusion

Calcific discitis seems to be a quite frequent incidental finding on CT of the spine. Knowledge of this entity is important to establish proper differential diagnosis.

### Conflict of Interest

None declared.

## References

- Schulze P, O'Connor C, Anand N, Wynne P. A rare cause of acute back pain in a young adult: case report and discussion of calcific discitis. *J Radiol Case Rep* 2020;14(10):1–9
- Iampreechakul P, Lertbutsayanukul P, Suanprasert N. Acute calcific discitis or symptomatic Schmorl's nodule of the upper thoracic spine in an adult: a case report and literature review. *Asian J Neurosurg* 2019;14(03):1021–1029
- Hershkovich O, Koch JEJ, Grevitt MP. Schmorl node- a cause of acute thoracic pain: a case report and pathophysiological mechanism. *Int J Spine Surg* 2020;14(03):441–446
- Shah A, Botchu R, Grainger MF, Davies AM, James SL. Acute symptomatic calcific discitis in adults: a case report and review of literature. *Skeletal Radiol* 2015;44(12):1819–1824
- Mittal P, Saggar K, Sandhu P, Gupta K. Case report: Acute calcific discitis with intravertebral disc herniation in the dorsolumbar spine. *Indian J Radiol Imaging* 2010;20(03):205–207
- Dussault RG, Kaye JJ. Intervertebral disk calcification associated with spine fusion. *Radiology* 1977;125(01):57–61
- Chanchairujira K, Chung CB, Kim JY, et al. Intervertebral disk calcification of the spine in an elderly population: radiographic prevalence, location, and distribution and correlation with spinal degeneration. *Radiology* 2004;230(02):499–503