



Circumferential Endoscopic Submucosal Dissection of long-segment Barrett's Esophagus with Multifocal High-Grade Dysplasia

Sukrit Sud¹ Smruti Ranjan Mishra¹ Randhir Sud¹

¹Institute of Digestive & Hepatobiliary Sciences, Medanta the Medicity, Gurugram, Haryana, India

Address for correspondence Randhir Sud, MD, DM, Institute of Digestive and Hepato-biliary Sciences, Medanta the Medicity, Sector 38, Gurugram 12200, Haryana, India (e-mail: drsud@gmail.com).

J Digest Endosc 2022;13:254–256.

Abstract

In a case of long-segment Barrett's esophagus with multifocal high-grade dysplasia with multiple comorbidities, circumferential endoscopic submucosal dissection was performed. Following the procedure, the esophageal stricture was also managed.

Keywords

- ▶ Barrett's esophagus
- ▶ endoscopic submucosal dissection
- ▶ esophageal stricture
- ▶ high-grade dysplasia

Case Report

A 79-year-old male patient presented with persistent retrosternal burning and high-volume reflux for the last 5 years. His reflux symptoms had become refractory to proton pump inhibitors and prokinetic agents since the past 2 years. He had significant comorbidities as he had panhypopituitarism and a history of stroke with left carotid artery stenosis. He underwent an esophagogastroduodenoscopy (EGD) at our center that revealed a long-segment circumferential Barrett's esophagus with irregular surface and vascular pattern (C8M8, Paris 0-IIb) (▶ **Fig. 1**). Multiple biopsies were taken as per the Seattle protocol that revealed Barrett's esophagus with multifocal high-grade dysplasia.

A multidisciplinary discussion about treatment options was done, and oesophagectomy was considered as high risk because of associated comorbidities. He underwent a CECT chest and abdomen, which revealed no significant mediastinal or abdominal lymphadenopathy. We decided to do a circumferential endoscopic submucosal dissection (ESD) of Barrett's segment in view of multiple comorbidities. The pocket-creation ESD method was employed.¹ We used a hybrid knife, VIO

200D electrogenerator (ERBE) with a forward-viewing endoscope (GIF-HQ190; Olympus, Tokyo, Japan) and a carbon dioxide (CO₂) insufflator (UCR; Olympus, Tokyo, Japan). We created an anterior and a posterior tunnel by submucosal dissection and these were subsequently joined, hence an en-bloc resection could be achieved. The procedure took 150 minutes to complete, and the 8-centimeter long circumferential specimen was sent for histopathological examination (▶ **Fig. 2-4**, ▶ **Video 1**). Histopathological examination confirmed an R0 resection of a long-segment Barrett's esophagus with multifocal high-grade dysplasia and foci of T1a adenocarcinoma. The patient underwent a gastrografin swallow study on a subsequent day and could be started on a liquid diet on that day.

Video 1

Final esoesd. Online content including video sequences viewable at: <https://www.thieme-connect.com/products/ejournals/html/10.1055/s-0042-1759743>.

DOI <https://doi.org/10.1055/s-0042-1759743>.
ISSN 0976-5042.

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India




Fig. 1 Retroflexion view showing long-segment Barrett's esophagus.




Fig. 4 Circumferential resected specimen with scale to measure (8 cm).



Fig. 2 Pocket-creation ESD using a hybrid knife.




Fig. 5 Post ESD esophageal stricture formation at 1 month.




Fig. 3 Proximal margin of the cut surface.




Fig. 6 Esophageal stricture CRE dilatation.

The rate of stricture occurrence after near-circumference or whole-circumference ESD has been reported to be between 60 and 100%.² The patient underwent an OGD after 1 month that revealed a concentric short-segment stricture along the proximal margin of dissection (► **Fig. 5**). He underwent balloon dilatation of the stricture with an injection of triamcinolone acetonide (40 mg/mL diluted as 1:1 with saline solution) using a 23-gauge, 5-mm long sclerotherapy needle in aliquots of 0.5 mL in each quadrant (► **Fig. 6**).³ The patient was already on oral corticosteroids for panhypopituitarism, and fortunately




Fig. 7 Follow-up after 3 months.

did not have a stricture recurrence. He underwent another EGD after 3 months that revealed reappearance of healthy squamous mucosa throughout the resected surface (► **Fig 7**).

Conflict of Interest
None declared.

References

- 1 Harada H, Murakami D, Suehiro S, et al. Water-pocket endoscopic submucosal dissection for superficial gastric neoplasms (with video). *Gastrointest Endosc* 2018;88(02):253–260
- 2 Martínek J, Juhas S, Dolezel R, et al. Prevention of esophageal strictures after circumferential endoscopic submucosal dissection. *Minerva Chir* 2018;73(04):394–409
- 3 Kochhar R, Makharia GK. Usefulness of intralesional triamcinolone in treatment of benign esophageal strictures. *Gastrointest Endosc* 2002;56(06):829–834