

Biologics in Rhinology – Forthcoming Personalized Concepts: the Future Starts Today



Author

Adam M. Chaker

Affiliation

Department of Otolaryngology and Center for Allergy and Environmental Medicine, Klinikum rechts der Isar, Technical University of Munich, Germany

Key words

Rhinitis, sinusitis, rhinosinusitis, CRS, CRSwNP, nasal polyps, biologics, precision medicine, personalized medicine

Bibliography

DOI <https://doi.org/10.1055/s-0043-123484>

Laryngo-Rhino-Otol 2018; 97: S164–S184

© Georg Thieme Verlag KG Stuttgart · New York

ISSN 0935-8943

Correspondence

Dr. med. Adam M. Chaker

Department of Otolaryngology

Center for Allergy and Environmental Medicine

University Hospital rechts der Isar

Ismaninger Straße 22

81675 München

Germany

adam.chaker@tum.de

ABSTRACT

Chronic diseases of the nose and the paranasal sinuses are most common, frequently associated with bronchial asthma, and result in substantial reduction of quality of life. Despite optimal treatment according to guidelines, approx. 20% of the patients will report inadequate control of symptoms. Apart from well-established surgical and conservative approaches in therapy, new therapeutic antibodies are available that aim specifically at pathophysiological targets. The optimal allocation of effective therapy for patients using appropriate biomarkers at the most suitable timepoint is the hallmark of stratified medicine and an important perspective in ENT.

Contents

Abstract	S164	2.1 Physiology of the nasal mucosa and the integrated mucosal immune system	S168
1. Biologicals and new treatment options of stratified medicine	S165	2.2 United Airways	S168
1.1 Biological medical substances: definition	S165	2.3 Entities and epidemiology of diseases of the nasal mucosa and the paranasal sinuses	S169
1.2 Development and historical milestones	S165	2.4 Endotypes of chronic airway inflammations	S171
1.3 First biologicals, hybrid molecules, and concept-related advantages and disadvantages	S165	2.5 Stratified downstream prevention	S172
1.4 Immunogenicity	S165	3. State-of-the-art	S172
1.5 Anaphylaxis against biologics: the example of cetuximab	S165	4. Biologicals in clinical trials with rhinological diseases	S173
1.6 Intended immunological effects and temporary immune phenomena	S166	4.1 Allergic rhinitis	S173
1.7 Biosimilars	S166	4.2 Chronic rhinosinusitis	S174
1.8 Small molecules and DNazyme	S167	4.3 Examples of rare indications	S176
1.9 Costs	S167	4.4 Biomarkers	S177
2. Rhinology: Epidemiology and pathophysiological concepts	S168	5. Conclusion and general reflections on personalized medicine	S177
		References	S178

1. Biologicals and new treatment options of stratified medicine

The design of biological pharmaceuticals is based on a profound mechanistic understanding of disease processes. Generally they consist of large-molecular substances that were synthesized by living organisms. Most commonly, biologics mediate their effects via selective binding of cytokines or their specific receptors [1]. In this introduction, first development and classification, biochemical specifics, immunological effects, current trends and finally characteristic side effects will be described.

1.1 Biological medical substances: definition

The European Medical Agency (EMA) defines biologics (or biological medicinal products) as biopharmaceutical medicinal products that are applied *in vivo* as biological medicinal substance and will usually be applied as therapeutic agents. Furthermore, they include biotechnologically produced active substances, e. g., therapeutic antibodies and recombinant proteins but also vaccines and allergens, blood and plasma products as well as recombinantly produced alternatives [2]. In the daily language of life-science and medicine, the term of biologics was established mainly for therapeutic antibodies and rarely also for recombinant therapeutic proteins.

1.2 Development and historical milestones

Based on mechanistic studies, molecular biology and genetic basic research, therapeutic target structures were developed in the 1990ies and validated in animal models as well as in human translational models before clinical testing. By means of a procedure developed by Milstein and Köhler (Nobel Prize for Medicine in 1984) [3], therapeutic antibodies may be produced by immortalized hybrid B cells from myeloma cell cultures for nearly every target structure. Further biotechnological procedures allowed synthesizing, validating, and approving new biopharmaceuticals, e. g., recombinant insulins (first approval by the FDA in 1982) [4]. This development was further fueled by the transfusion scandals of the 1980ies, accidental transmission of HIV or HCV to hemophilia patients and the contamination by Creutzfeld-Jakob in the context of pituitary extracts for growth hormone substitution. Also recombinant coagulation factors [5, 6] and growth hormone (approval by the FDA in 1985) [7, 8] were produced by means of biotechnology.

1.3 First biologicals, hybrid molecules, and concept-related advantages and disadvantages

In particular for usage in oncology and autoimmunity, promising monoclonal neutralizing antibodies have been constructed, fusion proteins with binding capacity (etanercept), with or without intrinsic activity (IL-4 mutein), receptor antagonists, bi- and trispecific antibodies that may bind different target structures and at the same time activate for example T cells. In the former dual immunological conception of Th1/Th2 inflammations [9, 10], recombinant cytokines were intended to restore the inflammatory balance, e. g., by application of recombinant IL-12 in cases of asthma [11] or e. g., IL-10 [12] and IL-11 [13] for psoriasis. In contrast as for example to interferon therapy for treatment of multiple sclerosis or viral hepatitis, those trials did not reach their endpoints consistently and someti-

mes showed relevant side effects. Hence, these therapeutic approaches have not been pursued.

Beside the possible variety of biologically potentially applicable therapeutic proteins, at the same time a high range of biotechnological expression systems has been developed [14]. After initial prokaryotic systems, successively eukaryotic expression systems have been developed that made available also larger and glycosylated proteins with complex tertiary and quaternary structure in reproducibly high and therapeutic quality. Even with regard to optimization of therapeutic antibodies to target epitopes and the development of extraction processes, enormous pioneering work was done [15–19].

While biologically active human insulin has a molecular weight of approx. 5,000 Daltons, the growth hormone somatotropin has a molecular weight of about 22,000 Daltons [20]. A monoclonal antibody has about 150,000 Daltons; for comparison, the molecular weight of acetylsalicylic acid is approx. 180 Daltons. The first therapeutic antibodies were developed as chimeric proteins with relatively high murine or other xenobiological portions and increasingly humanized. This led to a significantly improved tolerance, effectiveness, and safety for the patients. This development is also reflected in the nomenclature of the biologics: The anti-CEA antibody arcitumomab approved in 1996 was only murine, the therapeutic TNF- α antibody Infliximab, approved in 1999, was a chimeric antibody, the herceptin antibody trastuzumab and the anti-IgE antibody omalizumab were humanized antibodies, and the TNF- α antibody adalimumab was completely human (► Fig. 1 and [21]).

1.4 Immunogenicity

Beside an intended immunological effect, usually an undesired immunological reaction occurs on biologically active and mostly exogenous substances in the recipient or host organism, which is called immunogenicity. The murine or xenobiological portions on the one hand and the glycosylation pattern on the other hand determine significantly the immunogenicity of the active biological substance. This leads to for example ADAs (anti-drug antibodies) that at best are neutralizing and lead to ineffectiveness of the applied biological drug, but also potentially to immunological phenomena such as autoimmune hemolysis, cytokine release syndrome, mast cell activation, immune complex hypersensitivity and serum disease. Examples in this context are the relatively high rates of reactions against infliximab or rituximab, which consist of a significant murine proportion [22, 23]. Finally, those biologically active substances are complex, large, biotechnologically produced pharmaceuticals. Methodological details during production as well as cooling, light, pH, transportation, and application may lead to small molecular changes with potential alterations, such as loss of effectiveness, molecular aggregates, and increased immunogenicity. Furthermore, cross-reacting epitopes may lead to fatal anaphylaxis, especially when the antigens share amino acid sequences with IgE epitopes, e. g., cetuximab.

1.5 Anaphylaxis against biologics: the example of cetuximab

In 2003, the EGF receptor antibody cetuximab was approved in the USA (FDA) and in 2004 in Europe (at that time: EMEA), first for the treatment of advanced colon cancer. Since the publication of pivotal trials on combination therapy, it is also applied for advanced head




Fig. 1 Depiction of the international non-proprietary names (INN) for the development of humanized and human antibodies (Creative Commons Licence, cited from [21], Paul W.H.I. Parren et al., 2017, mABS).

and neck carcinomas [24–26]. Already during approval studies in the early 2000ies, regional differences became apparent in the USA with regard to the incidence of anaphylactic reactions. In 2007, O’Neil could show that the first administration of the drug had a global probability of less than 3 % to develop anaphylactic reactions. In the middle southeast of the USA, however, e. g., in Virginia, this rate amounted to up to 20 % [27, 28]. This reaction is of completely different nature compared to the relatively frequent cutaneous reactions [29]. In 2013, Platt-Mills and colleagues could clarify the mechanism of sensitization and association with tick bites and allergy against red meat [30, 31]. The regional distribution of a tick species, the so-called lone-star tick, leads to the development of galactose-alpha-1.3-galactose (alpha-GAL) specific IgE antibodies after tick bite. Alpha-GAL is a nearly ubiquitously expressed oligosaccharide of glycoproteins of the cell surface in all non-primate mammals, prosimians, and new world monkeys. In primates (old world monkeys and humans), about 1 % of the circulating IgG pool correspond to alpha-GAL antibodies, and alpha-GAL is a relevant biological obstacle of a simple xenotransplantation [32]. Since 2014, commercial test systems are available that can be applied to determine specific IgE against alpha-GAL. The exclusion of sensitization to avoid anaphylactic reactions prior to application of humanized antibodies can be life-saving for patients with high-risk profiles. High-risk patients are individuals with many tick bites (e. g., forestry professionals), with cat allergy, and pork-cat syndrome [33]. The determination of alpha-GAL in this high-risk cohort is a showcase of personalized medicine according to the WHO definition: avoiding side effects by specific characterization of patients.

1.6 Intended immunological effects and temporary immune phenomena

The effect that is inherent to most therapeutic antibodies is the impact on immunological processes, e. g., by antagonizing or eliminating messenger substances of the inflammation cascade (e. g., anti TNF α , anti-IL-1, anti-IL-5) (► Fig. 2).

The transition from undesired to intended immunological effects is fluid. Bi- or even trispecific antibodies such as for example catumaxomab that are applied for advanced oncologic diseases such as ma-


lignant ascites and hereby bind to the Fc receptor and to the activating T cell receptor CD3 and to the epithelial cell adhesion molecule EpCAM, activate the immune system in a targeted way and beside the antineoplastic main effect, they generate an inflammatory co-activation by cytokine release, however, with significant side effects.

Another intended effect is the inhibition of regulatory immunological processes on T cell level in vivo by checkpoint kinase inhibitors such as the inhibition of CTLA-4 by ipilimumab as well as inhibition of PD-1 by nivolumab or pembrolizumab (overview to be found in [34]).

In some cases, autoimmune phenomena or side effects caused by biologics, as for example rituximab, are causally co-treated de facto on B cell level by further continuously administering the same drug [35].

1.7 Biosimilars

The complexity and size of the molecules, their different expression systems and production processes do not allow real generics of biological products even after expiration of patent protection. Therefore, the concept of biosimilars was developed [36]. Ledford [37] defined biosimilars as an inexact copy of a reference substance. The major question from the immunological point of view is: How similar do biosimilars have to be compared to the original substance, or more bluntly: how similar is similar? The instrumental definition of the EMA considers a biosimilar as highly similar to the original substance in all relevant aspects, i. e., effective mechanism, safety, production process. In the European Union, at least one clinical trial is required from the manufacturers of biosimilars to document the biological equivalence which implicates a relevant switchability of the biosimilar with the originator substance. In addition, extensive immunogenicity data and data on the production process have to be made available. From this regulatory context it is mandatory for the manufacturers to commit themselves to permanent pharmacovigilance and extensive quality management. If the biological equivalence is confirmed, the biosimilar may be introduced onto the market under the INN name. Based on this certified equivalence, the approval authorities can allow an extrapolation, i. e., based on the



► **Fig. 2** On the left, a mucosal immune reaction against allergens under the influence of further pathogens in the sense of sensitization. Hereby, alarmins (IL-25, IL-33, and TSLP), proinflammatory cytokines (e. g., IL-4, IL-5, IL-13), antigen-presenting cells (APC), innate lymphoid cells (ILC) as well as T helper cells and B cells are depicted. On the right, the situation of a re-exposition is displayed involving mast cells as well as cellular late stage reaction with the influx of eosinophilic granulocytes and T helper cells under secretion of multiple cytokines and use of intercellular signaling pathways (according to [1], Boyman et al., Allergy, 2015). The mechanistic characterization of these different inflammatory pathways allows a targeted therapy by e. g., therapeutic antibodies for the processes in the epithelium as well as in the subepithelial stroma (licensed by RightsLink/John Wiley and Sons).

available data, it is assumed that the biosimilar works also in all other approved indications, equivalently to the originator substance [38, 39]. The switchability and the extrapolation are the aspects that are controversially discussed among Immunologists because the majority of patients with autoimmune diseases receive this medication in addition to other immunomodulatory treatments. Thus, the evaluation of the induction of ADAs and the switchability are particularly complex. On the basis of the data that are currently available for the approval authorities, however, the possibility of switching the approved biosimilars is assumed [38]. Pharmaceuticals that are very similar to an originator substance but that do not pass this regulatory process (e. g., in the EU controlled by the EMA, in the USA by the FDA), are called intended copies. Those drugs can be purchased for example in emerging countries but they are not approved in the EU.

Biobetters are substances that dispose of more favorable properties for therapy due to small selective changes leading to altered physico-chemical properties, changed binding capacities, or modified degradation or clearance. One example in this context are insulins that have a longer-lasting effects or pegylated antibodies that increase for example the bio-availability due to binding with polyethylene glycol (PEG) [40].

1.8 Small molecules and DNAzyme

Those pharmaceuticals have also been developed with a rational design based on mechanistic studies, but contrary to prevailing assumption, they are no biologics. Many of the small molecules often act as tyrosine kinase inhibitors and require certain genetic subty-

pes or expression of markers for successful therapy. In this way, they are of particular significance for precision medicine. However, because of their physico-chemical properties, their approach is completely different and so they are typical pharmaceuticals, generally with a size of up to 800 Daltons. They have been established especially in the field of clinical oncology. Activating EGFR mutations are found in 10–15% of the patients with lung cancer (e. g., non-smokers, adenocarcinomas) that are therefore suitable for treatment with gefitinib [41]. Inhibitors of janus kinases (so-called JAK inhibitors), which mediate the activity of cytokine receptors as cytoplasmic tyrosine kinase, can be applied as e. g., the small molecule tofacitinib in autoimmune diseases [42] and possibly also as supporting immune modulator in the allergen specific immune therapy [43]. As selective chloride channel enhancers, the substances ivacaftor and lumacaftor are a new and promising therapeutic option for patients with cystic fibrosis and a delta F508 mutation of the CFTR gene and lead to a reduction of pulmonary exacerbation [44, 45].

A topical effect induced by DNAzymes that are targeted for example effectively and specifically against the Th2 transcription factor GATA3 and thus inhibit all downstream immuno-pathological Th2 reactions [46] is also mechanistically elegant but according to the definition they are no biologics.

1.9 Costs

The global market for biologics in 2020 is estimated to amount to 350 billion Euro [47]. In 2016, in Germany biological drugs were prescribed for around 6.4 billion Euro within the statutory health-

care system, corresponding to 19% of the overall turnover of all drugs and 2.5% of all prescriptions [49]. Possible cost reductions by biosimilars are estimated to approx. 20–30% because the regulatory obstacles for approval and pharmacovigilance are very high. Currently, 28 biosimilars are approved for application in the EU. It should be noted that cost reductions by competition are possible in particular where several biosimilars are approved whereby the market shares of biosimilars do not necessarily correlate with the prizes [49].

This calculation however does not include the increasing incidence of prescriptions because the multitude of new molecules as well as the increasing number of indications and a corresponding market penetration can be observed. For the EU and the USA, IMS Health expects possible economies of 50–100 billion US\$ between 2016 and 2020 by application of biosimilars.

2. Rhinology: Epidemiology and pathophysiological concepts

Rhinology deals with diseases of the outer and inner nose, the paranasal sinuses as well as the frontobase and their surgical and conservative therapy. As outpost and door of the airways, the nose in this context is an organ that has esthetic, functional, and immunological tasks and works as sensory, filter, and reaction organ [50].

2.1 Physiology of the nasal mucosa and the integrated mucosal immune system

The nasal mucosa is a first-line of defense. The preservation of the physical integrity by the respiratory mucosa as active and passive barrier, which is at the same time the location of innate and adaptive immune response, the complex function of the mucociliary clearance, the permanent confrontation with particulate and soluble substances such as antigens, allergens, and pathogens result in the according nosology and pathology known to every ENT specialist [51]. Rational understanding of the inflammatory and protective mechanisms led to the definition of new therapeutic target structure and concepts (see review article [52]).

The respiratory epithelium that is found in the respiratory region of the nose and the paranasal sinuses as well as partially in the nasopharynx is a pseudostratified epithelium. The ciliary function contributes to transporting mucus, possible noxious substances, and particles that have been inhaled in direction of the ostia or the mouth, respectively, whereby pathologies of the ciliary apparatus fundamentally alter the physiology of the airways [53]. The cilia have multiple functions; it could be shown among others that they are chemosensory. At the same time, the intact respiratory epithelium is the basis for functionally intact defense [54, 55]. At transition points to enhanced mechanical use, e. g., in the pharynx, the epithelium merges into a stratified squamous epithelium. In this epithelium, goblet cells are found that produce mucins. These mucins play an elementary role in the unspecific mucosal immunity; they contribute to the barrier function, interact in an antimicrobial way with multiple antimicrobial peptides and defensins [56], develop a highly flexible and complex mucin-interactome [57], and act synergistically with inflammatory cytokines, e. g., IL-1 β [58]. Submucous glands produce nasal lining secretions, these consist of both, a transsudative (vasomotor) and a secretory layer. In the luminal secretory layer, substances with an unspecific antimicrobial activity are found, such as e. g.,

lysozyme, lactoferrin and defensins as well as adaptive mucosal antibodies of the IgA type (IgA1 and IgA2) [59] and the secretory component, which ensures the biostability of secretory mucosal IgA [60]. In addition, also IgM and IgG are found. IgA is induced by interaction with commensals and external influences and may additionally neutralize exotoxins as well as pathogens and transport sIgA commensals through the epithelium in luminal direction. Mucosal IgA has multiple functions of immune exclusion due to various affinities [61] and thus it contributes relevantly to the mucosal immunity of the respiratory mucosa and determines an active antigen-specific protection at this important external border of the upper airways, which is the most exposed external surface of the human body in relation to the covered surface.

Beside protective humoral mechanisms, also a synergistically acting cellular compartment is found in the upper airway mucosa. This involves tissue macrophages, eosinophilic, neutrophilic granulocytes, basophils, and mast cells, that act according to stimuli and pathology. Intraepithelial lymphocytes of the CD4 and CD8 type act in an antigen specific and adaptive manner. Together with submucous lymphocyte infiltrates of the lamina propria and special, functionally adapted NALT regions with adapted M cells and dome areas, $\gamma\delta$ T cells [62], innate lymphoid cells (ILC) [63] as well as NKT cells may induce tolerogenic, allergic, or cytotoxic immune responses [64, 65]. Furthermore, the pharyngeal tonsil, which usually should involute by school age, is part of the integrated mucosal system in the nasopharynx. Beside T cell infiltrates in e. g., allergic rhinitis or chronic rhinosinusitis with nasal polyposis, also areas of local IgE production by B cells [66] as well as dendritic cells [67, 68] exist. Together with macrophage-like dendritic CD11 + CD14 + CD68 cells, those professional, antigen presenting, partly immature CD1 + CD11 + cells form a dense network for regulation of adaptive immune responses [69] and express especially in atopics the highly affine IgE receptor Fc ϵ RI as well as costimulatory molecules such as CD80, CD86, and CD40 [70]. There is evidence, that apart from lymphoepithelial areas respiratory epithelium adapts functionally to existing immunological conditions. Hereby, it is modified by inflammatory stimuli such as e. g., IFN γ or IL-4 in its transcriptome and consecutively functional properties. This leads for example to a defective barrier function due to reduced expression of tight junction proteins [71]. However, the epithelium is acting as an immunointerface in the initiation and maintenance of e. g., allergic inflammation. This involves inflammatory cytokines such as TSLP [72], IL-25 [73], or IL-24 [74] and epithelial alarmins such as IL-33, that contribute mucosal inflammation [75]. In the full picture of chronic rhinosinusitis with nasal polyposis of the western subtype, predominantly a type 2 inflammation is found with eosinophilic infiltrates, local expression of IL-5, IL-13, and the presence of IgE [76–78] (see also chapter 2.4 in [52]).

2.2 United Airways

The functional and physiological connection of the upper and the lower airways is a daily experience in the treatment of ENT patients. The association of clinically severe courses of e. g., chronic sinusitis with chronic bronchitis and asthma or allergic rhinitis and asthma can be identified in daily practice as well as in the epidemiological perspective.

In 2002, a prospective longitudinal population cohort from Denmark convincingly revealed for the first time the correlation between

allergic rhinitis and allergic asthma in people suffering from allergy to pollen [79]. Previous trials, even performed in well-structured national cohorts [80] were able to show associations but they used non-adjusted datasets.

In 2008, Shaaban and colleagues [81] examined more than 6,460 patients in a European longitudinal cohort study in 14 European countries that patients with allergic rhinitis had a approx. four-fold risk to develop bronchial asthma within the observation period of nearly 9 years. The adjusted relative risk amounted to 3.53 (95% CI: 2.11–5.91) for allergic rhinitis and – that is also a relevant result – to 2.71 (95% CI: 1.64–4.46) for non-allergic rhinitis.

Even in cases of chronic rhinosinusitis, the association with asthma has been investigated epidemiologically [82] and mechanistically apart from the typical Vidal or Samter triad. Depending on the sample, the percentage of asthmatic patients in the patient cohort with CRS amounts to 25–70% [83, 84]. A research team from the UK analyzed a clinical cohort with 57 patients suffering from CRSwNP with regard to pulmonary function, bronchial hyperreactivity, and exhaled nitrogen monoxide (FeNO). In this context, 3 clinical groups with different phenomena could be identified. However, these groups were not different regarding the severity of their nasal symptoms [85]. Not only in cases of allergic asthma, but also in cases of non-allergic asthma and COPD, an association to pathologies of the nose and the paranasal sinuses could be demonstrated. Patients with non-allergic asthma and COPD had a higher symptom burden in the validated SNOT-20 questionnaire as well as increased concentrations of inflammatory cytokines in nasal lining secretions, e. g., IFN γ and G-CSF, but also eotaxin and MCP-1 [86].

The didactically and functionally interesting concept of the “atopic march” [87] fits well in such most likely T cell mediated disease concepts based on the assumption of antigen-specific inflammatory T cells migration.

In the early 2000ies, Braunstahl conducted segmental-bronchial and nasal allergen challenges and could show that a segmental bronchial allergen provocation leads to infiltration of IL-5 producing cells and of eosinophils in the nasal mucosa [88]. Nasal allergen provocation again leads to increased expression of epithelial (ICAM-1) and vascular (VCAM-1) adhesion molecules in the bronchial mucosa correlating with the number of locally expressed eosinophils [89]. Also infiltrates with basophils and mast cells could be revealed [90]. Despite the elegant study design, the major weakness of those studies is that the cellular source of the pro-eosinophilic IL-5 was neither functionally nor morphologically determined.

The interactions between the upper and lower airways are far from being only based on immunological mechanisms. The research team of Baroody and Naclerio could show that the nasal mucosa of patients suffering from allergic asthma can only poorly condition the air, i. e., warming it up and moistening it [91, 92].

2.3 Entities and epidemiology of diseases of the nasal mucosa and the paranasal sinuses

2.3.1 Infections

The most frequent origin of upper airway diseases is an acute viral infection with respiratory viruses; primarily bacterial rhinosinusitis is comparably rare. The etiology of chronic rhinitis and chronic rhinosinusitis is complex and it is mostly unclear if in the sense of “first hit hypothesis” at the beginning of the inflammatory cascade viral,

fungal, or bacterial infections must be assumed. Also the role of allergies, local or systemic, is not finally investigated.

The common viral cold, coryza, is the most frequently occurring infectious disease at all. Because of difficult distinction, Eccles considers it more as a cultural concept than a clearly defined clinical entity [93]. Meanwhile, 99 serotypes of human rhinoviruses have been phylogenetically examined [94] and completely sequenced [95]. The variability of those viruses is very high and thus rather difficult for the immune system: beside the variety of antigens and the specific virulence of the pathogens, also the pathogen-host interactions determine the pathogenicity and penetrance. One example is the genetic association of bronchial rhinovirus infections and childhood asthma [96].

In vitro, a Th2 microenvironment leads to an increased infection rate with human rhinoviruses under the influence of the cytokine IL-13 by upregulating the epithelial adhesion factor ICAM-1 [97]. In patient cohorts, this supposedly simple correlation remained unconfirmed up to now. However, it could be revealed that children with asthma have a clearly longer-lasting postviral hyperreagibility of the airways [98]. An afebrile rhinovirus infection may become an exacerbation of chronic rhinosinusitis or exacerbated asthma [99, 100]. Probably, rhinoviruses are responsible for about 50% of respiratory infections, beside influenza, adenoviruses, coronaviruses, and RSV (respiratory syncytial virus). In particular influenza viruses are accessible for vaccination strategies [101].

The early infection with RSV is also considered as risk factor for the development of allergic asthma [102, 103]. Developments of vaccination strategies for coronaviruses and RSV are relevant especially for high risk populations in the sense of stratified prevention. For prophylaxis of RSV pneumonia in susceptible children (e. g., premature children), the biologic pavilzumab is available as secondary preventive, passively immunizing antibody (Guideline in Pediatrics [104]).

The role of viral infections of the upper airways regarding genesis and exacerbation, is currently only insufficiently understood; in vitro models show in particular an induction of co-stimulating signals [105] in the epithelium and for example the synergistic impact in the Th2 microenvironment on pendlins, which contribute among others to dyscrinia and secretion in cases of chronic rhinosinusitis [106].

Acute bacterial superinfections of viral sinusitis occur in about 2% of uncomplicated cases [107–109]. The most frequent pathogens for simple acute bacterial rhinosinusitis without preexisting chronic rhinosinusitis are *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catharrhalis*, and *Staphylococcus aureus*. In the context of preexisting infection with rhinoviruses, an increased adhesion of different bacteria could be shown among others for *Staphylococcus aureus* [110].

Under the influence of IL-4, an altered function neutrophil granulocytes has been described in atopy [111], in that bacterial superinfection can be easily explained mechanistically in the model. The microbial interactions in chronic rhinosinusitis, however, are complex. On one hand, biofilms play a relevant role [112], on the other hand, the microbiome in the investigated patient population is probably iatrogenically altered and limited in its diversity in favor of single species. So Abreu et al. revealed a loss of lactic acid bacteria in favor of *Corynebacterium tuberculostearium* [113]. Choi and colleagues could identify colonization with *Staphylococcus aureus* [114]. Since especially severe cases of CRS with nasal polyposis are associated with the presence of specific IgE against *S. aureus* enterotoxin and

also the colonization of the mucosa with *S. aureus* is accompanied with nasal polyposis, tissue eosinophilia, and a Th2 microenvironment [115], the question of cause and effect has to be asked. Actually, *S. aureus* IgE is also found in severe adult asthma [116] and the involvement of superantigens in atopic eczema is documented clinically as well as experimentally [117]. Hence it is very difficult to estimate the role of a complex microbiome in these heterogenic cohorts that are often pretreated with antibiotics. Data on the exacerbation by bacterial superinfection in CRS are incomplete. Therapeutic microbiome interventions, e. g., as microbiota transfer may be promising tools in personalized medicine [118], however, this topic has not been published in rhinology up to now.

2.3.2 Tobacco smoke

In epidemiological studies, tobacco smoke could be consistently shown as independent risk factor for sinusitis (OR 1.7; 95% CI: 1.6–1.9) [119], an overview is found in Beule's contribution [120].

2.3.3 Allergies

Worldwide, between 10 and 40% of all people are affected by allergic rhinitis [121, 122]. Since many years, the epidemiology of allergic rhinitis is investigated on the basis of consistently structured international patient cohorts, mainly according to the ISAAC standard (International Study of Asthma and Allergy in Childhood [123]), that are further supported by population-based representative cohort studies. Unanimously, the picture of a globally increasing prevalence of respiratory allergies developed since the 1980ies that at least in Europe reached a high-level plateau in the last years. Approx. 500 million people probably suffer from allergic rhinitis worldwide; the socio-economic consequences are enormous (see overview in [122]). In Germany, the data of the study on adult health (DEGS1) reveal a lifetime prevalence for allergic diseases of about 30% [124]. In the context of population-related sampling with more than 7,000 blood tests of adults, at least 48% of the patients had allergic sensitization, hereby a total of 33.6% of the participants were sensitized against aeroallergens [125]. These percentages were confirmed by the Robert Koch Institut (RKI) in the general study on adult health in Germany from 2014 ("Gesundheit Erwachsener in Deutschland 2014" (GEDA study) enrolling 23,000 randomly selected individuals. The RKI reports about an "allergy tsunami": the 12-month prevalence for allergic diseases amounted to approx. 28% [126]. In the USA, also up to 30% of the population suffer from allergic rhinitis, about 60% of them cannot adequately control their symptoms by treatment with antihistamines or topical steroids [127]. In Germany, only about 10% of the patients are treated in compliance with the according guidelines [128]. The only causal therapeutic option for treatment of allergic rhinitis is specific immunotherapy [122, 129], which shows an excellently documented clinical effectiveness and safety as subcutaneous [130] as well as sublingual immunotherapy [131–133]. Furthermore, its effect is downstream preventive on the development of allergic bronchial asthma [134–136] (see also chapter 2.5).

2.3.4 Chronic rhinitis

Comparably reliable epidemiological data on chronic rhinitis in adolescents are found due to the Isle of Wight birth cohort that is also the association of different clusters on the risk to develop asthma

[137]. It is not easy to clinically and terminologically distinguish chronic rhinitis as differential diagnosis from chronic rhinosinusitis. Recent reports define the specific consensus, however, in contrast to allergic rhinitis sound data are not available [138, 139]. In the context of non-allergic chronic rhinitis, several subtypes are defined such as non-allergic rhinitis with eosinophilia (NARES) [140], hormonal rhinopathy, rhinitis with neurogenic inflammation [141] (which is possibly a specific endotype, see chapter 2.4), and idiopathic rhinitis (which is still called vasomotor rhinitis in English-speaking countries). Those subtypes can be treated selectively with topical nasal steroids, e. g., ipratropium bromide (off-label use in Germany) or capsaicin (if available).

2.3.5 Chronic rhinosinusitis

The category of chronic rhinosinusitis defines inflammatory diseases of the nose and the paranasal sinuses with durations of more than 12 weeks. The EPOS2 guideline from 2012 [107] defines chronic rhinosinusitis as a disease of the nose and the paranasal sinuses of at least 12 weeks duration characterized by more than 2 of the following symptoms: a) nasal obstruction, b) rhinorrhea, c) pressure sensation or facial pains, d) hyposmia or anosmia, while either at least nasal obstruction or rhinorrhea should be present as obligatory symptoms. Clinically, either endoscopic signs of nasal polyposis, mucopurulent discharge from the middle nasal meatus and/or CT morphological signs of mucosal alterations in the osteomeatal unit or the paranasal sinuses should be found. This definition taken from the European position paper has been increasingly accepted in the last years. Furthermore, independent from the etiology and pathogenesis of the disease, chronic rhinosinusitis was classified into chronic rhinosinusitis with nasal polyposis (CRSwNP) and chronic rhinosinusitis without (lat.: sine) nasal polyposis (CRSsNP) [107, 142].

It is difficult to perform methodically and regionally consistent evaluations of the epidemiology of chronic rhinosinusitis. On one hand, significant regional differences are found regarding the identical assessment methods, on the other hand, there are various approaches for assessment, for example in the context of medical diagnosis and – which is meanwhile generally established in epidemiological methods – questionnaires and/or the assessment of the disease-related quality of life. A comprehensive overview can be found in [120]. The most consistent assessment by means of standardized questionnaires in the context of the "Global Allergy and Asthma Network of Excellence" (GA2LEN) was uniquely performed for chronic rhinosinusitis [119] and in a combined way for asthma associated with chronic rhinosinusitis [82].

Hereby, the first trial with more than 57,000 returned questionnaires from 12 different European countries with patients between the ages of 15 and 75 years could reveal a global prevalence of 10.9% (interval: 6.9–27.1%) according to the definition of European guidelines. Those data were confirmed among others by an American study with a global prevalence of CRS of 11.9% [143]. The second trial [82] additionally investigated the association with asthma in this dataset. Despite the heterogeneity in the prevalence of CRS between the single regions in Europe, the association between CRS and asthma was comparable over all centers and age groups (adjusted OR: 3.47; 95% CI: 3.20–3.76). Patients who had allergic rhinitis beside the symptoms of CRS had an additionally increased risk to be affected from asthma (adjusted OR: 11.85; 95% CI: 10.57–13.17). In non-

allergic individuals, however, an association to so-called late-onset asthma was found. CRS is accompanied by a clearly reduced quality of life and has a high socio-economic relevance [144].

AERD (aspirin exacerbated respiratory disease), is associated with rather severe course of disease [107]. A recent meta-analysis [145] reported about a 7–15 % prevalence of AERD in asthmatic individuals and an increased prevalence in severe cases. In about 8.7 % of the cases, patients with CRS showed symptoms of AERD, patients with confirmed CRSwNP had those symptoms in approx. 9.7 %. According to EPOS3 [107], about 15 % of the patients with CRS had AERD. Zhang et al. could demonstrate that specific IgE against *S. aureus* enterotoxin is expressed disproportionately frequently in those patients [146].

Important, although less frequent differential diagnoses are autoimmune system diseases with involvement of the nasal mucosa and the paranasal sinuses. In this context, in particular vasculitis has to be pointed out. Depending on the stage, its clinical image may appear similar to chronic rhinosinusitis [147, 148]. Recently, the American College of Rheumatology (ACR) performed changes of the definitions, in particular GPA (formerly called Wegener's granulomatosis [149]) and eosinophilic granulomatosis with polyangiitis (formerly called Churg-Strauss vasculitis) must be mentioned. The diagnostic criteria of GPA of the European League against Rheumatism (EULAR) and the ACR are currently being revised based on the current "Diagnostic and Classification Criteria in Vasculitis Study" (DCVAS) [150]. Regarding the treatment of GPA with rituximab, meanwhile a recommendation degree A has been stated [151]. From a mechanistic point of view, also eosinophilic GPA (EGPA) is an interesting target for antibodies that target IL-5 or its receptor as recently investigated in clinical studies [152] (see chapter 4.2 below).

Cystic fibrosis (also: mucoviscidosis) is the most frequent autosomal-recessively inherited genetic disease (1:2,000 live births) and is associated with a dysfunction of membraneous chloride channels (CFTR, cystic fibrosis transmembrane regulator) that leads to modified mucus composition in the airways and the gastrointestinal tract. In the airways, the viscous mucus that can only hardly be mobilized leads to recurrent, opportunistic, and often life-threatening broncho-pulmonary and upper-airway infections. In cases of missing enzyme substitution, the altered function in the gastrointestinal tract results in malnutrition syndromes due to exocrine pancreatic insufficiency. Nearly 2,000 mutations of the CFTR gene are known, so the phenotypes can vary considerably (an overview can be found in [44, 45]). In at least 25–40 % of all patients, sinonasal polyposis is found [107, 153]. Phenotypically, histological subtypes are observed depending on the sample, with rather neutrophilic or eosinophilic inflammation patterns [154, 155]. Regarding the treatment of CF, new selective therapies with small molecules could be established (see above, chapter 1.8).

More rarely, the Kartagener syndrome [156] is found, which is the prototype of ciliary dyskinesia [53] that is obligatorily associated with nasal polyps beside bronchiectasis and situs inversus. The polyps may also contain eosinophilic infiltrates, interestingly also a clearly reduced expression of NO synthetase in the tissue is found [157]. These findings correspond a priori to the low exhaled NO values in the diagnosis [158]. In exhalations, the use of an eNose may differentiate between healthy patients, Kartagener syndrome, and cystic fibrosis with or without chronic infection with *Pseudomonas aeruginosa* [159]. In this way, it is an interesting new tool for potential endotyping.

2.4 Endotypes of chronic airway inflammations

The phenotypic definition of chronic sinusitis with or without nasal polyposis does not give any hint to the molecular developmental mechanism, it cannot predict the therapeutic outcome, and does not contribute to the allocation of optimal therapy, i. e., for example conservative standard therapy with or without surgery [142, 160]. A sound mechanistic understanding, however, is the basis for a targeted, ideally precise and personalized therapy. Hypothesis-driven research in immunology and inflammation of the past 30 years has identified the association of allergic airway diseases with phenotypically predominant eosinophilia to the Th2 diseases mainly on the basis of the also didactically elegant Th1/Th2 model [9, 10]. In nasal polyps this was first shown by Bachert [76], further intensified investigations revealed cytokine patterns [77] and also the involved T cell clones [161] followed. However, for example in Asian cohorts, distinct IL-17 associated inflammatory pattern were [162]. Interestingly, mucosal inflammatory patterns have revealed a transition to a more "western" cytokine profile over time, probably associated with socio-cultural transition [163]. Furthermore, patients were diagnosed with CRSwNP in central China in whom none of the expected or "classic" cytokine microenvironments was dominant, i. e., neither Th1 nor Th2 nor Th17 inflammation [164]. Even in special types of CRS, e. g., cystic fibrosis, other inflammatory signatures are found [161, 165]. The presence of different infiltrates correlates with the response to the therapy. So Wen [166] could show that nasal polyposis with predominantly neutrophilic infiltrates does not clinically respond to oral steroid treatment. Mucin-1 expression, however, could be identified as marker for a response to steroids [167].

The Th2-associated pathophysiology in inflammatory airway diseases in the western world led accordingly to biomedical therapy concepts which address mainly cytokines from the Th2 microenvironment (see chapter 4.2, clinical studies). Apart from the question if those new therapy procedures are affordable, a concept has been developed in the context of personalized medicine that describes endotypes. First described by Anderson [168], this concept was intended to result in a more selective and mechanistically rational therapeutic procedure. Regarding the definition of endotypes, for example genetic, mechanistic, histological, or functional properties are classified in order to define different and mechanistically coherent entities [169]. In a simplified way, Wenzel described endotypes as a "molecular phenotype" [170]. In the last years several position papers of highly reputed authors discussed this concept: at least – and this is a great progress – a European multicenter analysis and merely data-related study is now available for chronic rhinosinusitis [171].

For this cohort in the GA2LEN network, initially 917 patients from 8 European countries (respectively 10 European University Hospitals) were enrolled. Finally tissue could be gained with adequate quality and quantity from 173 patients with CRS and 89 controls. Based on a pre-defined biomarker selection, all specimens were characterized and assigned to 10 different groups by hypothesis-free analysis. Inflammatory parameters (MPO, IL-1 β , IL-6, and IL-8), type 2 inflammation markers (IgE, ECP, IL-5, and albumin), in the 3rd group IL-17, TNF- α , and IL-22, and additionally in 2 further groups IFN γ , TGF β ₁, *S. aureus* enterotoxin IgE were examined in the context of a hierarchic cluster analysis. The resulting 10 clusters were compared to clinical phenotypes and appeared mechanistically and clinically

plausible in order to depict coherent subtypes. IL-5 negative groups clinically showed mainly chronic rhinosinusitis without nasal polyposis and without asthma whereas the IL-5 positive groups, which also had a high systemic and local IgE level, contained a high percentage of asthma-affected patients. Patients who had particularly high IgE values as well as specific IgE against *S. aureus* enterotoxin generally suffered from chronic rhinosinusitis with nasal polyposis and nearly all of them from comorbid bronchial asthma. However, unfortunately this trial has one relevant weakness: Even if further clinical data have been assessed, they are not included, i. e., data on standardized symptom scores and/or disease-related quality of life. Ideally, also more than the initially 900 patients could and should have been included in the analysis, however, cautiously 3 axiomatic statements can be made that would have to be verified in real-life clinical data and pivotal clinical trials:

1. The more severe chronic rhinosinusitis affects a patient in Europe, the more mechanistically probable is an eosinophilic inflammation that is associated with a quantitative involvement of IL-5, IgE and IgE against *S. aureus* enterotoxin, depending on the severity.
2. Different inflammatory patterns can exist in parallel: in addition to Th2 cytokines and IL-5, generic-inflammatory activation of IL-6 and IL-8 is found in single clusters, furthermore IFN γ and depending on this IL-17 with IL-22, or IL-22 alone.
3. Functional endoscopic sinus surgery can only be one element among others for treatment of this complex inflammatory disease involving the upper and lower airways and should ideally be accompanied by rational anti-inflammatory measures.

Beside those Th2-associated endotypes, probably neurogenic-inflammatory endotypes exist in the phenotype of non-allergic inflammation that responds well to capsaicin therapy [141]. The terminological inexactness becomes apparent for example in the context of hormonally induced pregnancy rhinopathy: there is a very distinct hormonal pathomechanism that is sufficient for endotype definition, however, the disease can be defined only phenotypically in the patient.

The elaboration of endotypes of the upper airways is developing and there are many aspects where no adequate terminological consensus has been found yet [172, 173]. Even more significant is the data deficit which has to be met in the near future.

2.5 Stratified downstream prevention

The early involvement of the nose in cases of respiratory diseases makes the upper airways a potential proxy for the lower airways and an interesting tertiary-preventive lever.

Allergen-specific immunotherapy (AIT) as classic personalized therapy is the working horse in clinical allergy. In contrast to the ongoing administration of therapeutic antibodies that may be considered as disease response modifiers, there is compelling data on the disease modifying effects of AIT and based on these properties can be utilized as an instrument in downstream prevention.

By means of the randomized, controlled and open Preventive Allergy Treatment Study (PAT Study) with 183 children it could be shown for allergic rhinitis that the children who had received a 3-year specific immunotherapy (SIT) against birch and/or grass pollen suffered significantly less from allergic asthma after 5 years of therapy

onset (OR: 2.68: 1.3–5.7) [135]. Ten years after therapy onset, even more favorable outcomes were found in the immunotherapy group with an odds ratio of 4.6; 95 % CI: 1.5–13.7 not to develop asthma [134].

These data could be confirmed by analyses of retrospectively evaluated healthcare data from East Germany that included datasets of 118,754 patients between 2006 and 2012. In allergic patients who had received AIT, a lower number of newly diagnosed asthma cases was observed in comparison to allergic individuals who had not received AIT. The relative risk after regression analysis amounted to 60 % (RR, 0.60; 95 % CI: 0.42–0.84), but the non-adjusted risk did not vary [174].

With publication of the Grass-Asthma-Prevention Study (GAP Study), for the first time a double-blind, placebo-controlled trial revealed a downstream preventive effect with regard to asthma symptoms for the specific allergen immunotherapy. In this European trial, 812 children were randomized with a high-dose grass pollen tablet therapy for 3 years. The study did not reach its primary endpoint, i. e., time to onset of asthma resp. wheeze based on pre-defined criteria that included reversibility of obstruction after beta-2-agonist administration. However, consistently and significantly less asthma symptoms and less consumption of asthma medication could be shown in the verum group (OR: 0.66, $p < 0.036$) [136].

In contrast to the specific immunotherapy for allergic rhinitis, the data on chronic rhinosinusitis have to be evaluated in a much more heterogenic way and the cohorts for interventions are generally smaller. In this context, in particular healthcare studies are interesting. Retrospective data of the National Health Service (NHS) in the United Kingdom show that late surgery of patients with chronic rhinosinusitis may often lead to severe and recurrent cases as well as to bronchial asthma. Market research data seem to support these statements [175, 176]. A study encompassing prospective data of a national British audit reports that especially patients who suffered from allergies and asthma underwent surgery particularly late. This may be due to the slow process of symptom adaptation as well as to possibly conservative therapy attitudes. However, the data reveal that patients who underwent surgery at early disease stages have a better symptom control and disease-related quality of life in the context of follow-up examinations after 12 and 60 months [177]. Furthermore, patients who had undergone surgery at an earlier time cause less subsequent healthcare expenses [178].

Whether this conclusion proves correct and early surgery really protects patients with chronic rhinosinusitis against recurrent disease courses and against comorbid asthma can only be demonstrated by means of consistent data from randomized interventional studies, cohort, and healthcare data. Only such data allow suggesting treatment recommendations for stratified prevention approaches in the sense of personalized medicine.

3. State-of-the-art

The existing S2k guideline for the treatment of Rhinosinusitis for German-speaking countries was revised in 2017 [179]. The European position paper EPOS that had been revised in 2012 in its 3rd edition basically corresponds to a European guideline regarding the dimension and the scientific claim and encompasses acute as well as chronic rhinosinusitis. For allergic rhinitis, the extensive document entitled "Allergic rhinitis and its impact on asthma" is referenced as

standard manual in the area of the European Academy of Allergy and Clinical Immunology (EAACI). An important milestone for causal therapy of allergic rhinitis is the German guideline on specific immunotherapy published in 2014 [129].

The significance of functional endoscopic sinus surgery for therapy of CRS is estimated as very high. This is supported on one hand by the data of healthcare systems with limited access to high-quality surgical care and on the other hand by merely rational mechanistic reflections of the pathological anatomy of the osteomeatal unit [180, 181]. Hereby, surgical therapy should be performed in all cases where conservative therapy only allows insufficient control of the CRS symptoms and when endoscopy and/or imaging defines a confirmed pathological correlate as curative surgical objective. Despite 2 recently published meta-analyses, the currently available evidence is unsatisfactory [182, 183] because the meta-analyses can only evaluate high-quality standardized studies, which are currently not available.

Based on the existing evidences (level 1a) for 2 main pillars of conservative therapy of chronic rhinosinusitis without nasal polyposis, the European guideline EPOS3 recommends (recommendation grade A) to apply topical nasal steroids and nasal rinsing with saline solution. For patients suffering from chronic rhinosinusitis with nasal polyposis, topical nasal and oral steroids are recommended as conservative treatment, in EPOS3 data from meta-analyses are provided that support this recommendation [107].

A clear recommendation in favor of topical therapy is found in an evidence-based review published by Rudmik [184]. Furthermore, 2 recent meta-analyses of the Cochrane Society are available regarding topical steroids after a meta-analysis had been withdrawn because of the availability of new data published by Kalish [185]. In 2016, Chong presented a meta-analysis that could include 18 randomized, controlled trials with a pool of 2,738 patients. The authors of the meta-analysis criticize that in the context of trials on topical nasal steroids a moderate effect on symptoms such as nasal obstruction and the overall severity of the disease could be measured in pooled datasets but the overall quality of the evidence, in relation to the number of studies and patients, was only moderate or even poor and the follow-up periods were too short. In addition, the disease-associated quality of life had not been sufficiently taken into account [186]. In a second meta-analysis, nearly the same group of authors compared different intranasal topical steroids. Based on the available trials, however, no recommendation could be given for single substances or application modalities [187].

In the European position paper, a large variety of therapies achieves the recommendation levels of C or D (not recommended). Based on the recent meta-analyses and also the confirmatory trials with biologics, the next revision will most probably implement new recommendations.

Without any doubt, the currently guideline-based therapy treats many patients in an adequate way, especially in such a well-structured and accessible healthcare system as it is found in Germany. However, data from phenotyping studies show that the therapy responses are very heterogenic regarding surgical as well as conservative therapy approaches and that some patients do not benefit at all [160]. Based on an algorithm with non-supervised machine learning, Soler and colleagues tried to predictively identify these groups. This was partially successful in a pilot study, but it led to clinical groups that were not attributable by known and intuitively understood features and that ad-

ditionally completely lack of an – at least currently known – distinct pathophysiological correlate. Nonetheless, such cohorts have to be evaluated and reproduced in order to prospectively verify a possible usability also in algorithms. In summary of endoscopic and radiological findings with systematic eosinophilia in a cohort of more than 1,700 Japanese patients, a scoring system entitled JESREC (Japanese Epidemiological Survey of Refractory Eosinophilic Chronic Rhinosinusitis) provided a cut-off value that allows describing a diagnostic criterion for eosinophilic CRSwNP and an increased risk for recurrences [188]. These results should be reproduced on an international level with a modified score, if needed. One weakness of the trial is that it could not be verified to what extent the pretreatment with steroids has an effect on the applicability of these tests because patients who had been pretreated with steroids were excluded from the study.

A so-called “unmet clinical need” of the current guideline-based therapy results because of different reasons. On one hand, it has become clear that assumedly more than 20% of the patients can only partially or not at all control their symptoms despite guideline-based, adequate, effective, and safe therapy [189]. In addition, the percentage of recurrences seems to have been underestimated for years. In the United Kingdom, the rate of surgical revisions amounts to 19.1% for CRS and 20.6% for CRSwNP in a 5-years interval and has not improved despite optimized surgical techniques and the availability of further-developed topical steroids [190]. In single cohorts, up to 80% of the patients with CRSwNP develop recurrences [84].

It is highly probable that the patients will have to be informed more realistically; however, better characterized clinical cohorts will be needed. Further, it is completely unclear to which percentage for example missing cooperation and suboptimal compliance regarding the topical steroid intake amount in case of severe courses [191]. The lack of disease control in these epidemiologically and socio-economically relevant cohorts of chronic patients with severe or persisting allergic rhinitis and/or chronic rhinosinusitis is a factor that is currently not adequately quantified. A rather relevant percentage of those patients will probably have to be classified as SCUAD (severe chronic upper airway disease) with multifactorial underlying disease processes [192].

In this context, stratified medicine may allow individual therapy and downstream prevention. Possible measures are pathophysiologically rational and targeted interventions with biologics.

4. Biologicals in clinical trials with rhinological diseases

4.1 Allergic rhinitis

4.1.1 Trials with omalizumab

The until now last and fifth human immunoglobulin class IgE was discovered in 1967 after a scientific race competition of Johansson and Bennich against Kimishige and Teruko Ishizaka. Ishizaka called it immunoglobulin E [193], but was not able to isolate and characterize it. Reagin of the Prausnitz-Küster reaction that had not been identified up to that time was characterized by Bennich and Johansson [194] and the association to asthma was revealed in the same year [195]. The IgE mediated sensitization against an allergen is the pathomechanistic basis of modern allergology and completes the concept created by Pirquet that allergy is a specific hypersensitivity reaction [196].

Omalizumab is a humanized monoclonal antibody against IgE and was approved by the FDA in 2003 and by the European Medical Agency (former EMEA) in 2005 for therapy of severe allergic asthma. Mechanistic studies on causal treatment of allergic disease did not meet the expectations first because further allergic reactions occurred despite the effective pharmacological antagonism and elimination of IgE from the serum.

In a double-blind and placebo-controlled trial with a cohort of 536 patients, Casale could demonstrate the dose-related effect of omalizumab as monotherapy for patients with allergic rhinitis against ragweed [197]. In 2003, this effect could be confirmed for perennial allergens [198].

Mechanistically interesting was the combination of omalizumab with allergen-specific immunotherapy. Hereby, omalizumab could clearly reduce the incidence and severity of the side effects in the phase of increased dosing in particular in the context of accelerated dose increase of the allergen for effective and tolerogenic maintenance dosage, so-called rush-immunotherapy. Kopp applied omalizumab in a pediatric cohort and could observe *in vitro* a reduced release of leukotrienes in children who received immunotherapy against pollen under the protection of omalizumab [199].

In the context of a trial of the Immune Tolerance Network of the National Institute of Health (USA), Casale conducted a double-blind, placebo-controlled 4-arm study with a rush immunotherapy against ragweed pollen in combination with and without omalizumab as well as a double placebo group. In each arm 39–40 patients were randomized. From a total of 159 patients, 123 completed the trial that started preseasonally, included the pollen season, and lasted for 21 weeks. The primary endpoint of the study was the comparison of the seasonal symptom score between the group with combined application of omalizumab and allergen immunotherapy and the group with immunotherapy alone. The primary endpoint could be achieved when the effect size was relatively low (0.61 vs. 0.85, $p=0.12$). Furthermore, a post-hoc analysis could show that the application of omalizumab significantly reduced the incidence of systemic allergic reactions by 80% during rush immunotherapy; however, with 25.6% the rate was extremely high in the group of immunotherapy without omalizumab and was based on a blinded self-assessments of the patients. The increased effectiveness of the combined application of omalizumab with immunotherapy was explained by the fact that omalizumab was further therapeutically administered at the time of pollen occurrence. If the effect of immunotherapy was better in the group treated with the combined therapy beyond the time of systemic elimination of omalizumab, cannot be stated based on the published data.

In vitro specimens of a total of $n=36$ individuals from the same patient cohort, were investigated in further mechanistic studies by Klunker et al. Hereby, a validated facilitated antigen binding (FAB) assay was measured for assessment of the serum inhibitory activity against IgE binding to B cells. In the group of patients who underwent combined therapy and in the group with omalizumab, binding of IgE was inhibited of nearly 100%, even beyond the pollen season. The group with immunotherapy alone reached an inhibition of 50%. Mechanistically, it could be shown that allergen-specific IgE was no longer available in both therapy arms with omalizumab. Interestingly, the serum-inhibitory effect could be maintained up to 42 weeks after treatment and was even more pronounced in the group with combined treatment [200].

In clinical routine, the combined application of omalizumab plus allergen could be established in Europe, in particular for patients who require an increased safety profile regarding specific immunotherapy. This concerns for example the presence of severe bronchial asthma, urticaria, severe food allergies [202], or pre-existing anaphylaxis. Hereby it must be taken into consideration that not all these applications are in-label. A sound review article is found under [203].

4.1.2 Trials with VAK694

The induction of regulatory T cells that express the transcription factor Foxp3 and other subtypes, e. g., Tr1 cells, is considered a mechanistic key event of specific immunotherapy [204]. Actually, monitoring of this T cell conversion to regulatory populations is extremely difficult and in humans it is only possibly by extensive *ex vivo* examinations [205, 206]. Antigen-specific monitoring is only performed via flow cytometric analyses with tetramers or ELISpots [207, 208]. Mechanistic studies on specific immunotherapy could show that the induction of regulatory T cells is associated with the secretion of inhibitory cytokines such as TGF- β_1 and IL-10; IL-4, IL-5, and IL-13 secreting T cells are reduced [209–212]. While IL-5 mainly contributes to eosinophilia in allergic inflammation as pro-eosinophilic cytokine, IL-4 and IL-13 induce the production of IgE in B cells and have direct proinflammatory effects on the respiratory epithelium [74], mucous secretion as well as goblet cell hyperplasia of allergic inflammation of the airways [213]. On this basis, the concept of combined specific immunotherapy with human anti-IL-4 antibodies was developed. The antibody VAK694 is a fully human antibody directed against the cytokine IL-4 and was applied in patients in the context of an exploratory, double-blind, placebo-controlled 3-arm study. Primary endpoint of the trial was the cutaneous allergen-specific late-stage reaction as antigen-specific surrogate for T cell suppression 12 months after therapy *in vivo*. Explorative surrogate endpoint was the antigen-specific production of IL-4 in ELISpots *in vitro*. Furthermore, the T cell populations were characterized by flow cytometry. The study was designed as proof-of-concept trial and not sufficiently powered for detection of symptom differences between the therapy groups in the pollen season. The primary endpoint, tolerance in the cutaneous late phase response compared to standard therapy, i. e., specific immunotherapy alone, was not achieved because standard therapy alone reached a suppression of the allergen-induced late phase response in the skin of more than 90% although a subeffective dosage of the allergen had been chosen. The proof of concept was achieved *in vitro*: a sustainable suppression of allergen-specific IL-4 producing cells could be shown 12 months after the end of combined therapy and in comparison to immunotherapy alone as well as placebo. This effect, however, did not translate into the clinical endpoint, the cutaneous late phase reaction [215]. If the combination of anti-IL-4 with specific immunotherapy is pursued, remains unclear because the development of VAK694 was stopped.


4.2 Chronic rhinosinusitis

4.2.1 Omalizumab

The role of IgE in asthma, allergic rhinitis, and atopic dermatitis as well as parasite defense was described already very early [216]. In 1970, IgE was found for the first time in tissue homogenates in nasal polyposis [217], and Whiteside examined it in 1975 [218] regarding local lymphocytes and the correlation with systemic IgE levels and

atopy. Already at that time, the question of local IgE production in nasal polyps was addressed which became a relevant research issue in the pathophysiology of CRS. In 2001, Bachert could show that the proportion of IgE in the tissue correlates with local eosinophilic infiltrates [78]. In the early 2000ies, the association with specific IgE against *S. aureus* enterotoxin [146, 219] was discovered, in particular in severe cases, which was another hint to pathophysiologic involvement of IgE. In 2005, Gevaert could document the local IgE synthesis in the tissue [66]. From here, it was a logic consequence to investigate the therapeutic application of omalizumab in patients with CRSwNP in clinical trials. First case reports and retrospective case series were published by Penn [220], Guglielmo [221], and Vennera [222], mainly in patients who were treated in-label with omalizumab because of their severe asthma. Pinto and colleagues were the first research group that published a double-blind, placebo-controlled trial. The study could reveal no superiority of the verum arm compared to placebo. This was also due to the inclusion criteria because in this study patients with as well as without nasal polyposis were included [223]. A placebo-controlled follow-up trial published by Gevaert in 2013 with application of omalizumab in patients with CRSwNP and comorbid asthma over a treatment period of 16 weeks, could consistently observe a reduction of the nasal polyp score (-2.67 ; $P=0.001$) as well as the nasal symptom score and the disease-related quality of life in comparison to placebo [224] (► **Fig. 3**). Further studies in the context of this indication have already been completed according to the study registries (NCT01066104), at the time of manuscript submission the results have not been published. Subsequent studies on CRS have been registered as actively recruiting, among others under clinicaltrials.gov.

Paradoxically, in the initial phase of the treatment with omalizumab, a transient increase of the overall IgE is observed because of the development of biologically inactive but measurable IgE antibody complexes. Only after about 16 weeks, the pharmacodynamic response can be measured adequately in the serum. Pharmacodynamic investigations under omalizumab treatment additionally show that the de novo IgE synthesis decreases of about 50 % per year with the treatment with omalizumab [225]. This observation can be explained by a possible change of the IgE homeostasis because of ne-



► **Fig. 3** Change of the nasal polyp score under the treatment with omalizumab vs. placebo (according to [224], Gevaert P., JACI 2013, licensed by RightsLink/Elsevier).

gative feedback mechanisms that involve the low-affine IgE receptor. Thus it is justified to assume a possible timely limitation of therapy and withdrawal trials. For this purpose, long-term data would have to be evaluated in order to define rational therapy corridors.

4.2. 2 Reslizumab


The pathophysiological role of IL-5 in CRSwNP of western type was described early [76, 77]. A therapeutic antagonism seemed to be obvious. The first trial in humans and actual milestone trial of CRSwNP was published by Gevaert in 2006. In this first double-blind, placebo-controlled trial, a total of 24 patients with bilateral CRSwNP received a single dose of the humanized anti-IL-5 antibody reslizumab or placebo. The effect on systemic eosinophils in the blood and ECP concentrations in the serum could be pharmacodynamically measured up to 8 weeks. Only half of the patients treated with the real substance showed a therapeutic effect on the endoscopically revealed polyp scores. In a post-hoc analysis, Gevaert and colleagues could demonstrate that the responders were those patients who had high IL-5 concentrations in nasal secretion at the time of treatment onset. By means of a regression analysis, a cut-off value for IL-5 > 40 pg/ml in nasal lining secretions could be defined that could predict a positive therapy outcome after reslizumab application (odds ratio: 21.0; 95 % CI: 1.5-293.3; $P=0.009$). A multicenter study with reslizumab conducted in the USA investigated the effectiveness of poorly controlled eosinophilic bronchial asthma. Interestingly, reslizumab was more effective in the subgroup of asthma patients with known CRSwNP compared to asthma patients without polyposis [226]. This subtyping corresponded quasi to an indirect endotyping. Another trial on CRS with reslizumab has been registered as recruiting at the time of manuscript submission under clinicaltrials.gov.

4.2.3 Mepolizumab

After first successful studies on the treatment of eosinophilic asthma [227, 228], Gevaert conducted a 2:1 randomized pilot study of 8 weeks duration with 2 therapeutic applications (750 mg i. v., each) of the humanized anti-IL-5 antibody mepolizumab. Compared to the dosage that is currently approved in the EU for eosinophilic asthma (100 mg s.c.), this is a rather high dose that was however well tolerated and showed a reduction of the polyp size in the majority of the patients, which could be consistently documented by radiological assessment. Interestingly, in contrast to the reslizumab investigation, local IL-5 concentration in nasal lining secretions did not predict the therapeutic outcome in this trial [229].

In 2017, Bachert and colleagues published a European multicenter confirmatory phase-II/III-study on mepolizumab for therapy of CRSwNP. In this study that had been initiated in 2009, 105 patients from all over Europe with severe therapy-refractory CRSwNP with indication for surgical therapy were randomized 1:1 either for 750 mg mepolizumab or placebo and received their treatment 6 times in 4 weeks as add-on therapy to topical nasal steroids. The primary endpoint was defined as indication or need of surgical therapy of CRSwNP 25 weeks after therapy initiation. 25 weeks after randomization, 30 % of the patients with mepolizumab did no longer need surgical therapy (ITT, 16 [30 %] vs. 5 [10 %]; $P=0.006$). Consistently, also the VAS scores reduced in the verum group (-1.8 in week 25; ITT 95 % CI: 22.9-20.8; $P=0.001$) as well as the SNOT-22 test and in the post-hoc analysis even the endoscopic polyp scores (► **Fig. 4**). Also

in this trial, mepolizumab was well tolerated despite the relatively high dosage. In the verum group, smelling based on VAS improved, unfortunately, different smell tests were applied in the trial so that this important parameter cannot be systematically included in the analysis [230].



► **Fig. 4** Effectiveness of mepolizumab vs. placebo in CRSwNP: consistent reduction of the need for surgery, VAS scores and the endoscopic polyp score (according to [230]; Bachert C., JACI 2017, licensed by RightsLink/Elsevier).

4.2.4 Benralizumab

Similar to mepolizumab and reslizumab, benralizumab functionally inactivates the mediation of biological, mainly pro-eosinophilic effects of IL-5 by binding the therapeutic humanized antibody to the IL-5-alpha subunit of the IL-5 receptor (overview in [231]). For CRS, no trials are currently available. At the time of manuscript submission, one study on CRS was registered as active and not recruiting under clinicaltrials.gov.

4.2.5 Dupilumab


The fully human antibody binds the common alpha-subunit of the IL-4 and IL-13 receptor and thus interrupts the pleiotropic Th2 signals into immunological compartments. The high effectiveness in cases of eosinophilic asthma [232] and in particular in the treatment of atopic eczema was convincingly described with pivotal trials [233, 234]. Regarding the indication of CRS, an international consortium around Bachert and colleagues investigated dupilumab vs. placebo as add-on therapy of topical nasal steroids in 60 patients in a phase-II/III-study over 16 weeks. In the endoscopic polyp score, a difference of -1.6 of verum (95% CI: -2.4 to -0.7); $P < 0.001$) could be shown in comparison to placebo. This effect was consistent to a reduction of the CT-morphologically determined Lund-MacKay score and the SNOT-22 (► **Fig. 5**).

In 2017, dupilumab was approved for treatment of atopic eczema in the USA by the FDA, the approval for the EU was recommended by the EMA.

Other target structures for therapeutic antibodies are currently investigated. In this context, for example anti-TSLP and anti-Siglec-8 should be mentioned. For further possibilities of cytokine modulation, furthermore a test substance PF-06817024 in this indication is evaluated (source: clinicaltrials.gov).

4.3 Examples of rare indications

For patients suffering from eosinophilic GPA (Churg-Strauss vasculitis), which is often associated with an involvement of the upper airways in the sense of CRSwNP, Wechsler evaluated the application of



► **Fig. 5** Change of the nasal polyp score under the treatment with dupilumab vs. placebo (according to [232]; Bachert C., JAMA 2016, licensed by RightsLink/American Medical Association).

mepolizumab in a multicenter trial. In this study, 136 patients were randomized. The percentage of patients with remission was significantly higher in the group treated with mepolizumab (32 vs. 3 %; odds ratio: 16.74; 95 % CI: 3.61–77.56; $P < 0.001$), in the verum group, the consumption of systemic steroids could be clearly reduced in more than 40 % of the patients. The recurrence of sinonasal symptoms was complained by 35 % of the patients of the mepolizumab group vs. 51 % of the placebo group within one year. The profile of side effects and the steroid-saving effects allow the statement that mepolizumab is an interesting therapeutic alternative [152].

For cystic fibrosis, new and successfully tested substances were approved with chloride channel enhancers (see chapter 1.8).

4.4 Biomarkers

A true personalization of the above-mentioned and very expensive therapy by means of biomarkers could meet the requirements of precision medicine to make available the adequate therapy at the right time of the disease for every patient in the best way possible. However, in particular the implementation of these markers turns out to be much more difficult. This is also due to the missing protection by patents and the resulting only low commercial interest. Apart from comprehensive clinical work-up of patients with CRS (see also the revised German guideline [179]), simple markers are a good working basis.

4.4.1 Eosinophils in full blood and overall IgE: simple inflammatory markers

Especially in “western” inflammation profiles of CRSwNP, the eosinophils as well as the overall IgE in the serum are excellent global and derivative markers of Th2 inflammation [235] that are ubiquitously established without any special request and have also been investigated partially in prospective trials such as for example the JESREC cohort (see above) [188]. Also ECP is a promising marker but not easily available [140]. Larger investigations on the sensitivity and specificity are necessary.

4.4.2 Exploratory biomarkers

This field opens an enormous spectrum of possibilities for stratified therapy, however, the number of identified potential markers does not at all correspond to the state of validation, which was recently shown also for the relatively well standardized AIT [236], nonetheless the new markers are scientifically interesting and would have to be verified. On an experimental level, for example new possibilities are seen based on the expression of DPP10 (dipeptidyl peptidase 10) as marker for AERD [237], transglutaminase-2 (TGM2) expression in the tissue as possible marker of AERD-negative endotype [238], and also the WNT signaling [239]. A comprehensive overview of possible local and systemic biomarkers in the airways is found in [240].

A notable example for a Th2-associated biomarker is periostin. The expression in patients with asthma correlated with the successful response to therapy with the anti-IL-13 antibody lebrikizumab [241]. Unfortunately, those very promising data could not be confirmed in the pivotal LAVOLTA trial [242, 243]. It must be discussed if this fact is due to a selection bias of the sample or the biomarker itself. In a post-hoc biomarker trial, Bachert and his research group investigated local and systemic expressions of periostin in different therapy modalities. In this context, it became obvious that the peri-

ostin expression reflects the local and systemic eosinophilic load and was reduced in serum and nasal secretion after application of systemic steroids as well as after application of mepolizumab and omalizumab in CRSwNP [244]. However, now prospective studies are required.

Even where the expression of the local pathophysiological factor would be expected, the results are very heterogenic. Thus, a trial on the treatment of CRSwNP with reslizumab could predict the therapeutic outcome based on the local expression of IL-5; in the context of treatment with mepolizumab, this correlation could not be shown (see above chapter 4.1) [229, 245].

It is unlikely in the authors’s view that only single markers will determine the biomarker situation. It may be speculated that ENT specialists will have to implement the complex data of the findings gained by hypothesis-free analyses and to make therapeutic and even surgical decisions for example supported by algorithms.

4.4.3 Molecular allergology

Due to the detection and characterization of molecular allergens, also the diagnostic precision has been relevantly improved using component-based diagnostics. In this way, allergic sensitizations can be more precisely assessed and thus allowing certain predictability with regard to the therapeutic outcome of AIT. Also for diagnosis of anaphylaxis or exclusion of sensitization against alpha-GAL before application of humanized antibodies such as cetuximab, the component-based diagnostics are essential. The EAACI established an application guideline [246], a more compact version focused on otolaryngology has recently been published also for aeroallergens [247].

5. Conclusion and general reflections on personalized medicine

“Personalized medicine” has become a buzzword even if it has been clearly defined by the WHO. It describes a medical model using molecular profiling technologies for tailoring the right therapeutic strategy for the right person at the right time, and determine the predisposition to disease at the population level and to deliver timely and stratified prevention. Individualized therapy as well as the population-based stratification and prevention play a key role in personalization [248]. The risk to neglect patients by molecular stratification should be taken seriously. We determine endotypes in our algorithms without knowing the complete pathophysiological picture. Avoiding side effects and not treating patients who do not express a certain biomarker or who are characterized as potential non-responders, has nothing in common with individualism and participation in the therapeutic decision-making process and does certainly not reflect the expectations that our patients have with regard to personalized medicine. Hence, this procedure is meanwhile current practice for oncological patients. A molecular determinism possibly puts at risk a well-informed healthcare behavior [249], however, the application of biomarkers (e. g., for blood type or the identification of allergic sensitization) in stratified therapy belongs to the standard of modern medicine (e. g., blood type compatible products, allergen-specific immunotherapy) [250].

If the term of personalized medicine or stratified medicine (which is favored by the author) is established, will have to be expected for the near future.

The difficulties of validating molecular markers in the context of clinical trials illustrate further problems: Is it still ethically justified to perform randomized, double-blind, and placebo-controlled trials with regard to a highly selective patient selection and present mechanistic standards? The example of the LAVOLTA trial demonstrates that we obligatorily need confirmatory and pivotal studies. On the other hand, there is the concern that patients do not want to be included in biomarker studies because they do not want to become victims of possible genetic discrimination [251].

Another important aspect for head and neck surgeons is the significance of endoscopic paranasal sinus surgery in the context of personalized medicine. It is important that we optimally apply this valuable and well-established instrument in the spectrum of therapeutic options and at the same time improve the database by means of controlled trials.

The tension of the terms between personalized medicine and precision medicine raises expectations and implies the illusion of control. Medicine is not only a science and may never be completely "precise" [252]. Competing connotations, interests, and expectations in patients, investors, and cost bearers require a realistic handling with the expectations, but also the well-informed and proactive use of new therapeutic options by human, participative, and rational medicine, also in the field of rhinology.

In this context, the otorhinolaryngologists have 2 interesting, already available perspectives: molecular therapy of chronic rhinosinusitis oriented at relatively simple biomarkers by means of new biologicals as well as specific immunotherapy precised by component-based allergy diagnosis.

Conflict of interest

Via the Technical University of Munich, Germany, Adam Chaker acts as consultant (e. g., advisory boards), gives lectures, or performs further activities for several third parties (Allergopharma, ALK-Abello, GSK, HAL Allergy, Mundipharma, and Lofarma); he received third party and research funding via the Technical University of Munich by ALK, Allergopharma, Novartis, Bencard/Allergen Therapeutics, ASIT Biotech, GSK, Roche, and Zeller AG, furthermore by the Federal Environmental Agency of Germany and the DZL (Deutsches Zentrum für Lungenforschung; German Center for Lung Research [BMBF]). In addition, Chaker received payments and reimbursements by the Bayerisches Ärzteblatt, Deutsche Gesellschaft für Allergologie und Klinische Immunologie (DGAKI, German Society for Allergology and Clinical Immunology), and the European Academy of Allergy and Clinical Immunology.

References

- [1] Boyman O et al. EAACI IG Biologicals task force paper on the use of biologic agents in allergic disorders. *Allergy* 2015; 70: 727–754
- [2] Agency EM. EMA GUIDELINE ON SIMILAR BIOLOGICAL MEDICINAL PRODUCTS. 2005 (http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003517.pdf (10th of October, 2017))
- [3] Kohler G, Milstein C. Continuous cultures of fused cells secreting antibody of predefined specificity. *Nature* 1975; 256: 495–497
- [4] Quianzon CC, Cheikh I. History of insulin. *J Community Hosp Intern Med Perspect* 2012; 2
- [5] Suiter TM. First and next generation native rFVIII in the treatment of hemophilia A. What has been achieved? Can patients be switched safely? *Semin Thromb Hemost* 2002; 28: 277–284
- [6] Beauvais P, Billette de Villemeur T. Prion diseases and blood transfusion. *Transfus Clin Biol* 1999; 6: 24–28
- [7] Frasier SD. The not-so-good old days: working with pituitary growth hormone in North America, 1956 to 1985. *J Pediatr* 1997; 131: S1–S4
- [8] Ayyar VS. History of growth hormone therapy. *Indian J Endocrinol Metab* 2011; 15 (Suppl 3): S162–S165
- [9] Mosmann TR et al. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 1986; 136: 2348–2357
- [10] Mosmann TR, Coffman RL. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989; 7: 145–173
- [11] Bryan SA et al. Effects of recombinant human interleukin-12 on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet* 2000; 356: 2149–2153
- [12] Friedrich M et al. Immunomodulation by interleukin-10 therapy decreases the incidence of relapse and prolongs the relapse-free interval in Psoriasis. *J Invest Dermatol* 2002; 118: 672–677
- [13] Trepicchio WL et al. Interleukin-11 therapy selectively downregulates type I cytokine proinflammatory pathways in psoriasis lesions. *J Clin Invest* 1999; 104: 1527–1537
- [14] Goel N, Chance K. Biosimilars in rheumatology: understanding the rigor of their development. *Rheumatology (Oxford)* 2017; 56: 187–197
- [15] Dorvignit D et al. Expression and biological characterization of an anti-CD20 biosimilar candidate antibody: A case study. *MAbs* 2012; 4: 488–496
- [16] Elliott MJ et al. Randomised double-blind comparison of chimeric monoclonal antibody to tumour necrosis factor alpha (cA2) versus placebo in rheumatoid arthritis. *Lancet* 1994; 344: 1105–1110
- [17] Taylor PC. Developing anti-TNF and biologic agents. *Rheumatology (Oxford)* 2011; 50: 1351–1353
- [18] Upchurch KS, Kay J. Evolution of treatment for rheumatoid arthritis. *Rheumatology (Oxford)* 2012; 51 (Suppl 6): vi28–vi36
- [19] Roque-Navarro L et al. Humanization of predicted T-cell epitopes reduces the immunogenicity of chimeric antibodies: New evidence supporting a simple method. *Hybrid Hybridomics* 2003; 22: 245–257
- [20] Croft M, Benedict CA, Ware CF. Clinical targeting of the TNF and TNFR superfamilies. *Nat Rev Drug Discov* 2013; 12: 147–168
- [21] Parren P, Carter PJ, Pluckthun A. Changes to International Nonproprietary Names for antibody therapeutics 2017 and beyond: of mice, men and more. *MAbs* 2017; 9: 898–906
- [22] Vultaggio A et al. Manifestations of Antidrug Antibodies Response: Hypersensitivity and Infusion Reactions. *J Interferon Cytokine Res* 2014; 34: 946–952
- [23] Vultaggio A et al. Circulating T cells to infliximab are detectable mainly in treated patients developing anti-drug antibodies and hypersensitivity reactions. *Clin Exp Immunol* 2016; 186: 364–372
- [24] Robert F et al. Phase I study of anti-epidermal growth factor receptor antibody cetuximab in combination with radiation therapy in patients with advanced head and neck cancer. *J Clin Oncol* 2001; 19: 3234–3243
- [25] Bourhis J et al. Phase I/II study of cetuximab in combination with cisplatin or carboplatin and fluorouracil in patients with recurrent or metastatic squamous cell carcinoma of the head and neck. *J Clin Oncol* 2006; 24: 2866–2872
- [26] Bonner JA et al. Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck. *N Engl J Med* 2006; 354: 567–578

- [27] O'Neil BH et al. High incidence of cetuximab-related infusion reactions in Tennessee and North Carolina and the association with atopic history. *J Clin Oncol* 2007; 25: 3644–3648
- [28] Chung CH et al. Cetuximab-induced anaphylaxis and IgE specific for galactose-alpha-1,3-galactose. *N Engl J Med* 2008; 358: 1109–1117
- [29] Budach W, Bolke E, Homey B. Severe cutaneous reaction during radiation therapy with concurrent cetuximab. *N Engl J Med* 2007; 357: 514–515
- [30] Commins SP, Platts-Mills TA. Delayed anaphylaxis to red meat in patients with IgE specific for galactose alpha-1,3-galactose (alpha-gal). *Curr Allergy Asthma Rep* 2013; 13: 72–77
- [31] Commins SP, Platts-Mills TA. Tick bites and red meat allergy. *Curr Opin Allergy Clin Immunol* 2013; 13: 354–359
- [32] Galili U. The alpha-gal epitope and the anti-Gal antibody in xenotransplantation and in cancer immunotherapy. *Immunol Cell Biol* 2005; 83: 674–686
- [33] Fischer J et al. Galactose-alpha-1,3-galactose sensitization is a prerequisite for pork-kidney allergy and cofactor-related mammalian meat anaphylaxis. *J Allergy Clin Immunol* 2014; 134: 755–759 e1
- [34] Kleponis J, Skelton R, Zheng L. Fueling the engine and releasing the break: combinational therapy of cancer vaccines and immune checkpoint inhibitors. *Cancer Biol Med* 2015; 12: 201–208
- [35] Perez-Alvarez R et al. Biologics-induced autoimmune diseases. *Curr Opin Rheumatol* 2013; 25: 56–64
- [36] Weise M et al. Biosimilars: the science of extrapolation. *Blood* 2014; 124: 3191–3196
- [37] Ledford H. First biosimilar drug set to enter US market. *Nature* 2015; 517: 253–254
- [38] Kurki P et al. Interchangeability of Biosimilars: A European Perspective. *BioDrugs* 2017; 31: 83–91
- [39] Agency EM. Guideline on similar biological medicinal products. 2013 (http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2014/10/WC500176768.pdf) Accessed on 10th October 2017
- [40] Glue P et al. Pegylated interferon-alpha2b: pharmacokinetics, pharmacodynamics, safety, and preliminary efficacy data. Hepatitis C Intervention Therapy Group. *Clin Pharmacol Ther* 2000; 68: 556–567
- [41] Mok TS et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009; 361: 947–957
- [42] Krueger J et al. Tofacitinib attenuates pathologic immune pathways in patients with psoriasis: A randomized phase 2 study. *J Allergy Clin Immunol* 2016; 137: 1079–1090
- [43] Aguilar-Pimentel A et al. Improved efficacy of allergen-specific immunotherapy by JAK inhibition in a murine model of allergic asthma. *PLoS One* 2017; 12: e0178563
- [44] Boyle MP, De Boeck K. A new era in the treatment of cystic fibrosis: correction of the underlying CFTR defect. *Lancet Respir Med* 2013; 1: 158–163
- [45] Quon BS, Wilcox PG. A new era of personalized medicine for cystic fibrosis – at last!. *Can Respir J* 2015; 22: 257–260
- [46] Krug N et al. Allergen-induced asthmatic responses modified by a GATA3-specific DNase. *N Engl J Med* 2015; 372: 1987–1995
- [47] IMS Institute for Healthcare Informatics. Delivering on the Potential of Biosimilar Medicines. The Role of Functioning Competitive Markets. 2016 (<http://www.medicinesforeurope.com/wp-content/uploads/2016/03/IMS-Institute-Biosimilar-Report-March-2016-FINAL.pdf> am 04.01.2018)
- [48] GKV Spitzenverband Deutschland. Biosimilars aus Sicht der Kostenträger. 2016; https://www.bfarm.de/SharedDocs/Downloads/DE/Service/Termine-und-Veranstaltungen/dialogveranstaltungen/dialog_2016/160627/08_Folien_Haas.pdf?__blob=publicationFile&v=2 zuletzt am 04.01.2018)
- [49] Health, I. Impact of Biosimilar Competition 2016. 2016 ([https://www.imshealth.com/files/web/Market Insights/IMS_Health_Impact_of_Biosimilar_Competition_EU_2016.pdf](https://www.imshealth.com/files/web/Market%20Insights/IMS_Health_Impact_of_Biosimilar_Competition_EU_2016.pdf) online am 10.10.2017)
- [50] Chaker AM, Wagenmann M. Schleimhaut der Nase. *Allergo J* 2014; 23: 1–4
- [51] Pabst R, Chaker A. Integriertes Schleimhautimmunsystem der oberen Atemwege: Intraepitheliale Lymphozyten, NALT und der Waldeyer-sche Rachenring. Biedermann, Heppt, Renz, Röcken: *Allergologie* 2016; 1: 147–155
- [52] Bachert C, Holtappels G. Pathophysiology of chronic rhinosinusitis, pharmaceutical therapy options. *Laryngorhinootologie* 2015; 94 (Suppl 1): S32–S63
- [53] Afzelius BA. Cilia-related diseases. *J Pathol* 2004; 204: 470–477
- [54] Stanke F. The Contribution of the Airway Epithelial Cell to Host Defense. *Mediators Inflamm* 2015; 2015: 463016
- [55] Sisson JH et al. Smoke and viral infection cause cilia loss detectable by bronchoalveolar lavage cytology and dynein ELISA. *Am J Respir Crit Care Med* 1994; 149: 205–213
- [56] Ganz T, Lehrer RI. Defensins. *Pharmacol Ther* 1995; 66: 191–205
- [57] Radicioni G et al. The innate immune properties of airway mucosal surfaces are regulated by dynamic interactions between mucins and interacting proteins: the mucin interactome. *Mucosal Immunol* 2016; 9: 1442–1454
- [58] Cobo ER et al. Colonic MUC2 mucin regulates the expression and antimicrobial activity of beta-defensin 2. *Mucosal Immunol* 2015; 8: 1360–1372
- [59] Delacroix DL et al. IgA subclasses in various secretions and in serum. *Immunology* 1982; 47: 383–385
- [60] Brandtzaeg P. Pillars Article: Mucosal and Glandular Distribution of Immunoglobulin Components: Differential Localization of Free and Bound SC in Secretory Epithelial Cells. *J Immunol* 1974; 112: 1553–1559 *J Immunol* 2017; 198: 1768–1774
- [61] Macpherson AJ et al. The immune geography of IgA induction and function. *Mucosal Immunol* 2008; 1: 11–22
- [62] Pawankar RU et al. Phenotypic and molecular characteristics of nasal mucosal gamma delta T cells in allergic and infectious rhinitis. *Am J Respir Crit Care Med* 1996; 153: 1655–1665
- [63] Mjosberg JM et al. Human IL-25- and IL-33-responsive type 2 innate lymphoid cells are defined by expression of CRTH2 and CD161. *Nat Immunol* 2011; 12: 1055–1062
- [64] Brandtzaeg P, Pabst R. Let's go mucosal: communication on slippery ground. *Trends Immunol* 2004; 25: 570–577
- [65] Chaker AM. Anatomy and Microanatomy of Tonsils. *Encyclopedia of Immunobiology* 2016; 3: 420–426
- [66] Gevaert P et al. Organization of secondary lymphoid tissue and local IgE formation to *Staphylococcus aureus* enterotoxins in nasal polyp tissue. *Allergy* 2005; 60: 71–79
- [67] Fokkens WJ et al. Langerhans cells in nasal mucosa of patients with grass pollen allergy. *Immunobiology* 1991; 182: 135–142
- [68] Fokkens WJ et al. Characterization and quantification of cellular infiltrates in nasal mucosa of patients with grass pollen allergy, non-allergic patients with nasal polyps and controls. *Int Arch Allergy Appl Immunol* 1990; 93: 66–72
- [69] Jahnsen FL et al. Human nasal mucosa contains antigen-presenting cells of strikingly different functional phenotypes. *Am J Respir Cell Mol Biol* 2004; 30: 31–37
- [70] Allam JP et al. Comparative analysis of nasal and oral mucosa dendritic cells. *Allergy* 2006; 61: 166–172
- [71] Soyka MB et al. Defective epithelial barrier in chronic rhinosinusitis: the regulation of tight junctions by IFN-gamma and IL-4. *J Allergy Clin Immunol* 2012; 130: 1087–1096 e10

- [72] Zhou B et al. Thymic stromal lymphopoietin as a key initiator of allergic airway inflammation in mice. *Nat Immunol* 2005; 6: 1047–1053
- [73] Xu G et al. Opposing roles of IL-17A and IL-25 in the regulation of TSLP production in human nasal epithelial cells. *Allergy* 2010; 65: 581–589
- [74] Zissler UM et al. Interleukin-4 and interferon-gamma orchestrate an epithelial polarization in the airways. *Mucosal Immunol* 2016; 9: 917–926
- [75] Soyka MB et al. The Induction of IL-33 in the Sinus Epithelium and Its Influence on T-Helper Cell Responses. *PLoS One* 2015; 10: e0123163
- [76] Bachert C et al. IL-5 synthesis is upregulated in human nasal polyp tissue. *J Allergy Clin Immunol* 1997; 99: 837–842
- [77] Van Zele T et al. Differentiation of chronic sinus diseases by measurement of inflammatory mediators. *Allergy* 2006; 61: 1280–1289
- [78] Bachert C et al. Total and specific IgE in nasal polyps is related to local eosinophilic inflammation. *J Allergy Clin Immunol* 2001; 107: 607–614
- [79] Linneberg A et al. The link between allergic rhinitis and allergic asthma: a prospective population-based study. The Copenhagen Allergy Study. *Allergy* 2002; 57: 1048–1052
- [80] Anderson HR, Bland JM, Peckham CS. Risk factors for asthma up to 16 years of age. Evidence from a national cohort study. *Chest* 1987; 91 (6 Suppl): 1275–1305
- [81] Shaaban R et al. Rhinitis and onset of asthma: A longitudinal population-based study. *Lancet* 2008; 372: 1049–1057
- [82] Jarvis D et al. Asthma in adults and its association with chronic rhinosinusitis: the GA2LEN survey in Europe. *Allergy* 2012; 67: 91–98
- [83] Settiple GA, Chafee FH. Nasal polyps in asthma and rhinitis. A review of 6,037 patients. *J Allergy Clin Immunol* 1977; 59: 17–21
- [84] Bachert C, Zhang L, Gevaert P. Current and future treatment options for adult chronic rhinosinusitis: Focus on nasal polyposis. *J Allergy Clin Immunol* 2015; 136: 1431–1440
- [85] Williamson PA et al. Airway dysfunction in nasal polyposis: A spectrum of asthmatic disease? *Clin Exp Allergy* 2011; 41: 1379–1385
- [86] Hens G et al. Sinonasal pathology in nonallergic asthma and COPD: 'united airway disease' beyond the scope of allergy. *Allergy* 2008; 63: 261–267
- [87] Spergel JM, Paller AS. Atopic dermatitis and the atopic march. *J Allergy Clin Immunol* 2003; 112 (6 Suppl): S118–S127
- [88] Braunstahl GJ et al. Segmental bronchial provocation induces nasal inflammation in allergic rhinitis patients. *Am J Respir Crit Care Med* 2000; 161: 2051–2057
- [89] Braunstahl GJ et al. Nasal allergen provocation induces adhesion molecule expression and tissue eosinophilia in upper and lower airways. *J Allergy Clin Immunol* 2001; 107: 469–476
- [90] Braunstahl GJ et al. Segmental bronchoprovocation in allergic rhinitis patients affects mast cell and basophil numbers in nasal and bronchial mucosa. *Am J Respir Crit Care Med* 2001; 164: 858–865
- [91] Assanasen P et al. The nasal passage of subjects with asthma has a decreased ability to warm and humidify inspired air. *Am J Respir Crit Care Med* 2001; 164: 1640–1646
- [92] Naclerio RM et al. Observations on the ability of the nose to warm and humidify inspired air. *Rhinology* 2007; 45: 102–111
- [93] Eccles R. Is the common cold a clinical entity or a cultural concept? *Rhinology* 2013; 51: 3–8
- [94] Lewis-Rogers N, Bendall ML, Crandall KA. Phylogenetic relationships and molecular adaptation dynamics of human rhinoviruses. *Mol Biol Evol* 2009; 26: 969–981
- [95] Palmenberg AC et al. Sequencing and analyses of all known human rhinovirus genomes reveal structure and evolution. *Science* 2009; 324: 55–59
- [96] Caliskan M et al. Rhinovirus wheezing illness and genetic risk of childhood-onset asthma. *N Engl J Med* 2013; 368: 1398–1407
- [97] Bianco A et al. Th2 cytokines exert a dominant influence on epithelial cell expression of the major group human rhinovirus receptor, ICAM-1. *Eur Respir J* 1998; 12: 619–626
- [98] Xepapadaki P et al. Duration of postviral airway hyperresponsiveness in children with asthma: effect of atopy. *J Allergy Clin Immunol* 2005; 116: 299–304
- [99] Toussaint M et al. Host DNA released by NETosis promotes rhinovirus-induced type-2 allergic asthma exacerbation. *Nat Med* 2017; 23: 681–691
- [100] Bartlett NW et al. Mouse models of rhinovirus-induced disease and exacerbation of allergic airway inflammation. *Nat Med* 2008; 14: 199–204
- [101] Institut, R.K. Influenza. Robert Koch Website, online am 10.10.2017: p. http://www.rki.de/DE/Content/InfAZ/I/Influenza/IPV/IPV_Node.html
- [102] Zomer-Kooijker K et al. Increased risk of wheeze and decreased lung function after respiratory syncytial virus infection. *PLoS One* 2014; 9: e87162
- [103] Sigurs N et al. Respiratory syncytial virus bronchiolitis in infancy is an important risk factor for asthma and allergy at age 7. *Am J Respir Crit Care Med* 2000; 161: 1501–1507
- [104] American Academy of Pediatrics Committee on Infectious, D. and C. American Academy of Pediatrics Bronchiolitis Guidelines, Updated guidance for palivizumab prophylaxis among infants and young children at increased risk of hospitalization for respiratory syncytial virus infection. *Pediatrics* 2014; 134: e620–e638
- [105] Heinecke L et al. Induction of B7-H1 and B7-DC expression on airway epithelial cells by the Toll-like receptor 3 agonist double-stranded RNA and human rhinovirus infection: In vivo and in vitro studies. *J Allergy Clin Immunol* 2008; 121: 1155–1160
- [106] Seshadri S et al. Increased expression of the epithelial anion transporter pendrin/SLC26A4 in nasal polyps of patients with chronic rhinosinusitis. *J Allergy Clin Immunol* 2015; 136: 1548–1558 e7
- [107] Fokkens WJ et al. EPOS 2012: European position paper on rhinosinusitis and nasal polyps 2012. *Rhinology* 2012; 50 (Suppl 23): 1–305
- [108] Wang DY et al. A survey on the management of acute rhinosinusitis among Asian physicians. *Rhinology* 2011; 49: 264–271
- [109] Revai K et al. Incidence of acute otitis media and sinusitis complicating upper respiratory tract infection: the effect of age. *Pediatrics* 2007; 119: e1408–e1412
- [110] Wang JH, Kwon HJ, Jang YJ. Rhinovirus enhances various bacterial adhesions to nasal epithelial cells simultaneously. *Laryngoscope* 2009; 119: 1406–1411
- [111] Woytschak J et al. Type 2 Interleukin-4 Receptor Signaling in Neutrophils Antagonizes Their Expansion and Migration during Infection and Inflammation. *Immunity* 2016; 45: 172–184
- [112] Cohen M et al. Biofilms in chronic rhinosinusitis: a review. *Am J Rhinol Allergy* 2009; 23: 255–260
- [113] Abreu NA et al. Sinus microbiome diversity depletion and *Corynebacterium tuberculostrictum* enrichment mediates rhinosinusitis. *Sci Transl Med* 2012; 4: 151ra124
- [114] Choi EB et al. Decreased diversity of nasal microbiota and their secreted extracellular vesicles in patients with chronic rhinosinusitis based on a metagenomic analysis. *Allergy* 2014; 69: 517–526
- [115] Corriveau MN et al. Detection of *Staphylococcus aureus* in nasal tissue with peptide nucleic acid-fluorescence in situ hybridization. *Am J Rhinol Allergy* 2009; 23: 461–465

- [116] Song WJ et al. Staphylococcal enterotoxin sensitization in a community-based population: a potential role in adult-onset asthma. *Clin Exp Allergy* 2014; 44: 553–562
- [117] Akdis M et al. Skin homing (cutaneous lymphocyte-associated antigen-positive) CD8 + T cells respond to superantigen and contribute to eosinophilia and IgE production in atopic dermatitis. *J Immunol* 1999; 163: 466–475
- [118] Vrieze A et al. Transfer of intestinal microbiota from lean donors increases insulin sensitivity in individuals with metabolic syndrome. *Gastroenterology* 2012; 143: 913–916 e7
- [119] Hastan D et al. Chronic rhinosinusitis in Europe – an underestimated disease. A GA(2)LEN study. *Allergy* 2011; 66: 1216–1223
- [120] Beule AG. [Epidemiology of chronic rhinosinusitis, selected risk factors, comorbidities and economic burden]. *Laryngorhinootologie* 2015; 94 (Suppl 1): S1–S23
- [121] Bousquet J et al. Allergic Rhinitis and its Impact on Asthma (ARIA): Achievements in 10 years and future needs. *J Allergy Clin Immunol* 2012; 130: 1049–1062
- [122] Bousquet J et al. Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2)LEN and AllerGen). *Allergy* 2008; 63 (Suppl 86): 8–160
- [123] Pearce N et al. Self-reported prevalence of asthma symptoms in children in Australia, England, Germany and New Zealand: An international comparison using the ISAAC protocol. *Eur Respir J* 1993; 6: 1455–1461
- [124] Langen U, Schmitz R, Steppuhn H. Prevalence of allergic diseases in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013; 56: 698–706
- [125] Haftenberger M et al. Prevalence of sensitisation to aeroallergens and food allergens: Results of the German Health Interview and Examination Survey for Adults (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013; 56: 687–697
- [126] Schmitz R, Kuhnert R, Thamm M. 12-Monats-Prävalenz von Allergien in Deutschland. *Journal of Health Monitoring* 2017; 2: 77–82
- [127] Meltzer EO et al. Sleep, quality of life, and productivity impact of nasal symptoms in the United States: findings from the Burden of Rhinitis in America survey. *Allergy Asthma Proc* 2009; 30: 244–254
- [128] Biermann J, MHF Wehrmann W, Klimek L, Wasem J. Allergische Erkrankungen der Atemwege-Ergebnisse einer umfassenden Patientenkohorte in der deutschen gesetzlichen Krankenversicherung. *Allergo Journal* 2013; 22: 363–373
- [129] Pfaar O et al. Guideline on allergen-specific immunotherapy in IgE-mediated allergic diseases: S2k Guideline of the German Society for Allergology and Clinical Immunology (DGAKI), the Society for Pediatric Allergy and Environmental Medicine (GPA), the Medical Association of German Allergologists (AeDA), the Austrian Society for Allergy and Immunology (OGAI), the Swiss Society for Allergy and Immunology (SGAI), the German Society of Dermatology (DDG), the German Society of Oto- Rhino-Laryngology, Head and Neck Surgery (DGHNO-KHC), the German Society of Pediatrics and Adolescent Medicine (DGKJ), the Society for Pediatric Pneumology (GPP), the German Respiratory Society (DGP), the German Association of ENT Surgeons (BV-HNO), the Professional Federation of Paediatricians and Youth Doctors (BVKJ), the Federal Association of Pulmonologists (BDP) and the German Dermatologists Association (BVDD). *Allergo J Int* 2014; 23: 282–319
- [130] Calderon MA et al. Allergen injection immunotherapy for seasonal allergic rhinitis. *Cochrane Database Syst Rev* 2007; CD001936
- [131] Radulovic S et al. Sublingual immunotherapy for allergic rhinitis. *Cochrane Database Syst Rev* 2010; CD002893
- [132] Radulovic S et al. Systematic reviews of sublingual immunotherapy (SLIT). *Allergy* 2011; 66: 740–752
- [133] Nelson H et al. Network meta-analysis shows commercialized subcutaneous and sublingual grass products have comparable efficacy. *J Allergy Clin Immunol Pract* 2015; 3: 256–266 e3
- [134] Jacobsen L et al. Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy* 2007; 62: 943–948
- [135] Niggemann B et al. Five-year follow-up on the PAT study: specific immunotherapy and long-term prevention of asthma in children. *Allergy* 2006; 61: 855–859
- [136] Valovirta E et al. Results from the 5-year SQ grass sublingual immunotherapy tablet asthma prevention (GAP) trial in children with grass pollen allergy. *J Allergy Clin Immunol* 2017 Jul 6. pii: S0091–6749(17)31088–6
- [137] Kurukulaaratchy RJ et al. Identifying the heterogeneity of young adult rhinitis through cluster analysis in the Isle of Wight birth cohort. *J Allergy Clin Immunol* 2015; 135: 143–150
- [138] Glowania A, CM Klimek L, Chaker A. Rhinitis – allergic or not? *Allergo Journal* 2012; 08: 486–498
- [139] Hellings PW et al. Non-allergic rhinitis: Position paper of the European Academy of Allergy and Clinical Immunology. *Allergy* 2017; 72: 1657–1665
- [140] Becker S et al. Non-allergic rhinitis with eosinophilia syndrome is not associated with local production of specific IgE in nasal mucosa. *Eur Arch Otorhinolaryngol* 2016; 273: 1469–1475
- [141] Van Gerven L et al. Capsaicin treatment reduces nasal hyperreactivity and transient receptor potential cation channel subfamily V, receptor 1 (TRPV1) overexpression in patients with idiopathic rhinitis. *J Allergy Clin Immunol* 2014; 133: 1332–1339 e1–e3
- [142] Wagenmann M, Scheckenbach K, Chaker AM. Endotypes in Chronic Rhinosinusitis: Biomarkers Based on a Mechanistic Insight for Targeted Treatment? *ORL J Otorhinolaryngol Relat Spec* 2017; 79: 78–84
- [143] Hirsch AG et al. Nasal and sinus symptoms and chronic rhinosinusitis in a population-based sample. *Allergy* 2017; 72: 274–281
- [144] Rudmik L, Smith TL. Quality of life in patients with chronic rhinosinusitis. *Curr Allergy Asthma Rep* 2011; 11: 247–252
- [145] Rajan JP et al. Prevalence of aspirin-exacerbated respiratory disease among asthmatic patients: A meta-analysis of the literature. *J Allergy Clin Immunol* 2015; 135 (676/81): e1
- [146] Zhang N et al. An update on the impact of Staphylococcus aureus enterotoxins in chronic sinusitis with nasal polyposis. *Rhinology* 2005; 43: 162–168
- [147] Knopf A et al. Rheumatic disorders affecting the head and neck: Underestimated diseases. *Rheumatology (Oxford)* 2011; 50: 2029–2034
- [148] Hofauer B et al. Liposomal local therapy of sinusal symptoms in ANCA associated vasculitis. *Laryngorhinootologie* 2014; 93: 461–466
- [149] Woywodt A, Matteson EL. Wegener's granulomatosis – probing the untold past of the man behind the eponym. *Rheumatology (Oxford)* 2006; 45: 1303–1306
- [150] Craven A et al. ACR/EULAR-endorsed study to develop Diagnostic and Classification Criteria for Vasculitis (DCVAS). *Clin Exp Nephrol* 2013; 17: 619–621
- [151] Yates M et al. EULAR/ERA-EDTA recommendations for the management of ANCA-associated vasculitis. *Ann Rheum Dis* 2016; 75: 1583–1594
- [152] Wechsler ME et al. Mepolizumab or Placebo for Eosinophilic Granulomatosis with Polyangiitis. *N Engl J Med* 2017; 376: 1921–1932
- [153] De Gaudemar I et al. Is nasal polyposis in cystic fibrosis a direct manifestation of genetic mutation or a complication of chronic infection? *Rhinology* 1996; 34: 194–197

- [154] Claeys S et al. Nasal polyps in patients with and without cystic fibrosis: a differentiation by innate markers and inflammatory mediators. *Clin Exp Allergy* 2005; 35: 467–472
- [155] Steinke JW, Borish L. Chronic rhinosinusitis phenotypes. *Ann Allergy Asthma Immunol* 2016; 117: 234–240
- [156] Kartagener M. Zur Pathogenese der Bronchiektasien bei Situs viscerum inversus. *Beiträge Klin. Tuberk* 1933; 83: 489–511
- [157] Degano B et al. Expression of nitric oxide synthases in primary ciliary dyskinesia. *Hum Pathol* 2011; 42: 1855–1861
- [158] Marthin JK, Nielsen KG. Choice of nasal nitric oxide technique as first-line test for primary ciliary dyskinesia. *Eur Respir J* 2011; 37: 559–565
- [159] Joensen O et al. Exhaled breath analysis using electronic nose in cystic fibrosis and primary ciliary dyskinesia patients with chronic pulmonary infections. *PLoS One* 2014; 9: e115584
- [160] Soler ZM et al. Cluster analysis and prediction of treatment outcomes for chronic rhinosinusitis. *J Allergy Clin Immunol* 2016; 137: 1054–1062
- [161] Derycke L et al. Mixed T helper cell signatures in chronic rhinosinusitis with and without polyps. *PLoS One* 2014; 9: e97581
- [162] Zhang N et al. Different types of T-effector cells orchestrate mucosal inflammation in chronic sinus disease. *J Allergy Clin Immunol* 2008; 122: 961–968
- [163] Katotomichelakis M et al. Inflammatory patterns in upper airway disease in the same geographical area may change over time. *Am J Rhinol Allergy* 2013; 27: 354–360
- [164] Ba L et al. The association between bacterial colonization and inflammatory pattern in Chinese chronic rhinosinusitis patients with nasal polyps. *Allergy* 2011; 66: 1296–1303
- [165] Derycke L et al. IL-17A as a regulator of neutrophil survival in nasal polyp disease of patients with and without cystic fibrosis. *J Cyst Fibros* 2012; 11: 193–200
- [166] Wen W et al. Increased neutrophilia in nasal polyps reduces the response to oral corticosteroid therapy. *J Allergy Clin Immunol* 2012; 129: 1522–1528 e5
- [167] Milara J et al. Mucin 1 downregulation associates with corticosteroid resistance in chronic rhinosinusitis with nasal polyps. *J Allergy Clin Immunol* 2015; 135: 470–476
- [168] Anderson GP. Endotyping asthma: New insights into key pathogenic mechanisms in a complex, heterogeneous disease. *Lancet* 2008; 372: 1107–1119
- [169] Lotvall J et al. Asthma endotypes: A new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol* 2011; 127: 355–360
- [170] Wenzel SE. Complex phenotypes in asthma: Current definitions. *Pulm Pharmacol Ther* 2013; 26: 710–715
- [171] Tomassen P et al. Inflammatory endotypes of chronic rhinosinusitis based on cluster analysis of biomarkers. *J Allergy Clin Immunol* 2016; 137: 1449–1456 e4
- [172] De Greve G et al. Endotype-driven treatment in chronic upper airway diseases. *Clin Transl Allergy* 2017; 7: 22
- [173] Akdis CA et al. Endotypes and phenotypes of chronic rhinosinusitis: A PRACTALL document of the European Academy of Allergy and Clinical Immunology and the American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol* 2013; 131: 1479–1490
- [174] Schmitt J et al. Allergy immunotherapy for allergic rhinitis effectively prevents asthma: Results from a large retrospective cohort study. *J Allergy Clin Immunol* 2015; 136: 1511–1516
- [175] Hopkins C, Andrews P, Holy CE. Does time to endoscopic sinus surgery impact outcomes in chronic rhinosinusitis? Retrospective analysis using the UK clinical practice research data. *Rhinology* 2015; 53: 18–24
- [176] Benninger MS et al. Impact of medically recalcitrant chronic rhinosinusitis on incidence of asthma. *Int Forum Allergy Rhinol* 2016; 6: 124–129
- [177] Hopkins C, Rimmer J, Lund VJ. Does time to endoscopic sinus surgery impact outcomes in Chronic Rhinosinusitis? Prospective findings from the National Comparative Audit of Surgery for Nasal Polyposis and Chronic Rhinosinusitis. *Rhinology* 2015; 53: 10–17
- [178] Benninger MS et al. Early versus delayed endoscopic sinus surgery in patients with chronic rhinosinusitis: impact on health care utilization. *Otolaryngol Head Neck Surg* 2015; 152: 546–552
- [179] Stuck BA et al. Guideline for “rhinosinusitis”-long version: S2k guideline of the German College of General Practitioners and Family Physicians and the German Society for Oto-Rhino-Laryngology, Head and Neck Surgery. *HNO* 2017
- [180] Messerklinger W. The ethmoidal infundibulum and its inflammatory illnesses (author’s transl). *Arch Otorhinolaryngol* 1979; 222: 11–22
- [181] Stammberger H, Posawetz W. Functional endoscopic sinus surgery. Concept, indications and results of the Messerklinger technique. *Eur Arch Otorhinolaryngol* 1990; 247: 63–76
- [182] Rimmer J et al. Surgical versus medical interventions for chronic rhinosinusitis with nasal polyps. *Cochrane Database Syst Rev* 2014; CD006991
- [183] Sharma R et al. Surgical interventions for chronic rhinosinusitis with nasal polyps. *Cochrane Database Syst Rev* 2014; CD006990
- [184] Rudmik L et al. Topical therapies in the management of chronic rhinosinusitis: an evidence-based review with recommendations. *Int Forum Allergy Rhinol* 2013; 3: 281–298
- [185] Kalish L et al. WITHDRAWN: Topical steroids for nasal polyps. *Cochrane Database Syst Rev* 2016; 4: CD006549
- [186] Chong LY et al. Intranasal steroids versus placebo or no intervention for chronic rhinosinusitis. *Cochrane Database Syst Rev* 2016; 4: CD011996
- [187] Chong LY et al. Different types of intranasal steroids for chronic rhinosinusitis. *Cochrane Database Syst Rev* 2016; 4: CD011993
- [188] Tokunaga T et al. Novel scoring system and algorithm for classifying chronic rhinosinusitis: the JESREC Study. *Allergy* 2015; 70: 995–1003
- [189] Hellings PW et al. Uncontrolled allergic rhinitis and chronic rhinosinusitis: where do we stand today? *Allergy* 2013; 68: 1–7
- [190] Philpott C et al. The burden of revision sinonasal surgery in the UK-data from the Chronic Rhinosinusitis Epidemiology Study (CRES): A cross-sectional study. *BMJ Open* 2015; 5: e006680
- [191] Nabi S et al. Nasal spray adherence after sinus surgery: Problems and predictors. *J Otolaryngol Head Neck Surg* 2012; 41 (Suppl 1): S49–S55
- [192] Bousquet J et al. Unmet needs in severe chronic upper airway disease (SCUAD). *J Allergy Clin Immunol* 2009; 124: 428–433
- [193] Ishizaka K, Ishizaka T, Hornbrook MM. Allergen-binding activity of gamma-E, gamma-G and gamma-A antibodies in sera from atopic patients. In vitro measurements of reaginic antibody. *J Immunol* 1967; 98: 490–501
- [194] Johansson SG, Bennich H. Immunological studies of an atypical (myeloma) immunoglobulin. *Immunology* 1967; 13: 381–394
- [195] Johansson SG. Raised levels of a new immunoglobulin class (IgND) in asthma. *Lancet* 1967; 2: 951–953
- [196] Ring J, Bergmann C. Geschichte der Allergologie. Biedermann, Heppt, Renz, Röcken: *Allergologie* 2016; 1: 3–9
- [197] Casale TB et al. Effect of omalizumab on symptoms of seasonal allergic rhinitis: A randomized controlled trial. *JAMA* 2001; 286: 2956–2967
- [198] Chervinsky P et al. Omalizumab, an anti-IgE antibody, in the treatment of adults and adolescents with perennial allergic rhinitis. *Ann Allergy Asthma Immunol* 2003; 91: 160–167

- [199] Kopp MV et al. The effect of anti-IgE treatment on in vitro leukotriene release in children with seasonal allergic rhinitis. *J Allergy Clin Immunol* 2002; 110: 728–735
- [200] Klunker S et al. Combination treatment with omalizumab and rush immunotherapy for ragweed-induced allergic rhinitis: Inhibition of IgE-facilitated allergen binding. *J Allergy Clin Immunol* 2007; 120: 688–695
- [201] Kopp MV et al. Combination of omalizumab and specific immunotherapy is superior to immunotherapy in patients with seasonal allergic rhinoconjunctivitis and co-morbid seasonal allergic asthma. *Clin Exp Allergy* 2009; 39: 271–279
- [202] MacGinnitie AJ et al. Omalizumab facilitates rapid oral desensitization for peanut allergy. *J Allergy Clin Immunol* 2017; 139: 873–881 e8
- [203] Kopp MV. Role of immunomodulators in allergen-specific immunotherapy. *Allergy* 2011; 66: 792–797
- [204] Akdis CA, Akdis M. Mechanisms and treatment of allergic disease in the big picture of regulatory T cells. *J Allergy Clin Immunol* 2009; 123: 735–746 quiz 747–748
- [205] Mobs C et al. Birch pollen immunotherapy results in long-term loss of Bet v 1-specific TH2 responses, transient TR1 activation, and synthesis of IgE-blocking antibodies. *J Allergy Clin Immunol* 2012; 130: 1108–1116 e6
- [206] Wambre E et al. Differentiation stage determines pathologic and protective allergen-specific CD4 + T-cell outcomes during specific immunotherapy. *J Allergy Clin Immunol* 2012; 129: 544–551 551 e1–e7
- [207] Gabrielsson S et al. Specific immunotherapy prevents increased levels of allergen-specific IL-4- and IL-13-producing cells during pollen season. *Allergy* 2001; 56: 293–300
- [208] Michaud B et al. Quantification of circulating house dust mite-specific IL-4- and IL-13-secreting T cells correlates with rhinitis severity in asthmatic children and varies with the seasons. *Clin Exp Allergy* 2014; 44: 222–230
- [209] Francis JN, Till SJ, Durham SR. Induction of IL-10 + CD4 + CD25 + T cells by grass pollen immunotherapy. *J Allergy Clin Immunol* 2003; 111: 1255–1261
- [210] Radulovic S et al. Grass pollen immunotherapy induces Foxp3-expressing CD4 + CD25 + cells in the nasal mucosa. *J Allergy Clin Immunol* 2008; 121: 1467–1472 1472 e1
- [211] Robinson DS, Larche M, Durham SR. Tregs and allergic disease. *J Clin Invest* 2004; 114: 1389–1397
- [212] Till S et al. IL-5 production by allergen-stimulated T cells following grass pollen immunotherapy for seasonal allergic rhinitis. *Clin Exp Immunol* 1997; 110: 114–121
- [213] Yang M et al. Interleukin-13 mediates airways hyperreactivity through the IL-4 receptor-alpha chain and STAT-6 independently of IL-5 and eotaxin. *Am J Respir Cell Mol Biol* 2001; 25: 522–530
- [214] Mantel PY et al. GATA3-driven Th2 responses inhibit TGF-beta1-induced FOXP3 expression and the formation of regulatory T cells. *PLoS Biol* 2007; 5
- [215] Chaker AM et al. Short-term subcutaneous grass pollen immunotherapy under the umbrella of anti-IL-4: A randomized controlled trial. *J Allergy Clin Immunol* 2016; 137: 452–461 e9
- [216] Johansson SG. IgE in allergic diseases. *Proc R Soc Med* 1969; 62: 975–976
- [217] Donovan R et al. Immunoglobulins in nasal polyp fluid. *Int Arch Allergy Appl Immunol* 1970; 37: 154–166
- [218] Whiteside TL et al. The presence of IgE on the surface of lymphocytes in nasal polyps. *J Allergy Clin Immunol* 1975; 55: 186–194
- [219] Van Zele T et al. Staphylococcus aureus colonization and IgE antibody formation to enterotoxins is increased in nasal polyposis. *J Allergy Clin Immunol* 2004; 114: 981–983
- [220] Penn R, Mikula S. The role of anti-IgE immunoglobulin therapy in nasal polyposis: A pilot study. *Am J Rhinol* 2007; 21: 428–432
- [221] Guglielmo M et al. Recalcitrant nasal polyposis: achievement of total remission following treatment with omalizumab. *J Investig Allergol Clin Immunol* 2009; 19: 158–159
- [222] Vennera Mdel C et al. Efficacy of omalizumab in the treatment of nasal polyps. *Thorax* 2011; 66: 824–825
- [223] Pinto JM et al. A randomized, double-blind, placebo-controlled trial of anti-IgE for chronic rhinosinusitis. *Rhinology* 2010; 48: 318–324
- [224] Gevaert P et al. Omalizumab is effective in allergic and nonallergic patients with nasal polyps and asthma. *J Allergy Clin Immunol* 2013; 131: 110–116 e1
- [225] Lowe PJ, Renard D. Omalizumab decreases IgE production in patients with allergic (IgE-mediated) asthma; PKPD analysis of a biomarker, total IgE. *Br J Clin Pharmacol* 2011; 72: 306–320
- [226] Castro M et al. Reslizumab for poorly controlled, eosinophilic asthma: a randomized, placebo-controlled study. *Am J Respir Crit Care Med* 2011; 184: 1125–1132
- [227] Haldar P et al. Mepolizumab and exacerbations of refractory eosinophilic asthma. *N Engl J Med* 2009; 360: 973–984
- [228] Nair P et al. Mepolizumab for prednisone-dependent asthma with sputum eosinophilia. *N Engl J Med* 2009; 360: 985–993
- [229] Gevaert P et al. Mepolizumab, a humanized anti-IL-5 mAb, as a treatment option for severe nasal polyposis. *J Allergy Clin Immunol* 2011; 128: 989–995 e1–e8
- [230] Bachert C et al. Reduced need for surgery in severe nasal polyposis with mepolizumab: Randomized trial. *J Allergy Clin Immunol* 2017; 140: 1024–1031 e14
- [231] Casale TB. Biologics and biomarkers for asthma, urticaria, and nasal polyposis. *J Allergy Clin Immunol* 2017; 139: 1411–1421
- [232] Wenzel S et al. Dupilumab in persistent asthma with elevated eosinophil levels. *N Engl J Med* 2013; 368: 2455–2466
- [233] Beck LA et al. Dupilumab treatment in adults with moderate-to-severe atopic dermatitis. *N Engl J Med* 2014; 371: 130–139
- [234] Simpson EL et al. Two Phase 3 Trials of Dupilumab versus Placebo in Atopic Dermatitis. *N Engl J Med* 2016; 375: 2335–2348
- [235] Jung YG et al. Predictive capabilities of serum eosinophil cationic protein, percentage of eosinophils and total immunoglobulin E in allergic rhinitis without bronchial asthma. *J Int Med Res* 2011; 39: 2209–2216
- [236] Shamji MH et al. Biomarkers for monitoring clinical efficacy of allergen immunotherapy for allergic rhinoconjunctivitis and allergic asthma: an EAACI Position Paper. *Allergy* 2017; 72: 1156–1173
- [237] Kim SH et al. Dipeptidyl-peptidase 10 as a genetic biomarker for the aspirin-exacerbated respiratory disease phenotype. *Ann Allergy Asthma Immunol* 2015; 114: 208–213
- [238] Dietz K et al. Age dictates a steroid-resistant cascade of Wnt5a, transglutaminase 2, and leukotrienes in inflamed airways. *J Allergy Clin Immunol* 2017; 139: 1343–1354.e6
- [239] Boscke R et al. Wnt Signaling in Chronic Rhinosinusitis with Nasal Polyps. *Am J Respir Cell Mol Biol* 2017; 56: 575–584
- [240] Zissler UM et al. Current and future biomarkers in allergic asthma. *Allergy* 2016; 71: 475–494
- [241] Corren J et al. Lebrikizumab treatment in adults with asthma. *N Engl J Med* 2011; 365: 1088–1098
- [242] Bujarski S, Parulekar AD, Hanania NA. Lebrikizumab in the treatment of asthma. *Expert Opin Biol Ther* 2016
- [243] Hanania NA et al. Efficacy and safety of lebrikizumab in patients with uncontrolled asthma (LAVOLTA I and LAVOLTA II): Replicate, phase 3, randomised, double-blind, placebo-controlled trials. *Lancet Respir Med* 2016; 4: 781–796

- [244] De Schryver E et al. The effect of systemic treatments on periostin expression reflects their interference with the eosinophilic inflammation in chronic rhinosinusitis with nasal polyps. *Rhinology* 2017; 55: 152–160
- [245] Gevaert P et al. Nasal IL-5 levels determine the response to anti-IL-5 treatment in patients with nasal polyps. *J Allergy Clin Immunol* 2006; 118: 1133–1141
- [246] Matricardi PM et al. EAACI Molecular Allergology User's Guide. *Pediatr Allergy Immunol* 2016; 27 (Suppl 23): 1–250
- [247] Klimek L, Becker S. Molecular component-resolved allergy diagnostics in ENT. *HNO* 2017; 65: 818–825
- [248] Chaker AM, Klimek L. Individualized, personalized and stratified medicine: a challenge for allergology in ENT? *HNO* 2015; 63: 334–342
- [249] Ethikrat Deutschland, Forum Bioethik: Die Medizin nimmt's persönlich. 2009 (http://www.ethikrat.org/dateien/pdf/fb_2009-06-24_simultanmitschrift.pdf am 04.01.2018)
- [250] Muller-Berghaus J et al. Special considerations for the regulation of biological medicinal products in individualised medicine. More than stratified medicine. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013; 56: 1538–1544
- [251] Green RC, Lautenbach D, McGuire AL. GINA, genetic discrimination, and genomic medicine. *N Engl J Med* 2015; 372: 397–399
- [252] Siest G. Systems medicine, stratified medicine, personalized medicine but not precision medicine. *Drug Metabol Drug Interact* 2014; 29: 1–2