



Relationship between Diabetes Self-Management and the Use of Health Care Apps: A Cross-Sectional Study

Satoshi Inagaki¹ Kenji Kato² Kozue Abe³ Hiroaki Takahashi³ Tomokazu Matsuda³

¹ Faculty of Nursing, Kobe City College of Nursing, Kobe, Japan

² Faculty of Nursing, Kobe Women's University, Kobe, Hyōgo, Japan

³ Matsuda Diabetes Clinic, Kobe, Japan

Address for correspondence Satoshi Inagaki, MS, Kobe City College of Nursing, 3-4 Gakuen-nishi-machi, Nishi-ku, Kobe City, Hyogo 651-2103, Japan (e-mail: s1inagaki@tr.kobe-ccn.ac.jp).

ACI Open 2023;7:e23–e29.

Abstract

Keywords

- ▶ patient self-care
- ▶ home care and e-health
- ▶ telemedicine and telehealth
- ▶ consumer health informatics
- ▶ diabetes mellitus
- ▶ chronic disease

Background People with diabetes are increasingly using smartphone health care applications (apps) to manage their health. However, few studies have examined the percentage of people with diabetes using health care apps and their relationship to self-care.

Objective The purpose of this study is to determine the prevalence of health care apps among people with diabetes and the relationship between app use and self-management.

Methods A cross-sectional study was conducted using an online survey among people with type 2 diabetes. Multiple linear regression analysis was conducted using the scores of the Japanese version of Summary of Diabetes Self-Care Activities and exercise and general diet subscales as the objective variables.

Results Of 253 participants included in this study, 61 (24.1%) used health care apps. Approximately 20% of those aged ≥ 60 also used health care apps. Use of health care apps was a significant predictor of physical activity frequency along with autonomous motivation ($p < 0.001$). Participants who used health care apps showed a 0.91 point higher physical activity score than those who did not. Regarding the general diet score, the use of health care apps was not significantly associated with dietary habits ($p = 0.29$).

Conclusion Among people with type 2 diabetes, 24.1% used health care apps, and self-management scores of exercise were significantly higher in people with diabetes who used health care apps than in those who did not.

Background and Significance

Up to 2019, the number of people with diabetes mellitus among adults was approximately 463 million (9.3% of the total population) worldwide.¹ People with diabetes are at a high risk of developing long-term vascular complications, causing considerable mortality.² Therefore, it is important to maintain good glycemic control through health care behaviors, such as healthy diets, exercises, and appropriate medications.

To maintain good blood glucose control and prevent diabetes and its complications, various Diabetes Self-Management Support (DSMS) programs have been developed.^{3,4} DSMS helps people implement and maintain coping strategies and behaviors necessary for diabetes self-management.⁵ Participation in DSMS programs improves and strengthens diabetes knowledge, self-management skills (e.g., meal planning and physical activity), and health status.^{6,7} DSMS is typically performed in community-based settings, such as

received

July 27, 2022

accepted after revision

February 16, 2023

DOI <https://doi.org/10.1055/s-0043-1766113>.

ISSN 2566-9346.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

clinics or community organizations; however, methods employing smartphone health care applications (apps) are also gaining attention.^{8–10}

Generally, health care apps are intended to provide health benefits to users, such as encouraging health management habits and disseminating health knowledge.^{11–13} Several types of health care apps have been developed, and currently, there are more than 500,000 health care apps available in various online stores,¹⁴ such as the Mac App Store and Google Play Store. Moreover, numerous health care apps have been developed to support self-management behaviors among people with diabetes.¹⁵ Several studies have reported changes in diabetes self-management behaviors after using these apps for several months.^{16–19} Although individual apps have been reported to be beneficial for people with diabetes, due to language and usability barriers,^{20,21} people with diabetes do not always have access to validated apps. In other words, it is assumed that some patients are using apps that have not been validated for their usefulness to people with diabetes. Therefore, to determine whether the use of health care apps is associated with self-care, we surveyed people with diabetes regarding their use of health care apps and their self-care status. Few studies have examined the percentage of people with diabetes using health care apps and their relationship to self-care in this way. It is important to conduct this basic study because determining the prevalence of health care apps and their relationship to self-care could provide evidence for clinicians to use such apps to promote DSMS. Health care apps generally introduced to people with diabetes are of the type that support self-management by recording exercise and diet and monitoring/reporting blood glucose levels and medication status.²² However, since the majority of people with diabetes in Japan, approximately 80%, do not manage their blood glucose levels,^{23,24} we conducted this survey focusing on self-care, particularly exercise and dietary lifestyle.

Objective

The purpose of this study is to investigate the use of health care apps by people with type 2 diabetes and to describe the relationship between diabetes self-management behaviors and the use of health care apps.

Methods

Study Design

An Internet-based survey of individuals who had preregistered with a research firm was used to conduct this cross-sectional study. Data were collected in June 2021.

Participants

The participants were recruited from a panel of people with type 2 diabetes who were registered with a research firm. The selection criteria were as follows: people with type 2 diabetes, aged ≥ 18 years, ability to read and write Japanese language, and ability to provide informed consent. No exclusion criteria were set.

All sampling/recruitment and the survey were conducted within the Web page of the research firm. A two-stage sampling method was employed. In the first stage, people with type 2 diabetes who had preregistered with a research firm were asked to complete a screening survey. In the screening survey, they were inquired regarding the reasons for their medical visits or hospitalizations in the past 5 years, and those who stated the reason as type 2 diabetes were selected. In the second stage of sampling, a research cooperation request form was provided and participants were asked whether they agreed to participate in the study; those who agreed were designated as research participants.

G*Power 3.1 was used to estimate the minimum sample size. To separately analyze participants with high and low Summary of Diabetes Self-Care Activities (SDSCA) scores (SDSCA score difference, 3.0; standard deviation, 6.0; significance level, 0.05; power, 0.95), the required minimum sample size was 210.

Possible Biases

It is crucial that people with diabetes be sampled without age or gender bias. Therefore, a research firm was selected based on its track record of academic research with people with diabetes. Over 15,000 individuals with diabetes have participated in previous surveys. The average age of respondents to previous survey was close to the average age of people with diabetes in Japan reported in 2019. Because of concerns about the low number of female respondents in past surveys, a lower limit was set in advance for the number of male and female respondents.

Second, a selection bias was considered, in which responses are concentrated among respondents who are digitally literate. Since this bias could not be completely avoided, we decided to establish a “do not have a smartphone” question to see if the responses were not excessively biased toward smartphone owners.

Ethical Approval and Consent to Participate

All participants provided written informed consent before the initiation of the study. The study protocol was reviewed and approved by the Institutional Review Board of Kobe City College of Nursing (20122-05).

Questionnaire Development and Variables

This cross-sectional survey was conducted using an Internet-based questionnaire. The entire survey was conducted on the research firm's Web page. Respondents logged into the Web page to complete the survey. As it is an Internet-based survey, the questionnaire was developed and designed by referring to the “Checklist for Reporting Results of Internet E-Surveys.”²⁵

Health Care App Use

The participants were asked regarding the use of health care- and fitness-related apps on their smartphones. The following answer choices were available: “currently using,” “have used in the past,” “have never used,” and “do not have a smartphone.” Additionally, the respondents were asked to provide the characteristics of the apps they use. The following options

were provided for app characteristics: “general condition (body temperature/sleep/general condition),” “weight management,” “physical activity (pedometer/activity meter),” “dietary management,” and “other type.” In this study, participants who answered “currently using” were defined as people with diabetes using health care apps.

Self-Management Measurement

The SDSCA-Revised is a scale that includes commonly recommended measures of diabetes self-management behaviors.²⁶ Japanese version of the SDSCA (J-SDSCA) questionnaire was applied,²⁷ and among various items in this questionnaire, “general diet” and “exercise” were used in this study. Higher scores on these items indicated better self-management behavior.

Other Independent Variables

Data on sociodemographic variables (e.g., age, sex, and family size) that influence diabetes self-management behaviors were collected.^{28,29} Moreover, motivational questions regarding treatment behaviors were included in the survey. Autonomous motivation is an important concept as it is associated with achieving and maintaining diabetes-related self-care goals.^{30–34} The Treatment Self-Regulation Questionnaire (Controlled and Autonomous motivation) was used to measure motivation.³⁵

Statistical Analysis

Data obtained from the entire questionnaire were coded and entered using Microsoft Excel. The data were then analyzed using SPSS version 26. Frequency and cross-tabulations were used to describe the data. To analyze the association between health care app use and diabetes self-management behaviors, multiple regression analysis was conducted using the J-SDSCA and subscales of general diet and exercise as objective variables. In this study, age, sex, and family size were set as covariates, and the health care app use and motivation scale were considered independent predictors. The unstandardized and standardized partial regression coefficients and 95% confidence intervals are presented. The Durbin–Watson (DW) test was used to confirm that the model was independent of multicollinearity and to demonstrate its adequacy, with a DW value of 1.5 to 2.5 as the preferred range.³⁶ Statistical significance was set at a *p*-value of < 0.05.

Results

Participant Characteristics and Health Care App Use on Smartphones

Overall, 466 people with diabetes participated in the screening survey, and 253 participants completed the survey. ▶ **Table 1** shows the demographic characteristics of the 253 participants. There were no incomplete data, and all

Table 1 Summary of the sociodemographic statistics of the survey population (*N* = 253)

Demographic details	Frequency (%)	Mean ± SD
Sex		
Female	103 (40.7)	
Male	150 (59.3)	
Age (y)		63.7 ± 10.1
< 60	86 (34.0)	
60–69	83 (32.8)	
> 70	84 (33.2)	
Family member		
Living alone	45 (17.8)	
Living with family members	208 (82.2)	
Use of health care apps on smartphone		
Currently using	61 (24.1)	
Have used in the past	18 (7.1)	
Have never used	129 (51)	
Do not have a smartphone	45 (17.8)	
Type of health care apps participants have used		
General condition apps (body temperature, sleep monitor)	26 (10.3)	
Weight management apps	31 (12.3)	
Physical activity apps (pedometer, activity monitor)	67 (26.5)	
Diet management apps	16 (6.3)	
Other apps	8 (3.2)	

Abbreviation: SD, standard deviation.

Table 2 Multiple regression of factors contributing to physical activity habits^a

Variables	Unstandardized Beta	SE	Standardized Beta	Significance	95% CI	
(Constant)	−3.92	0.96		0.00	−5.81	−2.03
Age	0.03	0.01	0.12	0.05	0.00	0.05
Sex (dummy ^b)	0.48	0.27	0.10	0.08	−0.05	1.00
Living alone (dummy ^c)	0.16	0.34	0.03	0.65	−0.52	0.83
Autonomous motivation	0.58	0.11	0.34	0.00	0.36	0.80
Controlled motivation	0.17	0.12	0.09	0.15	−0.06	0.40
Current use of apps <any health care apps>	0.91	0.32	0.17	0.00	0.29	1.53

Abbreviations: CI, confidence interval; J-SDSCA, the Japanese version of the Summary of Diabetes Self-Care Activities; SE, standard error.

^aThe dependent variable is exercise score on J-SDSCA.

^bSex is a dummy variable (female = 1, male = 2).

^cLiving alone is a dummy variable (living alone = 0, with family members = 1).

data were eligible for the final analysis. Thus, there was no handling of missing data.

Of the 253 participants, 24.1% (61/253) reported that they were using health care apps at the time. Moreover, 31.6% (25/79) of those aged < 60 years, 20.5% (17/83) of those aged 60 to 69 years, and 19.0% (16/84) of those aged > 70 years used health care apps. A physical activity app was the most used type of app, which was used by 26.5% (67/253) of the respondents.

Determinants of Physical Activity Habits

Multiple linear regression analysis was conducted to predict the frequency of individuals' physical activity based on their use of health care apps. A significant regression equation was obtained ($F(6,246) = 14.30$; $p < 0.001$), with an R^2 value of 0.24. The independent predictors that were significantly associated with this model were autonomous motivation and health care app use. Participants who used health care apps showed a higher physical activity score by 0.91 points

than those who did not (see ►Table 2 for all regression values). The DW statistic value in this model was 2.07.

Determinants of Dietary Habits

Multiple linear regression analysis was conducted to predict the general diet score on the use of health care apps. A significant regression equation was obtained ($F(6,246) = 9.82$; $p < 0.001$), with an R^2 value of 0.17. The use of health care apps was not shown to be significantly associated with dietary habits in this model. Autonomous motivation was the only independent predictor that was significantly associated with this model (see ►Table 3 for all regression values). The DW statistic value in this model was 1.92.

Discussion

Health Care App Use by People with Type 2 Diabetes

Of the total participants in this study, 24.1% used health care apps. In the 2018 survey, 33.3% of people with type 2 diabetes

Table 3 Multiple regression of factors contributing to dietary habits^a

Variables	Unstandardized Beta	SE	Standardized Beta	Significance	95% CI	
(Constant)	−0.39	1.16		0.74	−2.69	1.90
Age	0.00	0.02	−0.01	0.89	−0.03	0.03
Sex (dummy ^b)	−0.14	0.32	−0.02	0.67	−0.78	0.50
Living alone (dummy ^c)	0.37	0.42	0.05	0.38	−0.46	1.19
Autonomous motivation	0.77	0.14	0.39	0.00	0.50	1.03
Controlled motivation	0.09	0.14	0.04	0.55	−0.20	0.37
Current use of apps <any health care apps>	0.40	0.38	0.06	0.29	−0.35	1.16

Abbreviations: CI, confidence interval; J-SDSCA, the Japanese version of the Summary of Diabetes Self-Care Activities; SE, standard error.

^aThe dependent variable is general diet score on J-SDSCA.

^bSex is dummy variable (female = 1, male = 2).

^cLiving alone is dummy variable (living alone = 0, with family members = 1).

used health care apps for diabetes management compared with past surveys.³⁷ The mean age of participants in this study was 63.7 ± 10.1 years, which was more than that in previous studies; this could have influenced the slightly lower rate of app use. Several previous studies have noted design and usability issues specific to older adults and various psychological barriers.³⁸ However, this survey found that approximately 20% of respondents over the age of 60 years reported using health care apps to manage their health, indicating that even among the elderly, about one-fifth of the elderly use health care apps on a daily basis. This may be due to the widespread use of smartphones, which reduces barriers to health care management among older adults using health care apps. In fact, based on the penetration rates alone, smartphone use for health care purposes in people with diabetes was more prevalent in the present study than in a 2014 Latino survey in which only 3.3% of people with diabetes used health care apps.³⁹

Second, in terms of the type of app, the most common type of app used was associated with exercise management. However, previous studies have described effective and frequently used features of health care apps, such as food intake recording,³⁷ medication management,⁴⁰ and blood glucose management.⁴¹ According to a series of studies on the relationship between health care app and diabetes management, regardless of what is managed on their smartphone, there would be a factor that promote diabetes self-care in use of smartphones for health.

Third, 7% of individuals indicated that they have used health-related apps in the past. Issues related to usability were noted as a reason for lack of continued use.^{20,21,42} In addition, a previous survey in Australia reported that over 40% of respondents believed that smartphone-based management would not help them.⁴³ A past study had shown that although people with type 2 diabetes favored the features of health care apps and expressed interest in mobile health, their use of the apps was less than 10%.⁴⁴ These suggest that user engagement needs to be strengthened. To address these issues, a regular survey of easy-to-use yet effective diabetes support apps that clinicians can refer to may be effective. The Association of Diabetes Care & Education Specialists in the United States recommends diabetes support apps and regularly reviews the accuracy and relevance of the information,⁴⁵ and it may be helpful to refer to these activities in each language area.

Relationship between Health Care App Use and Exercise Habits

The participants who used health care apps showed a higher physical activity score by 0.91 points than those who did not. Thus, a difference of approximately 1 day in physical activity frequency was noted between the two groups, indicating a clear relationship between health care app use and physical activity frequency. This result is consistent with those of previous studies, suggesting that health care app use stimulates diabetes self-management behaviors along with individual self-efficacy and autonomous motivation.³²

Other studies have reported that only 9% of people with diabetes habitually engage in effective exercise.⁴⁶ A sedentary lifestyle, such as lack of exercise habits, has been

an independent risk factor for mortality^{47,48}; therefore, approaches to encourage physical activity are of growing interest among clinicians involved in diabetes management.⁴⁹ The results of this study indicate that health care apps can potentially be used to help people with type 2 diabetes acquire or maintain exercise habits.

Relationship between Health Care App Use and Healthy Eating Habits

Several studies have reported the association between dietary habits and health care app use^{37,50}; however, the results of this study showed no association between them. Multiple regression analysis revealed that the only independent predictor affecting dietary self-management was autonomous motivation. Another previous study reported that autonomous motivation was the only variable that improved dietary activity. Further, it reported that controlling motivation, self-efficacy, social support, and health literacy were not associated with dietary activity.³³ As a healthy diet requires several steps, such as acquiring the right knowledge, shifting the diet, and introducing a new food culture,^{34,51,52} using a health care app alone may not lead to the development of healthy eating habits. The result of lack of significant differences in the dietary scores using health care apps is consistent with those of previous intervention studies in which no change was reported in the dietary scores,^{16,53} thus reaffirming the difficulty in improving dietary behavior. The results of this study suggest that autonomous motivation plays a role in their eating habits, it is therefore important to help people with diabetes find meaning for themselves in healthy eating by encouraging reflection.⁵⁴ It can be stated that autonomous motivation requires several months of continuous positive coaching by the clinician.⁵⁵ In the future, it is necessary to confirm whether the use of smartphone interventions facilitates ongoing coaching and test whether they contribute to changes in dietary behavior.

Limitations

This study has several limitations. The survey was conducted among people with diabetes from a research firm through the Internet, which could have resulted in selection bias. However, the age range, sex ratio,²³ smartphone ownership,⁵⁶ and health care app use³⁷ of the respondents in this study approximated the data reported in other studies. These results suggest that the demographics of the respondents adequately reflect the population of people with type 2 diabetes in Japan. However, the potential lack of generalizability is a concern because the target population of this study included preregistered survey participants from a research firm.⁵⁷

Another important limitation is that although we examined app use, we did not assess the technique of using the apps and the duration for which participants used the apps (frequency and intensity). Furthermore, we cannot rule out the possibility that factors related to the socioeconomic status, such as education and economic status, may influence self-management. A resurvey using these parameters might be useful for refining and clarifying the factors that contribute to the variations in the diabetes self-management status.

Future studies are needed to determine whether the effects of using health care apps are sustainable and effective in preventing diabetes complications. Medical outcomes, such as hemoglobin A1c level, may be used in such studies to demonstrate the effects of health care apps in further detail.

Conclusion

This study reported an association between self-management behaviors and health care app use in people with diabetes. Approximately 25% of people with diabetes used health care apps, and those who used health care apps showed significantly higher exercise self-management scores than those who did not.

Clinical Relevance Statement

Exercise self-management scores were significantly higher among those using health care apps than those not using such apps. Therefore, including recommendations for using health care apps in the DSMS may be effective in acquiring exercise habits. The fact that one in four people with diabetes used the app in this study could be encouraging for potential health care app users.

Protection of Human and Animal Subjects

This study was conducted in compliance with the World Medical Association Declaration of Helsinki on ethical principles for medical research involving human subjects, and it was reviewed by the Institutional Review Board of Kobe City College of Nursing.

Ethical Approval

All participants provided written informed consent before initiation of the study. The study protocol was reviewed and approved by the Institutional Review Board of Kobe City College of Nursing (20122-05).

Data Sharing Policy

Data are available upon reasonable request. The data that support the findings of this study are available from the corresponding author upon reasonable request. The approval of the Institutional Review Board of Kobe City College of Nursing may be required to determine whether the data acquisition is reasonable.

Funding

This work was supported by JSPS KAKENHI Grant-in-Aid for Young Scientists 19K19533. The funding agreement ensured the authors' independence in designing the study, interpreting the data, writing, and publishing the report.

Conflict of Interest

T.M. received a speaker's honorarium from Eli Lilly Japan K.K. The other authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Acknowledgment

The authors would like to express their gratitude to the study participants.

References

- 1 International Diabetes Federation. IDF diabetes atlas - 9th edition 2019. 2020. Accessed August 18, 2021 at: <https://www.diabetesatlas.org/en/>
- 2 Dal Canto E, Ceriello A, Rydén L, et al. Diabetes as a cardiovascular risk factor: an overview of global trends of macro and micro vascular complications. *Eur J Prev Cardiol* 2019;26(2_suppl):25–32
- 3 Funnell MM, Tang TS, Anderson RM. From DSME to DSMS: developing empowerment-based diabetes self-management support. *Diabetes Spectr* 2007;20:221–226
- 4 Beck J, Greenwood DA, Blanton L, et al. 2017 National Standards for Diabetes Self-Management Education and Support. *Sci Diabetes Self Manag Care* 2021;47(01):14–29
- 5 Haas L, Maryniuk M, Beck J, et al; 2012 Standards Revision Task Force. National standards for diabetes self-management education and support. *Diabetes Care* 2012;35(11):2393–2401
- 6 Norris SL, Chowdhury FM, Van Le K, et al. Effectiveness of community health workers in the care of persons with diabetes. *Diabet Med* 2006;23(05):544–556
- 7 Powers MA, Bardsley J, Cypress M, et al. Diabetes self-management education and support in type 2 diabetes: a joint position statement of the American Diabetes Association, the American Association of Diabetes Educators, and the Academy of Nutrition and Dietetics. *Diabetes Care* 2015;38(07):1372–1382
- 8 He Q, Zhao X, Wang Y, Xie Q, Cheng L. Effectiveness of smartphone application-based self-management interventions in patients with type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *J Adv Nurs* 2022;78(02):348–362
- 9 Hamine S, Gerth-Guyette E, Faulx D, Green BB, Ginsburg AS. Impact of mHealth chronic disease management on treatment adherence and patient outcomes: a systematic review. *J Med Internet Res* 2015;17(02):e52
- 10 Wang Y, Xue H, Huang Y, Huang L, Zhang D. A systematic review of application and effectiveness of mHealth interventions for obesity and diabetes treatment and self-management. *Adv Nutr* 2017;8(03):449–462
- 11 Anderson K, Burford O, Emmerton L. Mobile health apps to facilitate self-care: a qualitative study of user experiences. *PLoS One* 2016;11(05):e0156164
- 12 Zhao J, Freeman B, Li M. Can mobile phone apps influence people's health behavior change? An evidence review. *J Med Internet Res* 2016;18(11):e287
- 13 Xie Z, Nacioglu A, Or C. Prevalence, demographic correlates, and perceived impacts of mobile health app use amongst Chinese adults: cross-sectional survey study. *JMIR Mhealth Uhealth* 2018;6(04):e103
- 14 Fleming GA, Petrie JR, Bergenstal RM, Holl RW, Peters AL, Heine mann L. Diabetes digital app technology: benefits, challenges, and recommendations. A consensus report by the European Association for the Study of Diabetes (EASD) and the American Diabetes Association (ADA) Diabetes Technology Working Group. *Diabetes Care* 2020;43(01):250–260
- 15 Doyle-Delgado K, Chamberlain JJ. Use of diabetes-related applications and digital health tools by people with diabetes and their health care providers. *Clin Diabetes* 2020;38(05):449–461
- 16 Lee DY, Park J, Choi D, Ahn HY, Park SW, Park CY. The effectiveness, reproducibility, and durability of tailored mobile coaching on diabetes management in policyholders: a randomized, controlled, open-label study. *Sci Rep* 2018;8(01):3642
- 17 Kim YJ, Rhee SY, Byun JK, et al. A smartphone application significantly improved diabetes self-care activities with high user satisfaction. *Diabetes Metab J* 2015;39(03):207–217

- 18 Gimbel RW, Rennert LM, Crawford P, et al. Enhancing patient activation and self-management activities in patients with type 2 diabetes using the US Department of Defense Mobile Health Care Environment: feasibility study. *J Med Internet Res* 2020;22(05):e17968
- 19 Greenwood DA, Blozis SA, Young HM, Nesbitt TS, Quinn CC. Overcoming clinical inertia: a randomized clinical trial of a telehealth remote monitoring intervention using paired glucose testing in adults with type 2 diabetes. *J Med Internet Res* 2015;17(07):e178–e178
- 20 Arnhold M, Quade M, Kirch W. Mobile applications for diabetics: a systematic review and expert-based usability evaluation considering the special requirements of diabetes patients age 50 years or older. *J Med Internet Res* 2014;16(04):e104
- 21 Harrington L, Parker C, Ulanday K, Harrington C. Heuristic evaluation of a top-rated diabetes self-management app. *Appl Clin Inform* 2021;12(05):1014–1020
- 22 Tran J, Tran R, White JR Jr. Smartphone-based glucose monitors and applications in the management of diabetes: an overview of 10 salient “apps” and a novel smartphone-connected blood glucose monitor. *Clin Diabetes* 2012;30:173–178
- 23 Japan Diabetes Clinical Data Management Study Group. Basic study material 2019. 2020. Accessed August 18, 2021 at: <https://jddm.jp/data/index-2019/>
- 24 Araki E, Goto A, Kondo T, et al. Japanese Clinical Practice Guideline for Diabetes 2019. *J Diabetes Investig* 2020;11(04):1020–1076
- 25 Eysenbach G. Improving the quality of Web surveys: the Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2004;6(03):e34–e34
- 26 Toobert DJ, Hampson SE, Glasgow RE. The Summary of Diabetes Self-Care Activities measure: results from 7 studies and a revised scale. *Diabetes Care* 2000;23(07):943–950
- 27 Daitoku M, Honda I, Okumiya A, et al. Validity and reliability of the Japanese translated “the Summary of Diabetes Self-care activities measure”. *J Japan Diabetes Soc* 2006;49:1–9
- 28 Miller TA, Dimatteo MR. Importance of family/social support and impact on adherence to diabetic therapy. *Diabetes Metab Syndr Obes* 2013;6:421–426
- 29 Caruso R, Rebora P, Luciani M, Di Mauro S, Ausili D. Sex-related differences in self-care behaviors of adults with type 2 diabetes mellitus. *Endocrine* 2020;67(02):354–362
- 30 Ryan RM, Deci EL. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am Psychol* 2000;55(01):68–78
- 31 Williams GC, McGregor HA, Zeldman A, Freedman ZR, Deci EL. Testing a self-determination theory process model for promoting glycemic control through diabetes self-management. *Health Psychol* 2004;23(01):58–66
- 32 Grønnegaard C, Varming A, Skinner T, Olesen K, Willaing I. Determinants of glycaemic control among patients with type 2 diabetes: testing a process model based on self-determination theory. *Heliyon* 2020;6(10):e04993
- 33 Shigaki C, Kruse RL, Mehr D, et al. Motivation and diabetes self-management. *Chronic Illn* 2010;6(03):202–214
- 34 De Man J, Wouters E, Delobelle P, et al. Testing a self-determination theory model of healthy eating in a South African township. *Front Psychol* 2020;11:2181
- 35 Inagaki S, Matsuda T, Abe K, et al. Development and validation of the Treatment Self-Regulation Questionnaire for Diabetics in Japanese (TSRQ-DJ). *J Japan Diabetes Soc* 2022;65:26–33
- 36 Backhaus K, Erichson B, Gensler S, et al. *Multivariate Analysis: An Application-Oriented Introduction*. Wiesbaden: Springer Gabler; 2021
- 37 Kebede MM, Pischke CR. Popular diabetes apps and the impact of diabetes app use on self-care behaviour: a survey among the digital community of persons with diabetes on social media. *Front Endocrinol (Lausanne)* 2019;10:135
- 38 Joe J, Demiris G. Older adults and mobile phones for health: a review. *J Biomed Inform* 2013;46(05):947–954
- 39 Williams JP, Schroeder D. Popular glucose tracking apps and use of mHealth by Latinos with diabetes: review. *JMIR Mhealth Uhealth* 2015;3(03):e84
- 40 Lim S, Kang SM, Shin H, et al. Improved glycemic control without hypoglycemia in elderly diabetic patients using the ubiquitous healthcare service, a new medical information system. *Diabetes Care* 2011;34(02):308–313
- 41 Boyle L, Grainger R, Hall RM, Krebs JD. Use of and beliefs about mobile phone apps for diabetes self-management: surveys of people in a hospital diabetes clinic and diabetes health professionals in New Zealand. *JMIR Mhealth Uhealth* 2017;5(06):e85
- 42 Scheibe M, Reichelt J, Bellmann M, et al. Acceptance factors of mobile apps for diabetes by patients aged 50 or older: a qualitative study. *Med* 2015;4:e1
- 43 Trawley S, Baptista S, Browne JL, Pouwer F, Speight J. The use of mobile applications among adults with type 1 and type 2 diabetes: results from the Second MILES-Australia (MILES-2) study. *Diabetes Technol Ther* 2017;19(12):730–738
- 44 Conway N, Campbell I, Forbes P, Cunningham S, Wake D. mHealth applications for diabetes: user preference and implications for app development. *Health Informatics J* 2016;22(04):1111–1120
- 45 Ministry of Internal Affairs and Communications. Outline of the 2020 White Paper on Information and Communications in Japan 2020. Accessed August 18, 2021 at: <https://www.soumu.go.jp/johotsusintokei/whitepaper/eng/WP2020/2020-index.html>
- 46 Thomas N, Alder E, Leese GP. Barriers to physical activity in patients with diabetes. *Postgrad Med J* 2004;80(943):287–291
- 47 Ekelund U, Steene-Johannessen J, Brown WJ, et al; Lancet Physical Activity Series 2 Executive Committee; Lancet Sedentary Behaviour Working Group. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* 2016;388100511302–1310
- 48 Wilmot EG, Edwardson CL, Achana FA, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia* 2012;55(11):2895–2905
- 49 Hamasaki H. Daily physical activity and type 2 diabetes: a review. *World J Diabetes* 2016;7(12):243–251
- 50 Ridat GS, Maybituin VCS, Bella CY, et al. Project DiabEHT: an approach to improve self-care management of diabetes. *Enferm Clin* 2020;30(Suppl 5):234–239
- 51 Fukuoka Y, Lindgren TG, Bonnet K, Kamitani E. Perception and sense of control over eating behaviors among a diverse sample of adults at risk for type 2 diabetes. *Diabetes Educ* 2014;40(03):308–318
- 52 Kasila K, Poskiparta M, Karhila P, Kettunen T. Patients' readiness for dietary change at the beginning of counselling: a transtheoretical model-based assessment. *J Hum Nutr Diet* 2003;16(03):159–166
- 53 Lee M-K, Lee DY, Ahn H-Y, Park CY. A novel user utility score for diabetes management using tailored mobile coaching: secondary analysis of a randomized controlled trial. *JMIR Mhealth Uhealth* 2021;9(02):e17573
- 54 Zoffmann V, Kirkevold M. Realizing empowerment in difficult diabetes care: a guided self-determination intervention. *Qual Health Res* 2012;22(01):103–118
- 55 Sebire SJ, Toumpakari Z, Turner KM, et al. “I’ve made this my lifestyle now”: a prospective qualitative study of motivation for lifestyle change among people with newly diagnosed type two diabetes mellitus. *BMC Public Health* 2018;18(01):204
- 56 Association of Diabetes Care and Education Specialists. DANA App Review for AADE members 2018. Accessed August 19, 2021 at: <https://www.danatech.org>
- 57 Remillard ML, Mazor KM, Cutrona SL, Gurwitz JH, Tjia J. Systematic review of the use of online questionnaires of older adults. *J Am Geriatr Soc* 2014;62(04):696–705