



The Effect of Correlation of Laboratory-Developed Test and Initial Symptoms and False Negatives in RT-PCR Strategies for COVID-19 Patients with Beta Variants

Shohreh Ghasemi^{1,2} Seyed Alireza Nadji³ Mahmood Dashti⁴ Mahboobeh Karimi-Galougahi³
Negar Raygani⁵ Mahla Nabi⁶ Ghazal Mohammadi⁷ Niyosha Kandez⁸ Amirali Ebrahimi⁹
Elaheh Askari¹⁰

¹OMFS Department of Augusta University, Georgia, United States

²Master of Craniofacial Reconstruction and Trauma Queen Mary, University of London, London, Great Britain

³Virology Research Center, National Research Institute of Tuberculosis and Lung Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁴Georgia School of Orthodontics, Atlanta, Georgia, United States

⁵Department of Otolaryngology, Masih Daneshvari Hospital, Tehran, Iran

⁶Department of Radiology, Ziyaian Hospital, Tehran University of Medical Sciences, Tehran, Iran

⁷Cancer Institute of Imam Khomeini, Faculty of Medicine, Tehran University of Medical Science, Tehran, Iran

⁸Psychology Department, University of South Alabama, Mobile, Alabama, United States

⁹Masih Daneshvari Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

¹⁰School of Dentistry, University of Pennsylvania, Philadelphia, Pennsylvania, United States

Address for correspondence Mahmood Dashti, DDS, Georgia School of Orthodontics, 1120 15th St, Augusta, Georgia 30912, United States (e-mail: dashti.mahmood72@gmail.com).

Eur Dent Res Biomater J 2022;3:21–25.

Abstract

Keywords

- ▶ SARS-CoV-2
- ▶ COVID-19
- ▶ saliva
- ▶ RT-PCR test
- ▶ laboratory-developed test

Objective Reverse transcription-polymerase chain reaction (RT-PCR) assays detect severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The number of viruses in the sample varies between patients; it depends on sample location, nasal or throat, and with time infection spreads. Previous studies showed that the viral load of coronavirus disease 2019 (COVID-19) infection is the peak just before symptoms onset. Furthermore, positive and negative results depend on test site, sampling, and timing method; RT-PCR can be 1 to 30% false-negative result.

Materials and Methods Within this study, we took RT-PCR test from COVID-19 positive patients who already had the confirmation of the disease either by lung

Introduction

Since the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an RNA virus, it is very mutable, so numerous variants of SARS-CoV-2 are spreading worldwide.

To prevent the spread of disease, countries all over the world have adopted social-distancing policies. This has limited the movement of people, disrupted their daily activities, and instituted work-from-home strategy for all employment sectors. The coronavirus disease 2019 (COVID-19) climate

article published online
April 21, 2023

DOI <https://doi.org/10.1055/s-0043-1768174>.
ISSN 2791-7452.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

computed tomography (CT)-scan or the symptoms such as dyspnea. The study was explained to all the patients, and they confirmed to take the RT-PCR test. Negative samples from those patients were retested, and if the result came back negative, we included them as negative in the result.

Result A total number of 49 patients (25 females) and (24 males) with a mean age of 53.24 years (ranging from 32 to 77) were enrolled. About 32.3% of patients, despite having COVID-19 disease, had a negative RT-PCR test. There is a positive and significant relationship between weight ($r = 0.253$) and CT at the time of hospitalization of COVID-19 patients and a negative and significant relationship with O₂ saturation without oxygen therapy ($r = -0.296$), the model can predict 67.7% of the disease due to the beta value, and the share of O₂ saturation without oxygen therapy is more than weight.

Conclusion We show that a pragmatic model can be designed to predict which patients have a higher chance of getting false-negative result, and should be retested for COVID-19. Among the variables, weight had a negative and significant relationship, and O₂ saturation without respiratory support had a negative and significant relationship with COVID-19 disease.

has devastatingly impacted global education as well.¹ In COVID-19, patient's lung is the first organ that is infected first and then it spreads in many organs such as the heart, liver, and kidneys. Before spreading viral pneumonia, SARS-CoV-2 can be measured in saliva, blood, sputum, and urine even though high viral loads of SARS-CoV-2 RNA are being found in salivary gland and saliva, potentially illustrating the importance of this biofluid for testing the disease in asymptomatic condition.² If the immune system can defend against COVID-19 at first, pneumonia will not be developed.

Although there are no specific symptoms in COVID-19 patients, there has been frequent symptoms such as fever, cough, diarrhea, and fatigue,^{3,4} and diagnosis of COVID-19 relies on the detection of viral ribonucleic acid (RNA) sequences by using reverse transcription-polymerase chain reaction (RT-PCR).

Sufficient viruses on the test site, sampling methods, and timing can result in a positive test. Previous research showed that in COVID-19 infection, viral loads rise just before onset of symptoms and at symptom; furthermore, false-negative results from respiratory samples for SARS-CoV-2 ranged from 1 to 30%.

The World Health Organization suggests that COVID-19 patients who have long symptoms with negative RT-PCR test, should undergo repeat testing (including the sampling of the lower respiratory tract) with continued infection prevention measures.^{5,6}

Even though RT-PCR has the outcome of false-negative results,⁷ it is still the most conventional method for testing because saliva collection is quite comfortable for patients as well as being easy, cheap, and noninvasive with minimal equipment required. It should also minimize the nosocomial transmission of 2019-nCoV to healthcare workers⁸ and saliva has shown true potential as an ideal noninvasive diagnostic specimen, with a high degree of sensitivity and specificity for the detection of the SARS-COV-2 virus.⁹ This study aimed to determine the relationship between the false-negative result

and the laboratory-developed test and initial symptoms in COVID-19 patients and their individual and additive power to predict in-hospital patients.^{10,11}

Materials and Methods

This Ethics Committee-approved prospective observational study includes the Masih Daneshvari Hospital. During the COVID-19 pandemic peak, this center was the first COVID-19 dedicated hospital in Iran.

Trial Registration

This research was supported by the Masih Daneshvari Hospital and Shahid Beheshti Medical Sciences, and the Shahid Beheshti Medical Sciences approved the protocol of this study with ethical code IR.SMBU.NRITLTD.1400.008.

Study Population

The study was conducted on 49 patients aged between 32 and 77 years old with an average of 53.24 years old referred to Masih Daneshvari Hospital. The illness of these patients was confirmed according to the symptoms such as dyspnea, diagnostic criteria, including laboratory tests (complete blood count, erythrocyte sedimentation rate, C-reactive protein, D-dimer, ...), radiopacity in radiographs, and computed tomography (CT) scans detection of the lungs. The study was explained to all the patients, and they agreed to take the RT-PCR test, both nasopharyngeal and oropharyngeal swabs. We repeated the test once again for negative tests, and if the result came back negative, we included them as negative in the result. Of 49 patients, 32.3% with COVID-19 had negative RT-PCR.

Exclusion Criteria of the Study Population

Patients were admitted to the intensive care unit with shortness of breath (>30 breaths/min), oxygen saturation of 90 at rest with

partial oxygen pressure, and nasal oxygenation (5–6 L/min) per fraction of inspired oxygen (FiO₂) less than 300 mm Hg. Patients with weakened immune systems, cough, patients with sepsis (Sequential Organ Failure Assessment (SOFA) score 2 or higher), shortness of breath, significant comorbidities, and fever (obstructive pulmonary disease, chronic kidney disease, chronic COVID-19 infection, diabetes, and congestive heart failure) were also admitted to the intensive care unit. The presence of hypoxemia, hypercapnic acidosis, despite the introduction of nasal oxygen at a high flow rate (oxygen flow rate \geq 40%, and FiO₂ \geq 60%) or severe shortness of breath with increased work (frequency) of breathing are all indicators of mechanical ventilation (a set of assistive products, intercostal recession or nasal enlargement and set of expiratory muscles). Patients with adult respiratory distress syndrome requiring mechanical respiration were assigned a lung-protective ventilation mode.

PCR Analysis and PCR Ct Values

Viral RNA extraction was performed using a high pure viral nucleic acid kit (Roche, Switzerland) following the manufacturer's protocol. PCR was performed using the extracted nucleic acid and a real-time COVID-19 commercial kit (Pish-taz, Iran). Briefly, a reaction included 15 mL of 9 U of the enzyme, 1 mL of primer & probe mix (RdRp/N/IC), 5 mL water, and 5 mL of extracted RNA. Then, PCR was carried out with a reverse transcription step at 50 °C for 20 minutes, cDNA initial denaturation at 95 °C for 3 minutes, and finally 45 cycles of 10 seconds at 94 °C and 40 seconds at 55 °C. Light Cycler 96 (LC96) PCR machine (Roche, Germany) was used to perform the amplification and to evaluate Ct (cycle threshold) values. Ct values less than or equal to 29 were considered strong positive reactions, Ct values of 30 to 37 indicated positive reactions, and Ct values of 38 to 40 represented weak reactions indicative of minimal amounts of target nucleic acid.

Result

Descriptive statistics used statistical indicators such as frequency, mean, and standard deviation. In inferential statistics, the default normality of the Kolmogorov–Smirnov test was checked and confirmed due to the need to use parametric tests to analyze the data. Then, the Pearson correlation test was used to examine the relationship between the variables, and multiple linear regression was used to predict after the Durbin Watson test and data alignment through SPSS software, version 25. Initially, the normality of the data was confirmed by the Kolmogorov–Smirnov test, and the conditions of Pearson correlation analysis were observed. Findings show that out of 49 patients, 32.3%, despite having COVID-19 disease, had a negative RT-PCR test; their age was between 32 and 77 years with an average of 53.24. Out of 49 patients 25 patients were male and 24 female. The results as shown in ►Table 1 are descriptive statistics from the center-orientation index and the dispersion index, which include the mean and standard deviation of research variables. Also, in inferential statistics, there has been a positive and significant relationship between weight ($r=0.253$) and CT at the time of hospitalization of COVID-19 patients and a negative and significant relationship between O₂ saturation without oxygen therapy ($r=-0.296$) and CT at the time of hospitalization of COVID-19 disease as ►Table 2 shows the results of the study showed multiple regression assumptions.

According to the value obtained from the multiple correlation coefficient, the model can predict 67.7% of the disease. Among the variables, weight had a negative and significant relationship, and O₂ saturation without respiratory support had a negative and significant relationship with COVID-19 disease, and due to the beta value, the share of O₂ saturation without respiratory support is more than weight.

Table 1 Mean, standard deviation, and correlation coefficients of research variables in patients with COVID-19 ($*p < 0.05$)

Variable	Half-value	ST	CT upon arrival	Age	Weight	Onset of symptoms	LDH	CRP	LYM	O ₂ Sat without oxygen therapy
CT upon arrival	32.31	0.89	1							
Age	53.24	1.74	-0.093	1						
Weight	83.02	1.95	0.253*	-0.16	1					
Onset of symptoms	9.08	0.51	-0.128	0.086	0.253	1				
LDH	608.06	25.89	0.150	0.033	0.198	-0.098	1			
CRP	36.45	2.31	0.106					1		
LYM	17.28	1.33	0.022	-0.053	-0.022	0.054	-0.326	-0.324	1	
O ₂ without oxygen	94.41	0.48	-0.296*	-0.078	-0.039	0.225	-0.272	-0.078	0.092	1

Abbreviations: COVID-19, coronavirus disease 2019; CRP, C-reactive protein; CT, computed tomography; LDH, lactate dehydrogenase; LYM,—.

Table 2 Multiple correlation coefficient, beta coefficients, determination coefficient, and standard estimation error (* $p < 0.05$)

Predictive variables	r		F	Df	SE	p -Value	Impact coefficient	p -Value
Gender	0.527	0.677	1.66	8	5.92	0.031	1.17	0.737
Age							-0.137	0.102
Weight							-0.328	0.028*
Onset of symptoms							0.066	0.8
LDH							0.006	0.275
CRP							-0.002	0.972
LYM							0.075	0.471
O2 sat without oxygen							-0.610	0.035*

Abbreviations: CRP, C-reactive protein; LDH, lactate dehydrogenase; LYM,—; SE, standard error.

Discussion

Our study included 32.2% of patients with COVID-19 symptoms and pulmonary involvement in CT-scan, had two negative PCR tests reported, taken from both oropharynx and nasopharynx swaps. Furthermore, the variant of the virus we worked on was the B.1.351 (beta variant) species.

For nasopharyngeal swaps, positive rates have been reported moderate, while oropharyngeal swaps have shown a low positive rate due to the systematic review and meta-analysis done by Bwire et al.¹²

We considered factors such as age, sex, weight, and CT scan upon arrival, lactate dehydrogenase, C-reactive protein, Lymphocyte count, LYM, O2 sat without oxygen, and symptom onset. Based on our findings, the patient's weight was inversely related to the CT value; patients with high body mass index (BMI) had a poor prognosis.

Also, in the case of O2 saturation, higher O2 saturation, the RT-PCR test was less likely to be false negative.

The systematic review by Ingrid Arevalo-Rodriguez et al reported a false-negative rate in the RT-PCR test with an average of 0.11 rate and emphasized retesting because nearly 54% of patients whose clinical signs were suspected of having COVID-19 had a negative RT-PCR test in the first trial.¹³

There are various factors involved in the negative and positive results of the RT-PCR test. According to the systematic review by Sue Mallett, the time of onset of symptoms and the sample transport tube are the most critical factors that affect the test result.¹⁴

In analytical considerations by Rahbari et al, which examined possible errors in the detection of COVID-19, preanalytical errors were considered as the main factor in detecting coronaviruses, such as equipment, location, and sampling time.¹⁵

In a study by Zhou et al, it has been said that the use of high-sensitivity kits is effective in reducing false-negative rates. Based on this, it is recommended to evaluate the ability of detection kits before routine usage. Moreover, continuous amplification can increase the detection rate of low viral load specimens. As a result, it can significantly reduce the false-negative rate of SARS-CoV-2.¹⁶

Due to the importance of this disease, the negative RT-PCR cannot be easily ignored. According to the retrospective

cohort study by Xiao et al, in addition to CT scan, in patients with a negative RT-PCR test, it was suggested to measure the levels of CD3, CD4, CD8, CD19, immunoglobulin M, C3 complement, and C4 complement for better screening.¹⁷

In addition to RT-PCR testing, various diagnostic methods have been studied in a systematic review by Mistry et al. Variable sensitivity was also reported in their study. However, this device could be a new form of the COVID-19 test.¹⁸

A systematic review and meta-analysis by Subali and Wiyono discuss that the reverse transcriptase loop-mediated isothermal amplification is more sensitive and specific compared to RT-PCR.¹⁹ Furthermore, droplet digital PCR (ddPCR) has been studied by Suo et al and ddPCR points to the distinctive clinical detection of SARS-CoV-2 to minimize random errors when compared to RT-PCR, suggesting that it could be a powerful addition to existing standard RT-PCR.²⁰ However, the RT-PCR test is still preferable to other tests, supplemented with other methods to achieve the best and most accurate results.^{8,9,21}

Lung involvement is not necessarily correlated to the result of real-time RT-PCR; in a study, Xingzhi Xie et al found chest CT findings in negative RT-PCR. A typical diagnostic finding, ground-glass opacity, was viral pneumonia due to (COVID-19) infection (chest CT for typical COVID-19).²²

RT-PCR tests are sensitive and specific, and false-positive rarely occurs, but insufficient quantities of the virus in the sample can result in a false negative.⁷

Conclusion

Our study is one of the first studies to describe and build a model for patients who initially confirmed negative for COVID-19, but were subsequently retested and confirmed positive for COVID-19. We show that a pragmatic model can be built to predict which patients should be retested for COVID-19, and those 49 patients are sufficient number of patients to be screened. Among the variables, weight had a negative and significant relationship, and O2 saturation without respiratory support had a negative and significant relationship with COVID-19 disease. Due to the beta value, the share of O2 saturation without oxygen therapy is more than weight at a BMI of more than 23 kg/m². We found an

increase in the risk of severe COVID-19 leading to positive tests and hospitalization. More research and studies are being conducted to assess the cost-effectiveness of the retesting method in clinical practice, as well as its usefulness.

Authors' Contribution

S.G.H. helped in conceptualization, methodology, supervision, project administration, and writing—review & editing. S.A.N. was involved in conceptualization, data curation, methodology, and supervision. M.D. contributed to conceptualization, formal analysis, methodology, project administration, and writing—review & editing. M.K.G. and N.R. helped in data curation, formal analysis, methodology, and investigation. M.N. and G.h.M. were involved in investigation, validation, and writing—original draft. N.K. and E.A. helped in investigation and writing—original draft. A.E. contributed to investigation, validation, and data curation.

Availability of Data and Materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Funding

No funding was taken for this research.

Conflict of Interest

None declared.

References

- Sarwar H, Akhtar H, Naeem MM, et al. COVID-19 pandemic and challenges of dentistry: self-reported effectiveness of e-learning classes during COVID-19 pandemic: a nation-wide survey of Pakistani undergraduate dentistry students. *Eur J Dent* 2020;14(S 01):S34–S43
- Hamid H, Khurshid Z, Adanir N, Zafar MS, Zohaib S. COVID-19 pandemic and role of human saliva as a testing biofluid in point-of-care technology. *Eur J Dent* 2020;14(S 01):S123–S129
- Guan WJ, Ni ZY, Hu Y, et al; China Medical Treatment Expert Group for Covid-19. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020;382(18):1708–1720
- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020;323(11):1061–1069
- Hanson KE, Caliendo AM, Arias CA, et al. Infectious Diseases Society of America Guidelines on the Diagnosis of COVID-19: Serologic Testing. *Clin Infect Dis* 2020:ciaa1343
- Clinical, Laboratory, and Radiologic Characteristics of Patients With Initial False-Negative Severe Acute Respiratory Syndrome Coronavirus 2 Nucleic Acid Amplification Test Results | Open Forum Infectious Diseases | Oxford Academic. Accessed January 16, 2023 at: <https://academic.oup.com/ofid/article/8/1/ofaa559/5999190>
- Wikramaratna PS, Paton RS, Ghafari M, Lourenço J. Estimating the false-negative test probability of SARS-CoV-2 by RT-PCR. *Euro Surveill* 2020;25(50):2000568
- Khurshid Z, Asiri FYI, Al Wadaani H. Human saliva: non-invasive fluid for detecting novel coronavirus (2019-nCoV). *Int J Environ Res Public Health* 2020;17(07):2225
- Warsi I, Khurshid Z, Shazam H, et al. Saliva exhibits high sensitivity and specificity for the detection of SARS-CoV-2. *Diseases* 2021;9(02):38
- Abdolrahimzadeh Fard H, Mahmudi-Azer S, Sefidbakht S, et al. Evaluation of Chest CT scan as a screening and diagnostic tool in trauma patients with coronavirus disease 2019 (COVID-19): a cross-sectional study. *Emerg Med Int* 2021; 2021:4188178
- Abdolrahimzadeh Fard H, Borazjani R, Sabetian G, et al. Establishment of a novel triage system for SARS-CoV-2 among trauma victims in trauma centers with limited facilities. *Trauma Surg Acute Care Open* 2021;6(01):e000726
- Bwire GM, Majigo MV, Njiro BJ, Mawazo A. Detection profile of SARS-CoV-2 using RT-PCR in different types of clinical specimens: a systematic review and meta-analysis. *J Med Virol* 2021;93(02): 719–725
- Arevalo-Rodriguez I, Buitrago-Garcia D, Simancas-Racines D, et al. False-negative results of initial RT-PCR assays for COVID-19: a systematic review. *PLoS One* 2020;15(12):e0242958
- Mallett S, Allen AJ, Graziadio S, et al. At what times during infection is SARS-CoV-2 detectable and no longer detectable using RT-PCR-based tests? A systematic review of individual participant data. *BMC Med* 2020;18(01):346
- Rahbari R, Moradi N, Abdi M. rRT-PCR for SARS-CoV-2: analytical considerations. *Clin Chim Acta* 2021;516:1–7
- Zhou Y, Pei F, Ji M, et al. Sensitivity evaluation of 2019 novel coronavirus (SARS-CoV-2) RT-PCR detection kits and strategy to reduce false negative. *PLoS One* 2020;15(11):e0241469
- Xiao Y, Shi X, She Q, et al. Exploration of turn-positive RT-PCR results and factors related to treatment outcome in COVID-19: a retrospective cohort study. *Virulence* 2020;11(01): 1250–1256
- Mistry DA, Wang JY, Moeser ME, Starkey T, Lee LYW. A systematic review of the sensitivity and specificity of lateral flow devices in the detection of SARS-CoV-2. *BMC Infect Dis* 2021; 21(01):828
- Subali AD, Wiyono L. Reverse transcriptase loop mediated isothermal amplification (RT-LAMP) for COVID-19 diagnosis: a systematic review and meta-analysis. *Pathog Glob Health* 2021; 115(05):281–291
- Suo T, Liu X, Feng J, et al. ddPCR: a more accurate tool for SARS-CoV-2 detection in low viral load specimens. *Emerg Microbes Infect* 2020;9(01):1259–1268
- Chaimayo C, Kaewnaphan B, Tanlieng N, et al. Rapid SARS-CoV-2 antigen detection assay in comparison with real-time RT-PCR assay for laboratory diagnosis of COVID-19 in Thailand. *Virol J* 2020;17(01):177
- Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for typical coronavirus disease 2019 (COVID-19) pneumonia: relationship to negative RT-PCR testing. *Radiology* 2020;296(02):E41–E45