


A New Arthroscopic Classification of Triangular Fibrocartilage Complex Disorders

Guillaume Herzberg, MD, PhD¹  Marion Burnier, MD² Lyliane Ly, MD³
Toshiyatsu Nakamura, MD, PhD⁴ Francisco del Piñal, MD⁵ Andrea Atzei, MD⁶

¹I-Trues Surgery Unit, Orthopedic Department, Clinique Parc Lyon, Lyon, France, Clinique Val Ouest, Lyon Ecully, France

²I-Trues Surgery Unit, Department of Orthopedic, Institut Main Membre Supérieur, Villeurbanne, France

³Hospices Civils Lyon, Lyon, France

⁴Department of Orthopedic Surgery, Minato-Ku, Tokyo, Japan

⁵Serrano 58-1B 28001-Madrid, Santander, Spain

⁶Chirurgia Della Mano, Polso e Gomito, Microchirurgia Ricostruttiva MediLAB Sottoportico Teatro Dofin, Treviso, Italy

Address for correspondence Guillaume Herzberg, MD, PhD, I-Trues Surgery Unit, Orthopedic Department, 97 Quai Charles de Gaulle 69006 Lyon, France (e-mail: profgherzberg@gmail.com).

J Wrist Surg 2024;13:2–8.

Abstract

Introduction The use of wrist arthroscopy has become a prerequisite for diagnosis and treatment of triangular fibrocartilage complex (TFCC) disorders. Since Palmer's landmark paper, many new arthroscopic descriptions of TFCC tears have been published but there is no currently available updated comprehensive arthroscopic classification of TFCC lesions.

Purpose We recently described the arthroscopic anatomy of the TFCC as viewed from a 3–4 portal. Our purpose was to propose a new TFCC disorders classification based on this new arthroscopic TFCC description.

Methods We included all currently described TFCC disorders to the best of our knowledge into our arthroscopic, functional, and vascular anatomical concept. We also included patient's specific ulnar variance and distal radial ulnar joint coronal inclination as baseline treatment-oriented parameters. The fresh or chronic, repairable or non-repairable nature of some types of TFCC tears were considered as separate parameters.

Results The proposed classification includes disc “D” (degenerative or traumatic), reins “R” (traumatic), and wall “W” (traumatic) lesions. Combined lesions of those three parts of the TFCC may be easily identified. This new classification should facilitate future analysis of isolated or combined TFCC disorders whether they are degenerative and/or traumatic.

Discussion The authors present a new three-dimensional-three-part arthroscopic updated description of TFCC disorders with relevance to etiology and treatment principles.

Keywords

- ▶ wrist
- ▶ TFCC tears
- ▶ TFCC disorders
- ▶ arthroscopy
- ▶ TFCC disorder classification

The use of wrist arthroscopy has become a prerequisite for the diagnosis and treatment of any triangular fibrocartilage complex (TFCC) disorders.^{1–4} Palmer's landmark paper was the first attempt at producing a comprehensive three-

dimensional (3-D) classification of TFCC lesions including wrist arthroscopy as one of its criteria.⁵

Since Palmer's description, there have been numerous arthroscopy-based contributions to the knowledge of TFCC

received

January 26, 2023

accepted

May 9, 2023

article published online

August 7, 2023

© 2023, Thieme. All rights reserved.
Thieme Medical Publishers, Inc.,
333 Seventh Avenue, 18th Floor,
New York, NY 10001, USA

DOI <https://doi.org/10.1055/s-0043-1769908>.
ISSN 2163-3916.

lesions with many newly recognized lesions or subtypes.^{2,6-13}

However, no comprehensive update of TFCC lesions classification based on arthroscopy has been proposed so far. A recently updated classification of TFCC disorders was based on classic 3-D anatomic drawings.¹⁴ We recently proposed a new arthroscopic description of the TFCC from a 3-4 radiocarpal portal (disc, reins, and wall) including functional and vascular aspects along with a first attempt at stratifying TFCC tears.³

Here, we propose a new comprehensive arthroscopic classification of TFCC lesions and their variants based on our arthroscopic description of the TFCC as seen from a 3-4 radiocarpal wrist arthroscopy portal with the addition of distal radial ulnar joint (DRUJ) arthroscopy when necessary.

Arthroscopic Classification of Triangular Fibrocartilage Complex Disorders

This new arthroscopic classification of TFCC disorders includes in background patient's specific ulnar variance¹⁵ and Tolat's type of sigmoid notch¹⁶ because these parameters may significantly influence the treatment of TFCC disorders whatever their etiology may be (►Fig. 1). Correspondence between Palmer's classification and our new classification is shown in ►Table 1 as well as the usefulness of DRUJ arthroscopy when appropriate.

Disc (D) Triangular Fibrocartilage Complex Tears

TFCC disc "D" tears may be called disc degenerative (DD) or disc traumatic (DT) as displayed in ►Fig. 1. "DD" and "DT" tears may be included in the same group because the avascularity and lack of healing potential of the disc make their treatment similar.

- "DD" central tears (previously Palmer's Class 2 A-E) are located within the flat central TFCC disc area. Palmer Class 2 stages provided a description of the natural history of "DD" tears⁵ including the addition of regional degenerative changes to the lunate and lunotriquetral ligament tears in class 2D and 2E. We have chosen to mention "DD" TFCC tears first because they are related to patient's specific anatomical features producing a static or dynamic¹⁵ positive ulnar variance and ulnocarpal impingement syndrome.⁵ Central "DD" TFCC lesions as seen from the 3-4 portal may be simple central wears or central ovoid perforations of various sizes from localized to massive, allowing the tip of the ulnar head to protrude into the radiocarpal space. The degree of ulnar head perforation of the TFCC may be related to the amount of positive ulnar variance. One or several TFCC pedicled flaps can be observed in "DD" TFCC tears. Chondral erosive changes of the ulnar proximal lunate facet and/or ulnar head tip with or without secondary attritional lunotriquetral dissociation may be observed depending on the severity of the ulnocarpal impingement syndrome. The treatment of "DD" TFCC lesions whatever their stage and arthroscopic aspect consists of tears/flaps debridement and ulnar

decompression (arthroscopic wafer procedure or ulnar shortening osteotomy).

- "DT" tears (previously Palmer class 1A) are most often sagittal slit tears located 2 to 3 mm medial to the radial insertion of the TFCC.⁴ They are commonly encountered during arthroscopy following distal radius fractures.³ They usually do not require any specific treatment. Two flap variants of "DT" tears have been recently described:

Distal horizontal flap "DT" tears were described by del Piñal² and mentioned by other authors.^{4,17,18} They are well visualized and debrided through 3-4 and 6R radiocarpal portals.

Proximal horizontal flap "DT" tears⁶ with potential interposition into the DRUJ joint space may be observed. They require a combination of radiocarpal and DRUJ arthroscopy for flap resection.

Reins (R) Triangular Fibrocartilage Complex Traumatic Tears

All "R" tears are potentially DRUJ destabilizing injuries. "R1" tears are located at the ulnar side of the TFCC, whereas "R2" tears are located at its radial side.

There is increasing evidence^{3,19} that the deep (proximal) and superficial (distal) dorsal and palmar radioulnar ligaments are intimately merged so that they act simultaneously during forearm rotation to stabilize the DRUJ. The well-vascularized²⁰ deep and superficial dorsal radioulnar ligaments are taut in pronation.^{3,19} By contrast, the deep and superficial volar radioulnar ligaments are taut in pronation.

During wrist arthroscopy from the 3-4 portal, the deep proximal part of the "R" TFCC component (deep dorsal and palmar radioulnar ligaments) inserting into the foveal area cannot be readily seen. DRUJ wet or dry arthroscopy has recently been improved and can reliably show the foveal insertion of the deep radioulnar ligaments.^{4,21-23}

At the ulnar side of the reins, although it has been recently challenged²⁴ most authors^{21,25} rely on the combination of a clinical DRUJ instability and positive hook test²⁶ to indirectly assess the competence of the deep part of the TFCC reins.

At the radial side of the reins, probing is used to assess the radial insertions of the dorsal and volar radioulnar ligaments on each side of the disc.

- The classic foveal avulsion of the TFCC "reins" (i.e., the deep proximal dorsal and palmar radioulnar ligaments) or Atzei class 2 TFCC lesion is a well-recognized injury^{5,21,25} typically resulting from forceful pronation or supination. It is called "R1" TFCC tear (►Fig. 1). "R1" TFCC tear requires an anchor or transosseous reinsertion. The use of DRUJ arthroscopy in addition to radiocarpal arthroscopy may be necessary.^{21,24,25}
- Proximal ulnar styloid fractures occasionally include the deep foveal insertion of the TFCC. This is called "R1 bony" TFCC tears or "R1 ulnar styloid fracture with foveal avulsion." Ulnar styloid fixation may be required to fix both the fracture and the foveal avulsion. However, ulnar styloid fracture may also occur following an avulsion of




Fig. 1 Arthroscopically based classification of TFCC Disorders: the fresh or chronic, reparable or nonreparable nature of TFCC tears are considered as separate parameters (see text).

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Table 1 Correspondence between our arthroscopically based classification of TFCC disorders and Palmer's classification of TFCC disorders

Palmer's TFCC disorders classification	New arthroscopic TFCC disorders classification
	First, consider ulnar variance and Tolat's type of DRUJ sigmoid notch
	D: Disc tears
2A-B-C-D-E	DD: Disc degenerative central tear (wear / perforation / massive with/without flap)
1A	DT: Disc traumatic sagittal slit tear
	DT proximal horizontal flap variant ^a (Arimitsu 2021)
	DT distal horizontal flap variant (del Piñal 2012)
	R: Reins traumatic tears
1B foveal avulsion	R1: Complete foveal avulsion ^a (Nakamura 2011- Atzei Class 2 - 2009)
1B with ulnar styloid fracture	R1 Bony: Ulnar styloid fracture with foveal avulsion
	R1 Deep: Hidden foveal avulsion ^a (Atzei Class 3 - 2009)
	R2/R2 Bony: Dorsal rim/bony dorsal rim avulsion (Nakamura - 2012)
	R3/R3 Bony: Volar rim ^a /bony volar rim avulsion (Nakamura - 2012)
1D	R4: Complete radial rim avulsion (including pre-1D injury variant [Luchetti - 2022])
	W: Wall traumatic tears
	W1: Dorsal peripheral tear (Fukuoka Nakamura - 2020)
	W2: Ulnar peripheral tear (Atzei Class 1 - Abe 2016 - Fukuoka Nakamura - 2020)
	W3: Ulnocarpal detachment syndrome (Nishikawa - 2012)
	W4: Volar peripheral tear (del Piñal - 2012)
	W5: Ulnotriquetral split tear (Tay Kakar Berger - 1997)
1C	W6: Ulnocarpal volar avulsion
	Combined tears
	Example DD + R1 (del Piñal - Nakamura - 2012)

Abbreviations: DRUJ, distal radial ulnar joint; TFCC, triangular fibrocartilage complex.

^aUsefulness of DRUJ Arthroscopy.

the foveal insertion of the reins, producing a "floating styloid."⁷ This condition does not benefit from ulnar styloid fixation to restore DRUJ stability.

- Isolated hidden avulsion of the foveal insertion of the proximal radioulnar ligament while its superficial counterpart has remained intact (Atzei class 3) is called "R1 deep" TFCC tear. The treatment may be the same as for "R1" lesions. The use of DRUJ arthroscopy in addition to radiocarpal arthroscopy is necessary.^{1,4,21}
- Avulsion of the radial dorsal insertion of the reins¹¹ is called "R2" or "R2 dorsal rim avulsion" TFCC lesion. Fixation with anchors is recommended¹¹ to avoid DRUJ dorsal subluxation. When there is a displaced bony avulsion as in some intra-articular distal radius fractures, the injury is called "R2 bony." It requires specific osteosynthesis to avoid DRUJ dorsal subluxation.
- Avulsion of the radial volar insertion of the reins¹¹ is called "R3" or "R3 volar rim avulsion" TFCC lesion. Fixation with anchors is recommended¹¹ to avoid DRUJ volar subluxation. When there is a displaced bony avulsion as in some intra-articular distal radius fractures, the injury is called "R3 bony." It requires specific osteosynthesis to avoid DRUJ volar subluxation.
- "R4" TFCC tear is a complete traumatic avulsion of the TFCC from its radial attachment (Palmer's class 1D). This severe high-energy injury is exceedingly rare.¹¹ It may be observed in conjunction with radiocarpal dislocation. A less severe "pre-1D" variant of Palmer's classification has recently been described by Luchetti.¹⁰

Wall (W) Triangular Fibrocartilage Complex Traumatic Tears

During wrist arthroscopy from the 3–4 portal, what we have described as the TFCC wall "W"³ is the radially concave vertical well-vascularized²⁰ fibrous layer raising from the reins at the periphery of the disc and inserting on the medial aspect of the proximal carpal row (i.e., on the ulnar part of the lunate and triquetrum).

- Dorsal peripheral TFCC tears^{27,28} which are typically painful but not (or little) DRUJ destabilizing are called "W1." These TFCC tears represent a laceration of the junction of the dorsal part of the reins to the dorsal wall. The suction test may be a useful arthroscopic maneuver to arthroscopically confirm "W1" TFCC tears.²⁹ "W1" TFCC tears are amenable to capsular repair.²⁸

- Ulnar peripheral TFCC tears³⁰ which are tears of the ulnar part of the capsule merging with the ulnar termination of the converging superficial part of the reins are called “W2.” These TFCC tears are amenable to capsular repair.
- Ulnocarpal detachment syndrome^{12,31,32} is a lesion of the dorsal carpal attachment of the wall to the triquetrum merging with the radiotriquetral ligament which is called “W3.”
- Volar Peripheral TFCC tears^{2,33} are called “W4” TFCC tears. These TFCC tears represent a laceration of the junction of the volar part of the reins to the volar wall. They are amenable to capsular fixation.
- Longitudinal split tears of the ulno-triquetral ligament^{9,34} are called “W5.” It may be an isolated TFCC lesion but combined regional pathology (other TFCC tear, ECU pathology, and ulnocarpal impingement) was associated in 17% in a recent series.³⁵ “W5” TFCC tears are amenable to capsular repair.⁹
- Ulnocarpal volar TFCC avulsion (former Palmer’s Class 1C) is an exceedingly rare and severe lesion called “W6.” They may be amenable to reattachment to the carpal bones.

Combined Triangular Fibrocartilage Complex Lesions

In the past 10 years, many authors highlighted the fact that a combination of different types of TFCC tears was sometimes observed. Del Piñal was the first author to formally describe several cases of combined degenerative and traumatic TFCC tears,² that is, for example, combined “DD” and “R1” (if DRUJ destabilizing; ► **Figs. 2–3**) or “DD” and “W2.” Other combined TFCC disorders may be identified in the future and easily added to our arthroscopic TFCC classification.

Discussion

Wrist arthroscopy has proven to be a prerequisite for the contemporary diagnosis and treatment of TFCC disorders. However, since the landmark Palmer’s classification of TFCC disorders and despite many recent papers about newly recognized TFCC injuries, no comprehensive arthroscopic classification of TFCC lesions has been published.

We recently proposed an arthroscopic description of the box-like TFCC structure as seen from the 3–4 radiocarpal portal including functional, vascular, and thus therapeutic aspects along with a preliminary attempt at stratifying TFCC tears.³ We now propose a final version of the classification including all the known disorders of the TFCC to the best of our knowledge and considering the need for combining radiocarpal and DRUJ arthroscopy.

The format of the comprehensive above-described arthroscopic classification is simple, user-friendly, and updatable with future new descriptions. Patient’s specific acute or chronic pattern of the traumatic lesions must be mentioned as well as the reparability or need for reconstruction of the injury.

Palmer’s landmark paper was the first attempt at producing a comprehensive 3-D classification of TFCC lesions including wrist arthroscopy as one of its criteria.⁵ Although it is still widely used, many authors have shown that it does not



Fig. 2 Example of combined “« DD » and « R1 » destabilizing lesion as described by del Piñal and Nakamura in reference.⁸

include some newly described injuries^{2,18,36} or allow identifying combined TFCC lesions⁸ as the combination of a degenerative and traumatic injury for example. Moreover, Palmer’s Class 1 lesions described different locations of traumatic TFCC injuries, whereas the Class 2 lesions described the natural history of a degenerative central tear due to ulnocarpal impingement.⁵

Numerous papers focusing on the analysis of subtypes of Palmer’s Class 1B lesions have been published.^{1,2,4} Some TFCC lesions could not be found in Palmer’s landmark paper.³⁶

Using 3-T MRI, Zhan et al^{17,18} reviewed 86 wrists with TFCC injuries. A total of 17 (20%) could not be classified according to Palmer’s classification.

Using arthroscopic diagnosis and treatment, Estrella et al³⁶ analyzed 35 wrists with traumatic dorsal tears of the TFCC. They found that 18 (51%) could not be classified according to Palmer’s classification.

Atzei and Luchetti^{1,7} included reparability as a key criterion in their classification of foveal TFCC tears to advise repair versus reconstruction. It is the author’s opinion that this key criterion should be used as an additional parameter for some specific types of “R” and “W” TFCC disorders. The acute or chronic nature of the TFCC tears should be included as an additional parameter. Usually, the 45th day is accepted as the landmark delay after which ligamentous injury is considered to be chronic.

The limitations of existing classifications of TFCC disorders have led Schmitt et al¹⁴ to recently propose a new classification system, the “CUP” classification (“C” for central, “U” for ulnar, and “P” for peripheral). This classification is based on a classic 3-D anatomical description of the TFCC and includes a “combined lesions” category as recommended by previous authors.^{3,8} However, this new classification is not




Fig. 3 Clinical example of combined « DD » and « R1 » TFCC tears. (A) Ulna + 2 mm variance in a 39 year-old lady (high-level volleyball player during 10 years). (B) She presented with both positive ulnocarpal stress test and positive painful DRUJ drawer test 1 year after a twisting and hyperflexion wrist trauma. (C) Obvious disc degenerative “DD” central tear as seen from 3–4 portal. (D) Positive arthroscopic hook test showing an additional “R1” tear.

easy to memorize and includes measurements in millimeters which may be difficult to reproduce.

We believe that our new arthroscopic TFCC classification is the simplest and most comprehensive current update about TFCC disorders.

Ethical Review Committee Statement

Not Applicable.

Conflicts of Interest

None declared.

References

- Atzei A, Luchetti R, Garagnani L. Classification of ulnar TFCC tears. A treatment algorithm for Palmer type IB tears. *J Hand Surg Eur* 2017;42E(04):405–414
- Del Piñal F. The 1 B constellation: an attempt to classify Palmer 1B Class. In: Del Piñal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:87–97
- Herzberg G, Burnier M, Nakamura T. Arthroscopic anatomy of the TFCC with relevance to function and pathology. *J Wrist Surg* 2021; 10(06):558–564
- Nakamura T, Takagi T. Differentiated approaches to treat lesions of the TFCC based on new arthroscopic classification. *Handchir Mikrochir Plast Chir* 2022;54(05):389–398
- Palmer AK. Triangular fibrocartilage complex lesions: a classification. *J Hand Surg Am* 1989;14(04):594–606
- Arimitsu S, Masatomi T, Shigi A, Yukioka C, Moritomo H. Proximal horizontal flap tears of TFCC diagnosed by computed tomography arthrography: six case series. *J Wrist Surg* 2021;10 (01):36–41
- Atzei A, Luchetti R. Foveal TFCC tear classification and treatment. *Hand Clin* 2011;27(03):263–272
- Del Piñal F, Nakamura T. Coronal tears of the TFCC. In: Del Piñal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:199–211
- Gillis J, Kakar S. Ulnar triquetral Ligament Split tears. In: Kakar S, Yao J, eds. *Ulnar-Sided Wrist Disorders*. Chicago: ASSH; 2022
- Luchetti R, Cozzolino R, Marcovici LL, Atzei A. The pre-1D lesion of the TFCC—a new variant of the Palmer 1D class. *Arthrosc Tech* 2022;11(08):e1435–e1440

- 11 Nakamura T. Radial side tear of the TFCC. In: Del Piñal F, Mathoulin C, Luchetti R, eds. *Arthroscopic Management of Distal Radius Fractures*. Springer 2010:89
- 12 Nishikawa S, Toh S, Miura H. The carpal detachment injury of the TFCC. *J Hand Surg Brit Eur Vol* 2002;27B:86–89
- 13 Shapiro LM, Richard MJ. Traumatic distal ulnar and sigmoid notch injuries. In: Kakar S, Yao J, eds. *Ulnar-Sided Wrist Disorders*. Chicago: ASSH; 2022:235–249
- 14 Schmitt R, Grunz JP, Langer M. Triangular fibrocartilage complex injuries—limitations of the current classification systems and the proposed new ‘CUP’ classification. *J Hand Surg Eur Vol* 2023;48(01):60–66
- 15 Tomaino MM. The importance of the pronated grip x-ray view in evaluating ulnar variance. *J Hand Surg Am* 2000;25(02):352–357
- 16 Tolat AR, Stanley JK, Trail IA. A cadaveric study of the anatomy and stability of the distal radioulnar joint in the coronal and transverse planes. *J Hand Surg [Br]* 1996;21(05):587–594
- 17 Zhan H, Zhang H, Bai R, et al. High-resolution 3-T MRI of the triangular fibrocartilage complex in the wrist: injury pattern and MR features. *Skeletal Radiol* 2017;46(12):1695–1706
- 18 Zhan H, Bai R, Qian Z, Yang Y, Zhang H, Yin Y. Traumatic injury of the triangular fibrocartilage complex (TFCC)—a refinement to the Palmer classification by using high-resolution 3-T MRI. *Skeletal Radiol* 2020;49(10):1567–1579
- 19 Crowe MM, Martin JT, Grier AJ, Spritzer CE, Richard MJ, Ruch DS. In vivo mechanical function of the DRU ligaments during rotation of the wrist. *J Hand Surg Am* 2020;45(11):1012–1021
- 20 Bednar MS. The microvasculature of the TFCC: clinical significance. *JHSA* 1991;16(06):1101
- 21 Atzei A, Luchetti R, Senesi L, Marcovici LL. Arthroscopic management of foveal tears of the TFCC with DRUJ instability. In: Kakar S, Yao J, eds. *Ulnar-Sided Wrist Disorders*. Chicago: ASSH; 2022:111–124
- 22 Burnier M, Herzberg G, Luchetti R, Del Piñal F, Kakar S. Dry wrist arthroscopy for ulnar sided wrist disorders. *J Hand Surg Am* 2021;46(02):133–141
- 23 Nakamura T, Sato K, Okazaki M, Toyama Y, Ikegami H. Repair of foveal detachment of the triangular fibrocartilage complex: open and arthroscopic transosseous techniques. *Hand Clin* 2011;27(03):281–290
- 24 Ecker J, Andrijich C. Dry arthroscopy distal radioulnar joint and foveal insertion: surgical technique. *J Wrist Surg* 2021;11(01):2–5
- 25 Nakamura T. DRUJ instability: arthroscopic repair of the detached TFCC to the fovea of the ulna. In: Del Pinal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:137–145
- 26 Atzei A, Luchetti R, Carletti D, Marcovici LL, Cazzoletti L, Barbon S. The hook test is more accurate than the trampoline test to detect foveal tears of the triangular fibrocartilage complex of the wrist. *Arthroscopy* 2021;37(06):1800–1807
- 27 Abe Y, Moriya A, Tominaga Y, Yoshida K. Dorsal tear of triangular fibrocartilage complex: clinical features and treatment. *J Wrist Surg* 2016;5(01):42–46
- 28 Haerle M. Treatment of stable peripheral TFCC tears. In: Del Pinal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:111–120
- 29 Greene RM, Kakar S. The suction test: a novel technique to identify and verify successful repair of peripheral triangular fibrocartilage complex tears. *J Wrist Surg* 2017;6(04):334–335
- 30 Fukuoka M, Nakamura T. Effects of peripheral tears of the triangular fibrocartilage complex on distal radioulnar joint stability: a biomechanical study. *J Orthop Sci* 2021;26(06):1008–1013
- 31 Nishikawa S, Toh S. Anatomical study of the carpal attachment of the triangular fibrocartilage complex. *J Bone Joint Surg Br* 2002;84(07):1062–1065
- 32 Nishikawa S, Toh S. The carpal detachment syndrome. In: Del Pinal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:161–168
- 33 Cerezal L. Ulnar imaging for arthroscopic surgeon. In: Del Pinal F, Mathoulin C, Nakamura T, eds. *Arthroscopic Management of Ulnar Wrist Pain*. New York: Springer; 2012:61–85
- 34 Tay SC, Berger RA, Parker WL. Longitudinal split tears of the ulnotriquetral ligament. *Hand Clin* 2010;26(04):495–501
- 35 Clark NJ, Berger RA, Kakar S. Outcomes of UT split tears repair: a report of 96 patients. *J Hand Surg (Eur Vol)* 2019;44E(10):1036
- 36 Estrella EP, Ho PC, Tse WL. Arthroscopic repair of triangular fibrocartilage complex tears. *Arthroscopy* 2007;23(07):729–737, 737.e1