



# CT-Guided Percutaneous Thrombin Injection of Posttraumatic Aortic Branch Pseudoaneurysm

Aron Michael Devane<sup>1</sup> Christine Marie Gilligan Schammel<sup>2</sup>

<sup>1</sup>Department of Radiology, Prisma Health Upstate, Greenville, South Carolina, United States

<sup>2</sup>Pathology Associates, Greenville, South Carolina, United States

Address for correspondence Christine Marie Gilligan Schammel, PhD, Department of Pathology, USCSOMG, 8 Memorial Medical Ct., Greenville, SC 29605, United States (e-mail: Christine.schammel@prismahealth.org).

J Clin Interv Radiol ISVIR 2024;8:49–52.

## Abstract

A pseudoaneurysm of an aortic branch artery is a potentially life-threatening uncommon occurrence, which may result in retroperitoneal hemorrhage. Imaging-guided percutaneous thrombin injection has a high technical success and effectiveness rate for treating pseudoaneurysms of the femoral, iliac, and popliteal arteries. We present a case of a retroperitoneal hemorrhage in a patient with a periaortic branch pseudoaneurysm, potentially from an avulsed lumbar artery. Anatomy prevented fixation with a covered stent; the comorbidities eliminated safe surgical repair, and endovascular embolization was not an option due to the avulsion. Computed tomography (CT) fluoroscopy-guided thrombin injection provided a safe, successful embolization with no postprocedural complications.

## Keywords

- ▶ aortic branch pseudoaneurysm
- ▶ lumbar artery pseudoaneurysm
- ▶ CT-guided thrombin injection

## Introduction

Posttraumatic aortic branch pseudoaneurysms (PSAs) are uncommon<sup>1</sup> and are typically associated with high-energy, blunt force or penetrating trauma with injury to intra-abdominal structures, resulting in a potentially life-threatening retroperitoneal hemorrhage<sup>1</sup> The risk of rupture requires prompt treatment as no reliable predictive symptoms exist.<sup>2</sup>

The recommended treatment for PSA is catheter angiography with endovascular embolization or endovascular stent graft exclusion. The most common embolic materials are coils; other embolic materials, such as polyvinyl alcohol (PVA), acrylic microspheres, gelatin sponge, and n-BCA are less commonly used.<sup>2</sup> Imaging-guided percutaneous thrombin injection has been widely accepted<sup>3</sup> as a treatment of femoral artery PSAs, with a suggestion for the broader use of thrombin for PSAs across other vessels.<sup>4</sup>

Here, we present a case of a retroperitoneal hemorrhage in an elderly patient presenting with a spinal fracture after a ground-level fall successfully treated with a computed tomography (CT) fluoroscopy-guided thrombin injection. Following institutional review board (IRB) approval and upon receiving publication consent, this report was completed.

## Case Report

An 81-year-old woman with a history of breast cancer presented to the emergency department with abdominal and back pain after a fall from a standing position. CT examination demonstrated a right periaortic PSA at the T12/L1 level (2.1 × 2.1 × 2.7 cm; ▶ **Fig. 1**) and a compression fracture of the L1 vertebral body. No apparent communication of the PSA with any branch of the abdominal aorta

article published online  
September 3, 2023

DOI <https://doi.org/10.1055/s-0043-1772493>.  
ISSN 2457-0214.

© 2023. Indian Society of Vascular and Interventional Radiology. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



**Fig. 1** Contrast-enhanced computed tomography (CT) axial plane demonstrating a pseudoaneurysm (*white arrow*) adjacent to the abdominal aorta.

was noted, suggesting an avulsed origin of a lumbar artery or the right subcostal artery given the T12/L1 location, eliminating the potential for catheter angiography and coil embolization. The adjacent aortic branches, such as the celiac axis, made fixation with a covered stent technically challenging and, as the spinal artery arises from this location, induced an increased risk of paraplegia. The location of the PSA and the patient's comorbidities made surgical repair high risk.

Utilizing CT fluoroscopy guidance, a 10-cm 19-gauge needle (Argon Medical, Plano, TX) was advanced into the PSA with a prone right paravertebral approach (►**Fig. 2**). Pulsatile arterial blood emanated from the needle spontaneously. Reconstituted thrombin (0.5 mL; 500 units; Pfizer, New York, NY) was administered with cessation of blood from the needle. No blood could be manually aspirated with a 5-mL syringe after a 10- to 15-second wait period. The needle was removed. CT angiography (CTA) demonstrated a lack of contrast opacification of the PSA in the arterial (►**Fig. 3**) and the delayed phase of intravenous contrast enhancement. No procedural-related complica-



**Fig. 2** Computed tomography (CT) fluoroscopy image demonstrating the right paravertebral approach of a 19-gauge needle into the pseudoaneurysm.



**Fig. 3** Axial computed tomography (CT) angiography (arterial phase) demonstrating resolution of contrast enhancement of the pseudoaneurysm after thrombin injection (*white arrow*).

tions were observed. At 3.5 months postinjection, the patient was asymptomatic, and a CTA illustrated a successful embolization.

## Discussion

While aortic branch PSAs are typically associated with blunt abdominal trauma with a spinal injury,<sup>1</sup> in an elderly patient, this should be considered a source of hemorrhage, even from a ground-level fall. On imaging, aortic branch PSAs present as an enhanced density close to the psoas major muscle, the lumen enhancing similar to that of the aorta.<sup>1,5</sup> A literature review of 26 cases (►**Table 1**) noted that all reports but two used embolization. Our report notes the limitations of the anatomy and lack of PSA communication with any abdominal aortic branch, suggesting an avulsed origin. This makes it impossible to catheterize the visceral aortic branch PSA endovascularly, and coil embolization was not feasible. The use of other embolic agents, such as glue or Onyx, would have required visualization, which was not a possibility in our case. Also, these embolic agents typically require larger volumes, risking reflux; glue would have been an unsafe option due to the possibility of attaching the needle to the PSA, risking a tear. While direct percutaneous injection of thrombin has been the preferred treatment for postcatheterization femoral artery PSAs,<sup>2,3</sup> it has had limited use in aortic branch PSA, except when embolization is not an option.<sup>2,5</sup> Given these data, thrombin appeared to be the safest option for this patient. A small amount of thrombin was administered initially to prevent reflux into the abdominal aorta, which could have resulted in visceral and lower extremity thrombus. Preparations were made for repeated small injections with repeated contrast-enhanced CT imaging if the first dose was ineffective; fortunately, this was unnecessary.

Given the favorable outcomes at other sites and that aortic PSAs are life-threatening, thrombin is a viable option for this site.

### Compliance with Ethical Standards

All procedures performed in studies involving human participants were in accordance with the ethical standards of

**Table 1** Survey of reported lumbar artery pseudoaneurysms

Study	Age	Cause	Presentation	Resolution
Ikubo et al <sup>6</sup>	62	Blunt trauma	Abdominal pain/hypotension	Embolization
Siablis et al <sup>7</sup>	45	Fall from 6 m	“Agonizing pain”	Embolization
Chan and Korivi <sup>8</sup>	24	Gunshot	Low back/flank pain	Embolization
Kessel et al <sup>9</sup>	21	Penetrating abdominal trauma	Missed during laparotomy	Embolization
Lee et al <sup>10</sup>	47	Fracture of lumbar spine	Hypotensive shock	Embolization
Domenicucci et al <sup>11</sup>	23	Thoracolumbar vertebral dislocation	Anemia/hemorrhage	Embolization
Ramsay and Marshall <sup>12,a</sup>	23	Renal biopsy		Thrombin
Sharma et al <sup>5,a</sup>	47	Spontaneous	Abdominal pain	Thrombin
Tomescot et al <sup>13</sup>	79	Osteomyelitis	Acute back pain	Embolization
Oh et al <sup>14</sup>	55	Lumbar posterolateral fusion	Retroperitoneal Hemorrhage	Embolization
Counihan et al <sup>15</sup>	18	Knife stab	Bleeding	Embolization
Suresh et al <sup>16</sup>	60	Postpercutaneous coronary intervention	Back ache/hypotensive	Embolization
Tsekouras et al <sup>17</sup>	64	Inferior vena cava filter	Abdominal pain	Embolization
Djuimo et al <sup>18</sup>	21	Nephrostomy removal	Bleeding on removal	Embolization
Giordano et al <sup>19</sup>	73	Cryoablation/vertebroplasty lumbar met	Abdominal pain	Embolization
Vashisht et al <sup>4</sup>	32	Lumbar/spine fracture from fall	Hypotensive shock	Symptom management
Salam and Khandwala <sup>20</sup>	71	Renal biopsy	Hemodynamic instability	Embolization
Panagiotopoulos et al <sup>21</sup>	39	Lumbar discectomy	Abdominal pain	Embolization
Latka et al <sup>22</sup>	58	Postlumbar fusion	Abdominal pain	Embolization
Méndez et al <sup>23</sup>	72	Postlumbar fusion	Monoplegia abdominal pain	Embolization
Eissa et al <sup>24</sup>	30	Fall from 2 m/postlumbar fusion	Back pain	Embolization
Ruffilli et al <sup>25</sup>	53	Postlumbar fusion	Back pain	Embolization
Keerthivasan et al <sup>26</sup>	80	Postlumbar fusion	Leg pain	Embolization
Kim et al <sup>27</sup>	73	Postlumbar fusion	Swelling of leg	Stent graft
Nam et al <sup>28</sup>	62	Postvertebral augmentation	Back pain	Embolization
Nam et al <sup>28</sup>	88	Postvertebral augmentation	Back pain	No intervention

Note: The table provides a survey of reported lumbar artery pseudoaneurysms, as this was the most likely location in our patient.

<sup>a</sup>The only report of thrombin injections for an aortic branch (lumbar artery) pseudoaneurysm.

the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

This study has obtained IRB approval from (Prisma Health), and the need for informed consent was waived. Dr. Devane is a paid speaker for Johnson and Johnson and TriSalus Life Sciences and is a consultant with Boston Scientific and Guerbet.

#### Funding

None.

#### Conflict of Interest

None declared.

#### References

- Shigematsu Y, Kudoh K, Nakasone Y, Fujisaki T, Uemura S, Yamashita Y. Nontraumatic rupture of lumbar artery causing an intra-vertebral body pseudoaneurysm: treatment by transcatheter embolization. *Cardiovasc Intervent Radiol* 2006;29(05):870–874
- Thomson B, Patel V, Moser S, Diamantopoulos A. Successful Treatment of a left gastric artery pseudoaneurysm by image guided percutaneous thrombin injection. *Vasc Endovascular Surg* 2023;57(03):306–310
- Eheli WL, Bozdogan E, Janas G, et al. Imaging-guided percutaneous thrombin injection for the treatment of iatrogenic femoral artery pseudoaneurysms. *Abdom Radiol (NY)* 2019;44(03):1120–1126
- Vashisht S, Bachhal V, Kumar V, Kumar D. Lumbar artery pseudoaneurysm following blunt trauma: A case report and literature review. *Trauma Case Rep* 2019;23:100232

- 5 Sharma D, Elmore JRE, Nadal LLS, Sheldon DGI, Franklin DPI. Treatment of spontaneous lumbar artery pseudoaneurysm by computed tomography: guided thrombin injection. *Vasc Endovascular Surg* 2008;42(03):284–288
- 6 Ikubo A, Komura M, Matoba N, et al. Lumbar artery pseudoaneurysm: an unusual cause of a retroperitoneal hematoma: report of a case. *Surg Today* 1993;23(07):635–638
- 7 Siablis D, Panagopoulos C, Karamessini M, et al. Delayed diagnosis of a false aneurysm after lumbar arterial injury: treatment with endovascular embolization: a case report. *Spine* 2003;28(04):E71–E73
- 8 Chan KT, Korivi N. Lumbar artery pseudoaneurysm in traumatic spinal cord injury: a case report. *Arch Phys Med Rehabil* 2003;84(03):455–457
- 9 Kessel BJ, Habib FA, Thompson B, Bajayo DE, Shatz D. Lumbar artery pseudoaneurysm: an unusual complication of penetrating abdominal trauma. *Eur J Trauma* 2004;30:187–190
- 10 Lee JS, Kim CW, Suh KT. Lumbar artery injury combined with a transverse process fracture of the lumbar spine presenting with hypovolemic shock after a fall: a case report. *J Korean Orthop Assoc* 2008;43:400–403
- 11 Domenicucci M, Ramieri A, Lenzi J, Fontana E, Martini S. Pseudoaneurysm of a lumbar artery after flexion-distraction injury of the thoraco-lumbar spine and surgical realignment: rupture treated by endovascular embolization. *Spine* 2008;33(03):E81–E84
- 12 Ramsay DW, Marshall M. Lumbar artery pseudoaneurysm following renal biopsy: treatment with ultrasound-guided thrombin injection. *Australas Radiol* 2002;46(02):201–203
- 13 Tomescot A, Dallaudiere B, Zurlinden O, Manelfe J. Lumbar artery pseudoaneurysm as a late complication of osteomyelitis with vertebral body destruction. *J Vasc Surg* 2013;58(04):1084–1087
- 14 Oh YM, Choi HY, Eun JP. Delayed retroperitoneal hemorrhage due to lumbar artery pseudoaneurysm after lumbar posterolateral fusion. *J Korean Neurosurg Soc* 2013;54(04):344–346
- 15 Counihan M, Pontell ME, Selvan B, Trebelev A, Nunez A. Delayed presentation of a lumbar artery pseudoaneurysm resulting from isolated penetrating trauma. *J Surg Case Rep* 2015;2015(07):rjv083
- 16 Suresh A, Seshagiri Rao D, Rammurti S, Srinivas B, Sivaprasad A. Coil closure of lumbar artery pseudoaneurysm: an unusual complication associated with anomalous left circumflex artery primary percutaneous coronary intervention. *Chronic Dis Transl Med* 2015;1(01):55–58
- 17 Tsekouras N, Whalen RC, Comerota AJ. Lumbar artery pseudoaneurysm in a patient with inferior vena cava filter and history of strenuous physical exercise. *J Vasc Surg* 2015;61(03):796–799
- 18 Djuimo M, Aubé M, Beland M, Jeldres C, Carmel M, Benko A. Lumbar artery pseudoaneurysm: a complication of percutaneous nephrostomy. *Urol Case Rep* 2017;13:66–68
- 19 Giordano AV, Arrigoni F, Bruno F, et al. Interventional radiology management of a ruptured lumbar artery pseudoaneurysm after cryoablation and vertebroplasty of a lumbar metastasis. *Cardiovasc Intervent Radiol* 2017;40(05):776–779
- 20 Salam B, Khandwala K. Lumbar artery pseudoaneurysm following renal biopsy. *Cureus* 2018;10(05):e2634
- 21 Panagiotopoulos K, Gazzeri R, Bruni A, Agrillo U. Pseudoaneurysm of a segmental lumbar artery following a full-endoscopic transforaminal lumbar discectomy: a rare approach-related complication. *Acta Neurochir (Wien)* 2019;161(05):907–910
- 22 Latka K, Zurawel R, Maj B, Olbrycht T, Chowanec J, Latka D. Iatrogenic lumbar artery pseudoaneurysm after lumbar transpedicular fixation: case report. *SAGE Open Med Case Rep* 2019;7: X19835344
- 23 Méndez JC, Fandino E, Bermúdez-Coronel I, Prieto MA, Blázquez J. Endovascular occlusion of iatrogenic lumbar artery pseudoaneurysm using liquid embolic agent: case report. *Trauma Case Rep* 2020;29:100352
- 24 Eissa AT, Shahbaz A, Alhelal F, et al. Postoperative paralumbar pseudoaneurysm: a case report. *J Neurol Surg Rep* 2020;81(02): e33–e37
- 25 Ruffilli A, Barile F, Fiore M, Pasini S, Facchini G, Faldini C. Occurrence of L4 lumbar artery pseudoaneurysm after posterior reduction and L4–L5 fusion for grade I anterolisthesis: a case report. *JBJS Case Connect* 2020;10(04):00117
- 26 Keerthivasan P, Anupama NV, Kanna RM, Shetty AP, Rajasekaran S. Lumbar artery pseudoaneurysm: a rare case of delayed onset incomplete cauda equina syndrome following transforaminal lumbar interbody fusion. *Eur Spine J* 2020;29(Suppl 2):156–161
- 27 Kim C, Hwang D, Yun WS. Endovascular repair of an ilio-iliac arteriovenous fistula with pseudoaneurysm after lumbar disc surgery: a case report. *Vasc Spec Int* 2021;37:30
- 28 Nam D, Johnson MH, Mojibian H, Hebert RM, Cornman-Homonoff J. Development of paravertebral pseudoaneurysms following vertebral augmentation: a report of two cases. *Clin Imaging* 2022;90:39–43